



Universiteit
Leiden
The Netherlands

Comparative visual analytics for multi-modal single-cell and spatial omics data

Basu, S.

Citation

Basu, S. (2026, September 15). *Comparative visual analytics for multi-modal single-cell and spatial omics data*. Retrieved from <https://hdl.handle.net/1887/4310004>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4310004>

Note: To cite this publication please use the final published version (if applicable).

SUPPLEMENTARY INFORMATION

S1: Supplementary for Cytosplore Gradient Surfer ([Chapter 4](#))

S1.1: Technical validation

We first evaluated robustness using the MERFISH spatial transcriptomics profile of a mouse visual cortex slice. To maximize the number of genes available for analysis and obtain stronger expression signals, we included only the imputed genes from the MERFISH dataset. We ran Cytosplore Gradient Surfer on the original imputed MERFISH dataset (Dataset Technical 1, DT1; 2,150 cells) as the reference. We then generated a set of derivative datasets (collectively referred to as DT2 variants) from DT1 by applying individual transformations, including rotation, reduced spatial resolution, normalization, or dimensionality reduction, and evaluated each variant relative to DT1.

S1.1.1: Robustness against dataset rotation

We rotated the spatial coordinates in DT2 by 45°, 90°, 180°, and 270° (keeping DT1 fixed) and recomputed gene–position correlations. The same top-ranking gradient genes emerged despite the geometric transformation. Layer-specific marker genes that showed a clear laminar gradient in the unrotated data remained among the highest-correlated genes after rotation, with their spatial expression patterns correspondingly rotated. This result indicates that the method identifies consistent patterns even when one dataset is rotated, as expected from its design to detect stable gradients across spatial transformations.

S1.1.2: Robustness against cell subsampling

We assessed the robustness of the method under varying spatial cell densities by randomly subsampling cells from the MERFISH dataset in DT2, removing 10–90% of the cells (rows) and repeating the analysis. The method demonstrated strong stability across all levels of data reduction. Both Spearman’s and Kendall’s rank correlations remained high (>0.9) even after 90% data reduction, indicating excellent concordance in gene ranking. Analysis of gene rank changes revealed that most genes exhibited no or only minor rank changes, while moderate and major shifts were infrequent. Similarly, the top 50–100 ranked genes retained over 90% stability across all reduction levels. Although rank volatility increased progressively with greater data loss, the magnitude of change remained moderate, suggesting that the method

is resilient to sampling variability. The mean absolute rank shift was approximately 10 ranks out of 314 total genes at 90% reduction, further supporting the limited impact of subsampling. Statistical analyses confirmed strong rank consistency (Spearman and Wilcoxon signed-rank test, $p < 10^{-8}$ across all levels). Extreme ranks remained stable, and gene movements showed no directional bias under subsampling.


Collectively, these results indicate that the method maintains strong reproducibility and robustness under substantial spatial subsampling, ensuring that key gene rankings and biological interpretations remain consistent even with markedly reduced data density.

S1.1.3: Robustness against feature subsampling


To further assess robustness under varying feature dimensionality, we simulated a realistic scenario in which datasets share only a subset of genes. We first identified the top correlated genes (correlation coefficient > 0.5 ; $n = 41$) showing clear expression gradients across the dataset. Among these, the top 5 genes were retained for computation across all derivative datasets, while the remaining 36 genes were divided into two groups using two strategies: (i) random reduction, where genes were randomly split, and (ii) alternating reduction, where genes were assigned based on rank order.

Using these subsets, we generated four derivative datasets by removing the corresponding gene groups from the original dataset (DT1), resulting in two random-reduction and two alternating-reduction conditions. The analysis was repeated to evaluate whether gene rankings within each dataset's top 23 correlated genes (ranked by correlation strength), consisting of 5 universally retained genes and 18 subset-specific genes, were preserved under partial gene removal. In both random and alternating reduction settings, all genes showed high preservation of their relative rank positions within the top 23, with no major rank inversions observed.

Collectively, these results demonstrate that the method remains stable even under substantial reductions in feature number. The preservation of rank structure across all reduced-dimension datasets highlights its robustness and applicability to real-world scenarios where datasets may only partially overlap in gene coverage.



Supplementary Figure 1: Validation of conserved spatial gene expression gradients across primate species using a marmoset reference swipe. Spatial expression maps of representative gradient-associated genes identified by interactive line-swipe analysis across human, macaque, and marmoset brain slices (P7, P9, and P13, respectively), with the line swipe performed on the marmoset slice. In the figure, R denotes rank, P denotes positive expression rank, and N denotes negative expression rank. The results illustrate that gradients identified in marmoset correspond to conserved molecular patterns observed in other primate species.



Supplementary Figure 2: Validation of conserved spatial gene expression gradients across primate species using a macaque reference swipe. Spatial expression maps of representative gradient-associated genes identified by interactive line-swipe analysis across human, macaque, and marmoset brain slices (P7, P9, and P13, respectively), with the line swipe performed on the macaque slice. In the figure, R denotes rank, P denotes positive expression rank, and N denotes negative expression rank. The figure further supports the robustness of the approach in identifying conserved spatial gene expression gradients across primate species.

REFERENCES

- [1] B. Tasic, Z. Yao, L. T. Graybuck, et al. “Shared and distinct transcriptomic cell types across neocortical areas”. In: *Nature* 563.7729 (2018), pages 72–78.
- [2] A. Butler, P. Hoffman, P. Smibert, et al. “Integrating single-cell transcriptomic data across different conditions, technologies, and species”. In: *Nature Biotechnology* 36.5 (2018), pages 411–420.
- [3] L. Van der Maaten and G. Hinton. “Visualizing data using t-SNE.” In: *Journal of Machine Learning Research* 9.11 (2008), pages 2579–2605.
- [4] L. McInnes, J. Healy, and J. Melville. “Umap: Uniform manifold approximation and projection for dimension reduction”. In: *arXiv* (2018), pages 2008–02.
- [5] E. Becht, L. McInnes, J. Healy, et al. “Dimensionality reduction for visualizing single-cell data using UMAP”. In: *Nature Biotechnology* 37.1 (2019), pages 38–44.
- [6] J. Heer and B. Shneiderman. “Interactive dynamics for visual analysis: A taxonomy of tools that support the fluent and flexible use of visualizations”. In: *Queue* 10.2 (2012), pages 30–55.
- [7] F. A. Wolf, P. Angerer, and F. J. Theis. “SCANPY: large-scale single-cell gene expression data analysis”. In: *Genome Biology* 19.1 (2018), page 15.
- [8] A. Regev, S. A. Teichmann, E. S. Lander, et al. “The human cell atlas”. In: *eLife* 6 (2017), e27041.
- [9] I. Yanai and M. Lercher. “A hypothesis is a liability”. In: *Genome Biology* 21.1 (2020), page 231.
- [10] A. Maćkiewicz and W. Ratajczak. “Principal components analysis (PCA)”. In: *Computers & Geosciences* 19.3 (1993), pages 303–342.
- [11] D. J. Ardesch, L. H. Scholtens, L. Li, et al. “Evolutionary expansion of connectivity between multimodal association areas in the human brain compared with chimpanzees”. In: *Proc. Natl. Acad. Sci.* 116.14 (2019), pages 7101–7106.
- [12] T. E. Bakken, N. L. Jorstad, Q. Hu, et al. “Comparative cellular analysis of motor cortex in human, marmoset and mouse”. In: *Nature* 598.7879 (2021), pages 111–119.
- [13] N. L. Jorstad, J. H. Song, D. Exposito-Alonso, et al. “Comparative transcriptomics reveals human-specific cortical features”. In: *Science* 382.6667 (2023), ade9516.
- [14] Z. Dong and Y. Chen. “Transcriptomics: advances and approaches”. In: *Science China Life Sciences* 56 (2013), pages 960–967.
- [15] M. E. R. Shafer. “Cross-Species analysis of single-cell transcriptomic data”. In: *Front. Cell Dev. Biol.* 7 (2019), page 175.
- [16] A. Adil, V. Kumar, A. T. Jan, and M. Asger. “Single-cell transcriptomics: current methods and challenges in data acquisition and analysis”. In: *Frontiers in Neuroscience* 15 (2021), page 591122.

- [17] S. I. O'Donoghue, A.-C. Gavin, N. Gehlenborg, et al. "Visualizing biological data-now and in the future". In: *Nature Methods* 7.S3 (2010), S2–S4.
- [18] R. G. Briggs, O. Tanglay, N. B. Dadario, et al. "The unique fiber anatomy of middle temporal gyrus default mode connectivity". In: *Operative Neurosurgery* 21.1 (2021), E8.
- [19] J. Davey, H. E. Thompson, G. Hallam, et al. "Exploring the role of the posterior middle temporal gyrus in semantic cognition: Integration of anterior temporal lobe with executive processes". In: *NeuroImage* 137 (2016), pages 165–177.
- [20] S. Chen, Y. Chang, L. Li, et al. "Spatially resolved transcriptomics reveals genes associated with the vulnerability of middle temporal gyrus in Alzheimer's disease". In: *Acta Neuropathologica Communications* 10.1 (2022), pages 1–24.
- [21] R. D. Hodge, T. E. Bakken, J. A. Miller, et al. "Conserved cell types with divergent features in human versus mouse cortex". In: *Nature* 573.7772 (2019), pages 61–68.
- [22] S. Lu, D. Fürth, and J. Gillis. "Integrative analysis methods for spatial transcriptomics". In: *Nature Methods* 18.11 (2021), pages 1282–1283.
- [23] J. Qian, J. Liao, Z. Liu, et al. "Reconstruction of the cell pseudo-space from single-cell RNA sequencing data with scSpace". In: *Nature Communications* 14.1 (2023), page 2484.
- [24] S. B. Linker, I. Narvaiza, J. Y. Hsu, et al. "Human-specific regulation of neural maturation identified by cross-primate transcriptomics". In: *Current Biology* 32.22 (2022), pages 4797–4807.
- [25] J. R. Xue, A. Mackay-Smith, K. Mouri, et al. "The functional and evolutionary impacts of human-specific deletions in conserved elements". In: *Science* 380.6643 (2023), eabn2253.
- [26] E. Kun and V. M. Narasimhan. "Fast-evolving genomic regions underlie human brain development". In: *Nature* 613 (2023), pages 37–38.
- [27] K. S. Pollard, S. R. Salama, B. King, et al. "Forces shaping the fastest evolving regions in the human genome". In: *PLOS Genetics* 2.10 (2006), e168.
- [28] G. Konopka, J. M. Bomar, K. Winden, et al. "Human-specific transcriptional regulation of CNS development genes by FOXP2". In: *Nature* 462.7270 (2009), pages 213–217.
- [29] R. J. Mangan, F. C. Alsina, F. Mosti, et al. "Adaptive sequence divergence forged new neurodevelopmental enhancers in humans". In: *Cell* 185.24 (2022), pages 4587–4603.
- [30] S. I. O'Donoghue, B. F. Baldi, S. J. Clark, et al. "Visualization of biomedical data". In: *Annual Review of Biomedical Data Science* 1 (2018), pages 275–304.
- [31] C. B. Nielsen, M. Cantor, I. Dubchak, et al. "Visualizing genomes: techniques and challenges". In: *Nature Methods* 7.S3 (2010), S5–S15.
- [32] N. Gehlenborg, S. I. O'donoghue, N. S. Baliga, et al. "Visualization of omics data for systems biology". In: *Nature Methods* 7.Suppl 3 (2010), S56–S68.

- [33] S. Nusrat, T. Harbig, and N. Gehlenborg. “Tasks, techniques, and tools for genomic data visualization”. In: *Computer Graphics Forum*. Volume 38. 2019, pages 781–805.
- [34] C. B. Nielsen, S. D. Jackman, I. Birol, and S. J. Jones. “ABYSS-Explorer: visualizing genome sequence assemblies”. In: *IEEE Trans. Visual. Comput. Graphics* 15.6 (2009), pages 881–888.
- [35] T. O’Brien, A. Ritz, B. Raphael, and D. Laidlaw. “Gremlin: an interactive visualization model for analyzing genomic rearrangements”. In: *IEEE Trans. Visual. Comput. Graphics* 16.6 (2010), pages 918–926.
- [36] J. Aurisano, K. Reda, A. Johnson, and J. Leigh. “Bacterial gene neighborhood investigation environment: A large-scale genome visualization for big displays”. In: *2014 IEEE 4th Symposium on Large Data Analysis and Visualization (LDAV)*. 2014, pages 103–104.
- [37] M. Hess, D. Jente, J. Wiemeyer, et al. “Visual analysis and comparison of multiple sequence alignments”. In: *Proceedings of the Eurographics Workshop on Visual Computing for Biology and Medicine*. 2016, pages 31–40.
- [38] M. Meyer, T. Munzner, and H. Pfister. “MizBee: a multiscale synteny browser”. In: *IEEE Trans. Visual. Comput. Graphics* 15.6 (2009), pages 897–904.
- [39] M. Meyer, B. Wong, M. Styczynski, et al. “Pathline: A tool for comparative functional genomics”. In: *Computer Graphics Forum*. Volume 29. 2010, pages 1043–1052.
- [40] M. Meyer, T. Munzner, A. DePace, and H. Pfister. “MulteeSum: a tool for comparative spatial and temporal gene expression data”. In: *IEEE Trans. Visual. Comput. Graphics* 16.6 (2010), pages 908–917.
- [41] T. Höllt, N. Pezzotti, V. van Unen, et al. “Cytosplore: interactive immune cell phenotyping for large single-cell datasets”. In: *Computer Graphics Forum*. Volume 35. 2016, pages 171–180.
- [42] D. Feng, C. E. Whitehurst, D. Shan, et al. “Single Cell Explorer: collaboration-driven tools to leverage large-scale single-cell RNA-seq data”. In: *BMC Genomics* 20.1 (2019), page 676.
- [43] M. V. Patel. “iS-CellR: A user-friendly tool for analyzing and visualizing single-cell RNA sequencing data”. In: *Bioinformatics* 34.24 (2018), pages 4305–4306.
- [44] V. Gardeux, F. P. A. David, A. Shajkofci, et al. “ASAP: A web-based platform for the analysis and interactive visualization of single-cell RNA-seq data”. In: *Bioinformatics* 33.19 (2017), pages 3123–3125.
- [45] S. M. Huisman, B. van Lew, A. Mahfouz, et al. “BrainScope: Interactive visual exploration of the spatial and temporal human brain transcriptome”. In: *Nucleic Acids Res.* 45.10 (2017), gkx046, e83.
- [46] A. Somarakis, V. Van Unen, F. Koning, et al. “ImaCytE: Visual Exploration of Cellular Micro-Environments for Imaging Mass Cytometry Data”. In: *IEEE Trans. Visual. Comput. Graphics* 27.1 (2019), pages 98–110.

- [47] R. Krueger, J. Beyer, W.-D. Jang, et al. “Facetto: Combining unsupervised and supervised learning for hierarchical phenotype analysis in multi-channel image data”. In: *IEEE Trans. Visual. Comput. Graphics* 26.1 (2020), pages 227–237.
- [48] M. S. Keller, I. Gold, C. McCallum, et al. “Vitesse: Integrative visualization of multi-modal and spatially resolved single-cell data”. In: *Nature Methods* 22 (2025), pages 63–67.
- [49] S. Jänicke, P. Kaur, P. Kuzmicki, and J. Schmidt. “Participatory visualization design as an approach to minimize the gap between research and application”. In: *Proc. VisGap*. Eindhoven, NL: EUROGRAPHICS, 2020, pages 35–42.
- [50] M. Brehmer and T. Munzner. “A multi-level typology of abstract visualization tasks”. In: *IEEE Trans. Visual. Comput. Graphics* 19.12 (2013), pages 2376–2385.
- [51] A. Vieth, T. Kroes, J. Thijssen, et al. “ManiVault: A Flexible and Extensible Visual Analytics Framework for High-Dimensional Data”. In: *IEEE Trans. Visual. Comput. Graphics* 30.1 (2024), pages 175–185.
- [52] M. Bostock, V. Ogievetsky, and J. Heer. “D³ data-driven documents”. In: *IEEE Trans. Visual. Comput. Graphics* 17.12 (2011), pages 2301–2309.
- [53] L. Duhan, D. Kumari, M. Naime, et al. “Single-cell transcriptomics: background, technologies, applications, and challenges”. In: *Mol. Biol. Rep.* 51.1 (2024), page 600.
- [54] C. Darwin. *The origin of species: By means of natural selection, or the preservation of favoured races in the struggle for life*. Cambridge University Press, 1859.
- [55] A. A. Hoffmann and P. A. Parsons. *Extreme environmental change and evolution*. Cambridge University Press, 1997.
- [56] J. Tooby and L. Cosmides. “On the universality of human nature and the uniqueness of the individual: The role of genetics and adaptation”. In: *J. Pers.* 58.1 (1990), pages 17–67.
- [57] C. F. Underwood and L. C. Parr-Brownlie. “Primary motor cortex in Parkinson’s disease: Functional changes and opportunities for neurostimulation”. In: *Neurobiol. Dis.* 147 (2021), page 105159.
- [58] Y. Qiu, J. Wang, J. Lei, and K. Roeder. “Identification of cell-type-specific marker genes from co-expression patterns in tissue samples”. In: *Bioinformatics* 37.19 (2021). Edited by A. Mathelier, pages 3228–3234.
- [59] D. Rosati, M. Palmieri, G. Brunelli, et al. “Differential gene expression analysis pipelines and bioinformatic tools for the identification of specific biomarkers: A review”. In: *Comput. Struct. Biotechnol. J.* 23 (2024), pages 1154–1168.
- [60] J. Costa-Silva, D. Domingues, and F. M. Lopes. “RNA-Seq differential expression analysis: An extended review and a software tool”. In: *PLoS One* 12.12 (2017). Edited by Z. Wei, e0190152.

- [61] J. Ding, A. Alavi, M. R. Ebrahimkhani, and Z. Bar-Joseph. “Computational tools for analyzing single-cell data in pluripotent cell differentiation studies”. In: *Cell Reports Methods* 1.6 (2021), page 100087.
- [62] N. Bhattacharya, C. C. Nelson, G. Ahuja, and D. Sengupta. “Big data analytics in single-cell transcriptomics: Five grand opportunities”. In: *WIREs Data Min. Knowl. Discovery* 11.4 (2021), e1414.
- [63] Y. Jeong, J. Ronen, W. Kopp, et al. “ScMaui: A widely applicable deep learning framework for single-cell multiomics integration in the presence of batch effects and missing data”. In: *BMC Bioinf.* 25.1 (2024), page 257.
- [64] X. Gao, D. Hu, M. Gogol, and H. Li. “ClusterMap: Compare multiple single cell RNA-Seq datasets across different experimental conditions”. In: *Bioinformatics* 35.17 (2019), pages 3038–3045.
- [65] K. Davie, J. Janssens, D. Koldere, et al. “A single-cell transcriptome atlas of the aging *Drosophila* brain”. In: *Cell* 174.4 (2018), pages 982–998.
- [66] J. Zhao, A. Jaffe, H. Li, et al. “Detection of differentially abundant cell subpopulations in scRNA-seq data”. In: *Proc. Natl. Acad. Sci.* 118.22 (2021), e2100293118.
- [67] Y. Wang, L. Chen, J. Jo, and Y. Wang. “Joint t-SNE for comparable projections of multiple high-dimensional datasets”. In: *IEEE Trans. Visual. Comput. Graphics* 28.1 (2022), pages 623–632.
- [68] S. Basu, J. Eggermont, T. Kroes, et al. “Cytosplore Simian Viewer: Visual exploration for multi-species single-cell RNA sequencing data”. In: *Proc. VCBM*. Eindhoven, NL: EUROGRAPHICS, 2023, pages 111–120.
- [69] R. Fang, C. Xia, J. L. Close, et al. “Conservation and divergence of cortical cell organization in human and mouse revealed by MERFISH”. In: *Science* 377.6601 (2022), pages 56–62.
- [70] R. Machold and B. Rudy. “Genetic approaches to elucidating cortical and hippocampal GABAergic interneuron diversity”. In: *Front. Cell. Neurosci.* 18 (2024), page 1414955.
- [71] C. Scharff and J. Petri. “Evo-devo, deep homology and FoxP2: Implications for the evolution of speech and language”. In: *Philos. Trans. R. Soc. B Biol. Sci.* 366.1574 (2011), pages 2124–2140.
- [72] S. Herculano-Houzel. “The human brain in numbers: A linearly scaled-up primate brain”. In: *Front. Hum. Neurosci.* 3 (2009), page 31.
- [73] S. Bremm, T. von Landesberger, M. Hess, et al. “Interactive visual comparison of multiple trees”. In: *Proc. VAST*. New York: IEEE, 2011, pages 31–40.
- [74] M. Harrower and C. A. Brewer. “ColorBrewer.org: An Online Tool for Selecting Colour Schemes for Maps”. In: *The Cartographic Journal* 40.1 (2003), pages 27–37.
- [75] T. A. Hage, A. Bosma-Moody, C. A. Baker, et al. “Synaptic connectivity to L2/3 of primary visual cortex measured by two-photon optogenetic stimulation”. In: *eLife* 11 (2022), e71103.

- [76] J. Brooke. “SUS: A “quick and dirty” usability scale”. In: *Usability Evaluation in Industry*. Edited by P. W. Jordan, B. Thomas, B. A. Weerdmeester, and I. L. McClelland. Volume 189. 194. Taylor and Francis, 1996, pages 189–194.
- [77] C. Stoiber, D. Ceneda, M. Wagner, et al. “Perspectives of visualization onboarding and guidance in VA”. In: *Vis. Inform.* 6.1 (2022), pages 68–83.
- [78] G. A. Bouland, A. Mahfouz, and M. J. T. Reinders. “Consequences and opportunities arising due to sparser single-cell RNA-seq datasets”. In: *Genome Biol.* 24.1 (2023), page 86.
- [79] V. Svensson, S. A. Teichmann, and O. Stegle. “SpatialDE: Identification of spatially variable genes”. In: *Nature Methods* 15 (2018), pages 343–346.
- [80] J. Zhu, S. Sun, and X. Zhou. “SPARK-X: Non-parametric modeling enables scalable and robust detection of spatial expression patterns for large spatial transcriptomic studies”. In: *Genome Biology* 22.1 (2021), page 184.
- [81] C. Li, J. Thijssen, T. Kroes, et al. “SpaceWalker enables interactive gradient exploration for spatial transcriptomics data”. In: *Cell Reports Methods* 3.12 (2023), page 100645.
- [82] J. Boyle, G. Hamm, E. Williams, et al. “Persist: robust identification of spatially variable features in spatial omics datasets via topological data analysis”. In: *arXiv preprint* (2025). arXiv:2505.04360, arXiv–2505.
- [83] Y. Jin, Y. Zuo, G. Li, et al. “Advances in spatial transcriptomics and its applications in cancer research”. In: *Molecular Cancer* 23.1 (2024), page 129.
- [84] X. Wang, W. V. Li, and H. Li. “spCLUE: A contrastive learning approach to unified spatial transcriptomics analysis across single-slice and multi-slice data”. In: *Genome Biology* 26.1 (2025), page 177.
- [85] S. Basu, M. Wirthlin, J. Eggermont, et al. “Cytosplore EvoViewer: Visual analytics of conserved evolutionary patterns in multi-species single-cell sequencing data”. In: *2025 IEEE 18th Pacific Visualization Conference (PacificVis)*. IEEE, 2025, pages 149–159.
- [86] B. Long, J. Miller, and The SpaceTx Consortium. “SpaceTx: A roadmap for benchmarking spatial transcriptomics exploration of the brain”. In: *arXiv preprint* (2023). arXiv:2301.08436, pages 2023–01.
- [87] M. N. Hewitt et al. “A cross-species spatial transcriptomic atlas of the human and non-human primate basal ganglia”. In: *bioRxiv* (2025), pages 2025–11.
- [88] E. Lein, M. Hawrylycz, N. Ao, et al. “Genome-wide atlas of gene expression in the adult mouse brain”. In: *Nature* 445 (2006), pages 168–176.
- [89] B. Tasic, V. Menon, T. Nguyen, et al. “Adult mouse cortical cell taxonomy revealed by single cell transcriptomics”. In: *Nature Neuroscience* 19 (2016), pages 335–346.
- [90] L. Gray, Z. Yao, T. Nguyen, et al. “Layer-specific chromatin accessibility landscapes reveal regulatory networks in adult mouse visual cortex”. In: *eLife* 6 (2017), e21883.

- [91] A. Watakabe, S. Ohsawa, T. Hashikawa, and T. Yamamori. “Binding and complementary expression patterns of semaphorin 3E and plexin D1 in the mature neocortices of mice and monkeys”. In: *Journal of Comparative Neurology* 499 (2006), pages 258–273.
- [92] R. Ferland, T. Cherry, P. Preware, et al. “Characterization of Foxp2 and Foxp1 mRNA and protein in the developing and mature brain”. In: *Journal of Comparative Neurology* 460 (2003), pages 266–279.
- [93] J. Gunnensen, M. Kim, S. Fuller, et al. “Sez-6 Proteins Affect Dendritic Arborization Patterns and Excitability of Cortical Pyramidal Neurons”. In: *Neuron* 56 (2007), pages 621–639.
- [94] K. Miya, K. Keino-Masu, T. Okada, et al. “Expression of Heparan Sulfate Endosulfatases in the Adult Mouse Brain: Co-expression of Sulf1 and Dopamine D1/D2 Receptors”. In: *Frontiers in Neuroanatomy* 15 (2021), page 726718.
- [95] H. Zeng, E. Shen, J. Hohmann, et al. “Large-Scale Cellular-Resolution Gene Profiling in Human Neocortex Reveals Species-Specific Molecular Signatures”. In: *Cell* 149 (2012), pages 483–496.
- [96] Z. Yao, C. van Velthoven, M. Kunst, et al. “A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain”. In: *Nature* 624 (2023), pages 317–332.
- [97] A. Nagalski, L. Puellas, M. Dabrowski, et al. “Molecular anatomy of the thalamic complex and the underlying transcription factors”. In: *Brain Structure and Function* 221 (2015), pages 2493–2510.
- [98] M. Turner, T. Chartrand, M. Summers, et al. “Exploring the correspondence between gene expression and thalamic nuclei using the THALMANAC resource”. In: *openRxiv* (2025), pages 2025–09.
- [99] T. Vitalis, C. Fouquet, C. Alvarez, et al. “Developmental expression of monoamine oxidases A and B in the central and peripheral nervous systems of the mouse”. In: *Journal of Comparative Neurology* 442 (2002), pages 331–347.
- [100] V. Petyuk, W.-J. Qian, M. Chin, et al. “Spatial mapping of protein abundances in the mouse brain by voxelation integrated with high-throughput liquid chromatography–mass spectrometry”. In: *Genome Research* 17 (2007), pages 328–336.
- [101] N. Ochandarena, J. Niehaus, A. Tassou, and G. Scherrer. “Cell-type specific molecular architecture for mu opioid receptor function in pain and addiction circuits”. In: *Neuropharmacology* 238 (2023), page 109597.
- [102] T. Bakken, C. van Velthoven, V. Menon, et al. “Single-cell and single-nucleus RNA-seq uncovers shared and distinct axes of variation in dorsal LGN neurons in mice, non-human primates, and humans”. In: *eLife* 10 (2021), e64875.
- [103] Y. Miyamoto, I. Nagayoshi, A. Nishi, and T. Fukuda. “Three divisions of the mouse caudal striatum differ in the proportions of dopamine D1 and D2 receptor-expressing cells, distribution of dopaminergic axons, and composition of cholinergic and GABAergic interneurons”. In: *Brain Structure and Function* 224 (2019), pages 2703–2716.

- [104] C. Gerfen and D. Surmeier. “Modulation of Striatal Projection Systems by Dopamine”. In: *Annual Review of Neuroscience* 34 (2011), pages 441–466.
- [105] L. Marcuse, M. Langan, P. Hof, et al. “The thalamus: Structure, function, and neurotherapeutics”. In: *Neurotherapeutics* 22 (2025), e00550.
- [106] S. Sherman. “Thalamus plays a central role in ongoing cortical functioning”. In: *Nature Neuroscience* 19 (2016), pages 533–541.
- [107] X. Zhou, P.-Y. Risold, and G. Alvarez-Bolado. “Development of the GABAergic and glutamatergic neurons of the lateral hypothalamus”. In: *Journal of Chemical Neuroanatomy* 116 (2021), page 101997.
- [108] P. Bonnavion, L. Mickelsen, A. Fujita, et al. “Hubs and spokes of the lateral hypothalamus: cell types, circuits and behaviour”. In: *The Journal of Physiology* 594 (2016), pages 6443–6462.
- [109] R. A. Amezcua, A. T. Lun, E. Becht, et al. “Orchestrating single-cell analysis with Bioconductor”. In: *Nature Methods* 17.2 (2020), pages 137–145.
- [110] J. S. Sadick, M. R. O’Dea, P. Hasel, et al. “Astrocytes and oligodendrocytes undergo subtype-specific transcriptional changes in Alzheimer’s disease”. In: *Neuron* 110.11 (2022), pages 1788–1805.
- [111] S. Lehrer and P. H. Rhoenstein. “RORB, an Alzheimer’s disease susceptibility gene, is associated with viral encephalitis, an Alzheimer’s disease risk factor”. In: *Clin. Neurol. Neurosurg.* 233 (2023), page 107984.
- [112] J. B. Gurdon and P.-Y. Bourillot. “Morphogen gradient interpretation”. In: *Nature* 413.6858 (2001), pages 797–803.
- [113] M. S. Cembrowski, J. L. Bachman, L. Wang, et al. “Spatial gene-expression gradients underlie prominent heterogeneity of CA1 pyramidal neurons”. In: *Neuron* 89.2 (2016), pages 351–368.
- [114] H. Zeng. “What is a cell type and how to define it?” In: *Cell* 185.15 (2022), pages 2739–2755.
- [115] T. Lohoff, S. Ghazanfar, A. Missarova, et al. “Integration of spatial and single-cell transcriptomic data elucidates mouse organogenesis”. In: *Nature Biotechnology* 40.1 (2022), pages 74–85.
- [116] S. Dan, M. A. Turner, M. DeBerardine, et al. “Spatial patterning of transcriptional and regulatory programs in the primate subcortex”. In: *bioRxiv* (2025), pages 2025–11.