



Universiteit
Leiden
The Netherlands

Decoding telomere gene variation through saturation genome editing: from functional genomics to clinical interpretation

Obolenski, S.

Citation

Obolenski, S. (2026, September 24). *Decoding telomere gene variation through saturation genome editing: from functional genomics to clinical interpretation*. Retrieved from <https://hdl.handle.net/1887/4310110>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4310110>

Note: To cite this publication please use the final published version (if applicable).

Population-Based Analysis of *POT1* Variants in a Cutaneous Melanoma Case-Control Cohort

Journal of Medical Genetics 2023;60(7):692–696

Irving Simonin-Wilmer¹, Raul Ossio¹, Emmett M Leddin², Mark Harland³, Karen A Pooley⁴, Mauricio Gerardo Martil de la Garza⁵, Sofia Obolenski⁶, James Hewinson^{6,7}, Chi C Wong⁶, Vivek Iyer⁶, John C Taylor^{8,9}, Julia A Newton-Bishop³, D Timothy Bishop¹⁰, Gerardo Andrés Cisneros^{5,11}, Mark M Iles¹², David J Adams⁶, Carla Daniela Robles-Espinoza^{1,6,*}

Affiliations

¹Laboratorio Internacional de Investigación sobre el Genoma Humano, Universidad Nacional Autónoma de México, Campus Juriquilla, Querétaro, Qro, Mexico

²Department of Chemistry, University of North Texas, Denton, Texas, USA

³Section of Epidemiology and Biostatistics, Leeds Institute of Molecular Medicine, University of Leeds, Leeds, UK

⁴Centre for Cancer Genetic Epidemiology, Cambridge University, Cambridge, UK

⁵Department of Chemistry and Biochemistry, The University of Texas at Dallas, Richardson, Texas, USA

⁶CASM, Wellcome Sanger Institute, Hinxton, UK

⁷CeGaT GmbH, Tübingen, Germany

⁸Leeds Institute of Medical Research, University of Leeds, Leeds, UK

⁹Leeds Institute for Data Analytics, University of Leeds, Leeds, UK

¹⁰Section of Epidemiology and Biostatistics, University of Leeds, Leeds, UK

¹¹Department of Physics, The University of Texas at Dallas, Richardson, Texas, USA

¹²Leeds Institute of Cancer and Pathology, University of Leeds, Leeds, UK

Corresponding author drobles@liigh.unam.mx

ABSTRACT

Pathogenic germline variants in the protection of telomeres 1 gene (*POT1*) have been associated with predisposition to a range of tumour types, including melanoma, glioma, leukaemia and cardiac angiosarcoma. We sequenced all coding exons of the *POT1* gene in 2928 European-descent melanoma cases and 3298 controls, identifying 43 protein-changing genetic variants. We performed POT1-telomere binding assays for all missense and stop-gained variants, finding nine variants that impair or disrupt protein–telomere complex formation, and we further define the role of variants in the regulation of telomere length and complex formation through molecular dynamics simulations. We determine that *POT1* coding variants are a minor contributor to melanoma burden in the general population, with only about 0.5% of melanoma cases carrying germline pathogenic variants in this gene, but should be screened in individuals with a strong family history of melanoma and/or multiple malignancies.

Keywords: Genetic Predisposition to Disease, Genetics, Genetic Variation, Genomics

MAIN TEXT

Since the discovery of pathogenic alleles of *CDKN2A* 25 years ago,¹ a number of other variants that increase melanoma risk have been uncovered by genome-wide association studies (GWAS)² and the genomic analysis of melanoma-predisposed families. These variants affect biological pathways related to pigmentation (such as alleles of *MC1R*, the ‘red hair’ gene), naevus count, including genetic variation adjacent to *PLA2G6*, cell cycle and senescence, comprising changes in *CDKN2A* and *CDK4*, and telomere regulation.³ Of note, pathogenic variants in the protection of telomeres 1 gene (*POT1*) have been associated with melanoma, as well as other types of cancer such as glioma,⁴ leukaemia⁵ and lymphoma.⁶ As such, pathogenic germline *POT1* variants have recently been recognised as defining a novel tumour predisposition syndrome.⁷ Genetic variation proximal to *POT1* has also been found to be associated with melanoma in recent large-scale GWAS studies.⁸


POT1 encodes a single-stranded DNA (ssDNA)-binding protein that forms part of the shelterin complex, a group of proteins that have functions in telomere protection by allowing cells to distinguish the ends of chromosomes from sites of DNA damage and also function in regulating telomere length.⁹ In recent years, sequencing of melanoma-predisposed individuals has revealed a number of pathogenic alleles of *POT1*, which affect the ability of *POT1* to bind to ssDNA and therefore lead to longer and abnormal telomeres.^{10–12} This, in turn, may promote carcinogenesis through the accumulation of damage at chromosome ends and a delay in the onset of cell senescence. Further, a recent study has identified *POT1* variants that lead to shorter telomeres,¹³ emphasising the need to identify and catalogue the consequences of these genetic changes in carriers.

As estimates have suggested that *POT1* may be the second major high-penetrance melanoma susceptibility gene after *CDKN2A*, being causal of disease predisposition in 2%–4% of *CDKN2A/CDK4*-negative families,^{10–14} it has been included in multiple panels for genetic testing of melanoma families. As such, and to inform genetic counselling, there is a need to identify which genetic variants abrogate *POT1* function leading to telomere dysregulation, as well as to determine their frequency in population-ascertained melanoma cases. In this study, we performed experimental and bioinformatic analyses to identify germline variants that disrupt the *POT1*–ssDNA complex and lead to telomere length alterations.

This study included 2928 melanoma cases and 3298 controls, making up a total of 6226 European-descent (British) individuals from two distinct melanoma cohorts plus a population cohort (**Online Supplemental Methods**). We sequenced all *POT1* coding exons on the MiSeq platform (reference transcript: ENST00000357628). After alignment, variant calling and quality

filtering, we identified 43 protein-altering variants in *POT1* by Fluidigm PCR-based amplicon sequencing and validated them by target capture with Agilent SureSelect probes and Illumina sequencing (**Online Supplemental Methods, Online Supplemental Figure 1, Online Supplemental Table 1, Online Supplemental File 6**). Of these, 19 have not been reported in the gnomAD 2.1 dataset.¹⁵

To assess whether the detected variants impair telomere regulation, we analysed the ability of *in vitro*-translated POT1 proteins containing all missense and stop-gained variants (38/43 variants in total (**Online Supplemental Table 1**) to bind to a telomere-like oligo via electrophoretic mobility shift assay (EMSA) experiments (**Online Supplemental Methods**). Our results indicate that four variants completely disrupted POT1–ssDNA complex formation (p.Cys59Tyr, p.Arg137His, p.Leu259Ter and p.Arg273Leu), whereas a further five appear to reduce the affinity of the interaction (p.Lys39Asn, p.Lys85Thr, p.Ser99Pro, p.Arg117His and p.Asp224Asn) (**Figure 1A; Online Supplemental Figure 2**). Of these, six had not been reported in the gnomAD 2.1 dataset, and, of note, as expected, all variants that altered POT1–ssDNA binding fall within the *N*-terminal oligonucleotide/oligosaccharide-binding (OB) fold domains.



Variants were classified in three groups according to their pathogenicity: Group 1 variants were confirmed by EMSA to disrupt the POT1–ssDNA complex or were those strongly suspected as pathogenic (frameshift and splice acceptor variants). We included variants with reduced binding in this category due to their high conservation across species (**Online Supplemental Figure 3**) and prior evidence that they may be pathogenic (p.Arg117His¹⁶ and p.Asp224Asn¹¹). In total, 14/43 variants were classified in this group, with 10 of these falling in the OB fold domains (**Figure 2; Online Supplemental Tables 1 and 2**). Group 2 variants were those predicted to be deleterious and probably damaging by both the SIFT and PolyPhen algorithms and did not disrupt POT1–ssDNA binding (4/43 variants). These variants may impair the function of the protein in other ways. The remaining variants (25) were classified into Group 3.




Figure 2: Schematic diagram of Group 1 *POT1* variants.

Variants are shown on the primary protein structure with their consequence (in a coloured circle or triangle) and their presence (red square) or absence (empty square) in publicly available datasets (gnomAD exomes v2.1, dbSNP build 151 and COSMIC v86). The ClinVar track indicates the pathogenicity prediction in ClinVar release 20220804. The OB domains are shown in green. Variants in red font colour are found in cases, those in blue font colour are found in controls and those in black are found in both cases and controls. For details on numbers of cases and controls, see **Online Supplemental Table 1**. Figure created with VCF/Plotein.22

The majority of cases and controls in this study did not carry a *POT1* variant (94.1% cases, 95.1% controls), and the majority of those with a variant had only one variant. No person had more than two variants. In total, three people had a Group 1 variant and a Group 3 variant (two cases, one control) while five people had two Group 3 variants (three cases, two controls). Given the limited number of persons with two variants, each case and control is classified by their most severe mutation. For Group 1, 15 cases (0.51%) carried a variant, while 8 (0.24%)

controls did (p value=0.08, OR for carrying a Group 1 variant compared with no variant (OR)=2.11, 95% CI (0.89 to 5.00)). For Groups 1+2 combined, 22 cases (0.75%) carried a variant, while 14 controls (0.42%) did (p value 0.096, OR=1.78). Finally, for Group 3, 126 cases (4.3%) carried a variant, as did 149 controls (4.6%) (two-tailed Fisher's exact test, p value 0.66), indicating no evidence of increased risk associated with this variant class. Overall, then while about twice as many cases as controls carried predicted pathogenic variants in *POT1*, this difference was not conventionally statistically significant, likely because of limited power even with a study this size. There were also no differences in age of onset, sex, family history or site of presentation by pathogenicity group when compared with those without one of the classified mutations (**Online Supplemental Tables 3–6**).

We next sought to determine whether the variants we detected had any effect on telomere regulation. For this, we measured telomere length in *POT1* variant carriers and non-carriers from the same populations (**Online Supplemental Methods**). After standardising lengths by plate and adjusting them for cohort via a linear model (**Online Supplemental Table 7, Online Supplemental Figure 4**), we observed that only the individuals carrying the p.Lys39Asn (percentile 98 when compared with controls) and the p.Arg273Leu (percentile 99 when compared with controls) variants had telomeres that were substantially longer than the mean (**Figure 1B**). We also observed that some individuals with splice variants or variants that showed reduced DNA binding also had telomeres on the longer side of the distribution (eg, Lys85Thr, percentile 91, p.Leu259Ter, percentile 90, one individual carrying c.1164–1G>A, percentile 97) but others did not (eg, p.Ser99Pro, percentile 31, most individuals with variants in splice sites). Individuals with the p.Asp224Asn variant had telomere lengths scattered throughout the whole distribution in contrast to previous reports suggesting that these variants increase telomere length¹¹ (**Figure 1B**).

Because the p.Lys39Asn, p.Cys59Tyr and p.Asp224Asn variants are found in controls and show POT1–ssDNA complex disruption, we further investigated those using molecular dynamics simulations (**Online Supplemental Methods**). Our results suggest that all three variants affect the dynamics of the system when compared with the wild-type structure, as evidenced by the first and second normal modes (**Online Supplemental Figure 5A–H, Online Supplemental Movie**). Existing protein structures for POT1 also imply that there are conformational differences between the POT1–ssDNA and POT1–ACD structures.^{17 18} As a result, the structural differences noted within the POT1 mutant proteins investigated here may affect shelterin complex formation, but further investigation is necessary. Additional analyses of root mean square deviation, root mean square fluctuation, residue-wise correlations, secondary structure, energy decomposition analysis and hydrogen bond interactions are all

consistent with the computational results reported herein (**Online Supplemental Figure 5I-L, 6–18, Online Supplemental Tables 8- 12**). MM- GBSA was used to assess the protein: DNA-binding affinities. We calculated a $\Delta\Delta H$ of -0.6 to -1.3 , and 21.6 kcal/mol for p.Lys39Asn, p.Asp224Asn and p.Cys59Tyr, respectively. These enthalpies are in agreement with the experimental binding pattern discussed above.

Even though *POT1* seems to be the second major melanoma susceptibility gene, with 2%–4% of *CDKN2A/CDK4*-WT families carrying a pathogenic coding variant in this gene, its contribution to melanoma risk burden in the general population is minor, with $\sim 0.5\%$ of cases carrying pathogenic variants. Telomere length calculations confirm known associations of variants with longer telomeres (p.Arg273Leu,¹⁰ p.Arg117His¹¹) and found associations with other pathogenic variants (p.Lys39Asn, p.Lys85Thr and confirmation of longer telomere length for p.Ala532Pro, percentile 93, a variant originally reported in Ref.¹¹), but for other variants the association with length was not clear (eg, all three carriers of c.1164–1G>A and six of p.Asp224Asn had telomere lengths scattered throughout the distribution). Although a prior study had shown slightly longer telomeres for carriers of p.Arg117His,¹¹ the carrier melanoma case in this cohort had normal-length telomeres. This may reflect the many mechanisms, including other genetic variants and lifestyle, by which telomere length can be affected or the assays used for telomere analysis. Telomere length for some control individuals (without reported melanoma) with pathogenic variants (eg, p.Lys39Asn and both controls carrying p.Asp224Asn) also showed an increase in telomere length, which may portend an increased risk of tumourigenesis in these individuals or indicate that other factors are necessary for melanoma genesis.

Although in this study we have attempted to identify pathogenic *POT1* variants through DNA-binding assays, the function of *POT1* proteins with variants outside the OB fold domains may be compromised by other mechanisms. For example, another study concluded that the *POT1* p.Ala532Pro variant shows impaired ACD binding, which may also lead to telomere dysregulation.¹⁹ Therefore, further systematic experiments are needed to address other *POT1* functions, such as telomere fragility, to provide a more complete catalogue of variants that alter protein function, and therefore that lead to cancer predisposition.

While the number of *POT1* variant carriers in this study is too limited to draw strong conclusions, the lack of any statistically significant difference in age of onset between variant carriers (54.7 years) and non-carriers (54.4 years) in the general population needs some consideration. By comparison and looking at another melanoma high- penetrance gene, in the Leeds Melanoma Cohort, *CDKN2A* variant carriers have an average age of onset of 50 years (based on data included in Ref. ²⁰). The literature contains many examples of families with

particularly early ages of onset for melanoma; these extreme families likely represent the product of interactions of high penetrance variants (in genes such as *CDKN2A* and *POT1*) with contributing lower penetrance variants and risk-associated lifestyle behaviours. Therefore, the analysis of population-based samples provides a more complete description of the impact of high-penetrance variants in the general population. A comparable scenario applies to breast cancer; recent analysis of the UK SEARCH study containing about 12,700 breast cancer diagnoses under the age of 70 years showed an average age of onset of 54.5 years for women without a known variant in a high-penetrance gene. Only *BRCA1* and *BRCA2* variant carriers had notably earlier ages of onset (46.7 and 50.6 years, respectively), while carriers of variants in rarer predisposing genes (*CHEK2*, *PALB2*, *ATM*, *RAD51C*) had average age of onset of between 51.1 years and 58.2 years (A Antoniou, University of Cambridge, personal communication based on data in Ref²¹).

SUPPLEMENTARY DATA

Materials and methods, ethics approval, as well as supplementary figures and data can be found in the online version of this manuscript: <https://doi.org/10.1136/jmg-2022-108776>.

AUTHOR CONTRIBUTIONS

IS-W: sequencing and telomere length data analysis, RO: sequencing data analysis, EML, MGMdIG: molecular dynamics simulation analysis, MH, KAP: qPCR assays and sample management, SO: data analysis, JH, JCT: sample management, CCW: telomere-binding assays, VI: sequence variant calling, JN-B: patient management, manuscript writing, TB: statistical analysis supervision, manuscript writing, GAC: molecular dynamics simulation analysis, manuscript writing, MI: statistical analysis supervision, DA: conceived and supervised study, manuscript writing, CDR-E: conceived and supervised study, sequencing and variant data analysis, manuscript writing.

REFERENCES

1. Kamb A, Gruis NA, Weaver-Feldhaus J, Liu Q, Harshman K, Tavitjian SV, Stockert E, Day RS, Johnson BE, Skolnick MH. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 1994;264:436–40.
2. Law MH, Bishop DT, Lee JE, Brossard M, Martin NG, Moses EK, Song F, Barrett JH, Kumar R, Easton DF, Pharoah PDP, Swerdlow AJ, Kypreou KP, Taylor JC, Harland M, Randerson-Moor J, Akslen LA, Andresen PA, Avril M-F, Azizi E, Scarrà GB, Brown KM, Dębniak T, Duffy DL, Elder DE, Fang S, Friedman E, Galan P, Ghorzo P, Gillanders EM, Goldstein AM, Gruis NA, Hansson J, Helsing P, Hočevár M, Höiom

- V, Ingvar C, Kanetsky PA, Chen WV, Landi MT, Lang J, Lathrop GM, Lubiński J, Mackie RM, Mann GJ, Molven A, Montgomery GW, Novaković S, Olsson H, Puig S, Puig- Butille JA, Qureshi AA, Radford- Smith GL, van der Stoep N, van Doorn R, Whiteman DC, Craig JE, Schadendorf D, Simms LA, Burdon KP, Nyholt DR, Pooley KA, Orr N, Stratigos AJ, Cust AE, Ward SV, Hayward NK, Han J, Schulze H- J, Dunning AM, Bishop JAN, Demenais F, Amos CI, MacGregor S, Iles MM, GenoMEL Consortium, Essen- Heidelberg Investigators, SDH Study Group, Q- MEGA and QTWIN Investigators, AMFS Investigators, ATHENS Melanoma Study Group. Genome- Wide meta- analysis identifies five new susceptibility loci for cutaneous malignant melanoma. *Nat Genet* 2015;47:987–95.
3. Ribero S, Glass D, Bataille V. Genetic epidemiology of melanoma. *Eur J Dermatol* 2016;26:335–9.
 4. Bainbridge MN, Armstrong GN, Gramatges MM, Bertuch AA, Jhangiani SN, Doddapaneni H, Lewis L, Tombrello J, Tsavachidis S, Liu Y, Jalali A, Plon SE, Lau CC, Parsons DW, Claus EB, Barnholtz- Sloan J, Il'yasova D, Schildkraut J, Ali- Osman F, Sadetzki S, Johansen C, Houlston RS, Jenkins RB, Lachance D, Olson SH, Bernstein JL, Merrell RT, Wrensch MR, Walsh KM, Davis FG, Lai R, Shete S, Aldape K, Amos CI, Thompson PA, Muzny DM, Gibbs RA, Melin BS, Bondy ML, Gliogene Consortium. Germline mutations in shelterin complex genes are associated with familial glioma. *J Natl Cancer Inst* 2015;107.
 5. Speedy HE, Kinnersley B, Chubb D, Broderick P, Law PJ, Litchfield K, Jayne S, Dyer MJS, Dearden C, Follows GA, Catovsky D, Houlston RS. Germ line mutations in shelterin complex genes are associated with familial chronic lymphocytic leukemia. *Blood* 2016;128:2319–26.
 6. McMaster ML, Sun C, Landi MT, Savage SA, Rotunno M, Yang XR, Jones K, Vogt A, Hutchinson A, Zhu B, Wang M, Hicks B, Thirunavukarason A, Stewart DR, Koutros S, Goldstein AM, Chanock SJ, Caporaso NE, Tucker MA, Goldin LR, Liu Y. Germline mutations in protection of telomeres 1 in two families with Hodgkin lymphoma. *Br J Haematol* 2018;181:372–7.
 7. Henry M- L, Osborne J, Else T. POT1 Tumor Predisposition. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJ, Mirzaa G, Amemiya A, eds. *GeneReviews®*. Seattle (WA: University of Washington, Seattle, 1993. <http://www.ncbi.nlm.nih.gov/books/ NBK563529/>
 8. Landi MT, Bishop DT, MacGregor S, Machiela MJ, Stratigos AJ, Ghiorzo P, Brossard M, Calista D, Choi J, Fargnoli MC, Zhang T, Rodolfo M, Trower AJ, Menin C, Martinez J, Hadjisavvas A, Song L, Stefanaki I, Scolyer R, Yang R, Goldstein AM, Potrony M, Kypreou KP, Pastorino L, Queirolo P, Pellegrini C, Cattaneo L, Zawistowski M, Gimenez- Xavier P, Rodriguez A, Elefanti L, Manoukian S, Rivoltini L, Smith BH, Loizidou MA, Del Regno L, Massi D, Mandala M, Khosrotehrani K, Akslen LA, Amos CI, Andresen PA, Avril M- F, Azizi E, Soyer HP, Bataille V, Dalmasso B, Bowdler LM, Burdon KP, Chen WV, Codd V, Craig JE, Dębniak T, Falchi M, Fang S, Friedman E, Simi S, Galan P, Garcia- Casado Z, Gillanders EM, Gordon S, Green A, Gruis NA, Hansson J, Harland M, Harris J, Helsing P, Henders A, Hočevár M, Höiom V, Hunter D, Ingvar C, Kumar R, Lang J, Lathrop GM, Lee JE, Li X, Lubiński J, Mackie RM, Malt M, Malvehy J, McAloney K, Mohamdi H, Molven A, Moses EK, Neale RE, Novaković S, Nyholt DR, Olsson H, Orr N, Fritsche LG, Puig- Butille JA, Qureshi AA, Radford- Smith GL, Randerson- Moor J, Requena C, Rowe C, Samani NJ, Sanna M, Schadendorf D, Schulze H- J, Simms LA, Smithers M, Song F, Swerdlow AJ, van der Stoep N, Kukutsch NA, Visconti A, Wallace L, Ward SV, Wheeler L, Sturm RA, Hutchinson A, Jones K, Malasky M, Vogt A, Zhou W, Pooley KA, Elder DE, Han J, Hicks B, Hayward NK, Kanetsky PA, Brummett C, Montgomery GW, Olsen CM, Hayward C, Dunning AM, Martin NG, Evangelou E, Mann GJ, Long G, Pharoah PDP, Easton DF, Barrett JH, Cust AE, Abecasis G, Duffy DL, Whiteman DC, Gogas H, De Nicolo A, Tucker MA, Newton- Bishop JA, Chanock SJ, Demenais F, Brown KM, Puig S, Nagore E, Shi J, Iles MM, Law MH, GenoMEL Consortium, Q- MEGA and QTWIN Investigators, ATHENS Melanoma Study Group, 23andMe, SDH Study Group, IBD Investigators, Essen- Heidelberg Investigators, AMFS Investigators, MelaNostrum Consortium. Genome- Wide association meta- analyses combining multiple risk phenotypes provide insights into the genetic architecture of cutaneous melanoma susceptibility. *Nat Genet* 2020;52:494–504.
 9. de Lange T. Shelterin: the protein complex that shapes and safeguards human telomeres. *Genes Dev* 2005;19:2100–10.
 10. Robles- Espinoza CD, Harland M, Ramsay AJ, Aoude LG, Quesada V, Ding Z, Pooley KA, Pritchard AL, Tiffen JC, Petljak M, Palmer JM, Symmons J, Johansson P, Stark MS, Gartside MG, Snowden H, Montgomery GW, Martin NG, Liu JZ, Choi J, Makowski M, Brown KM, Dunning AM, Keane TM, López- Otín C, Gruis NA, Hayward NK, Bishop DT, Newton- Bishop JA, Adams DJ. Pot1 loss- of- function variants predispose to familial melanoma. *Nat Genet* 2014;46:478–81.
 11. Shi J, Yang XR, Ballew B, Rotunno M, Calista D, Fargnoli MC, Ghiorzo P, Bressac- de Paillerets B, Nagore E, Avril MF, Caporaso NE, McMaster ML, Cullen M, Wang Z, Zhang X, Bruno W, Pastorino L, Queirolo P, Banuls- Roca J, Garcia- Casado Z, Vaysse A, Mohamdi H, Riazalhosseini Y, Foglio M, Jouenne F,

- Hua X, Hyland PL, Yin J, Vallabhaneni H, Chai W, Minghetti P, Pellegrini C, Ravichandran S, Eggermont A, Lathrop M, Peris K, Scarra GB, Landi G, Savage SA, Sampson JN, He J, Yeager M, Goldin LR, Demenais F, Chanock SJ, Tucker MA, Goldstein AM, Liu Y, Landi MT, NCI DCEG Cancer Sequencing Working Group, NCI DCEG Cancer Genomics Research Laboratory, French Familial Melanoma Study Group. Rare missense variants in POT1 predispose to familial cutaneous malignant melanoma. *Nat Genet* 2014;46:482–6.
12. Wong K, Robles- Espinoza CD, Rodriguez D, Rudat SS, Puig S, Potrony M, Wong CC, Hewinson J, Aguilera P, Puig- Butille JA, Bressac- de Paillerets B, Zattara H, van der Weyden L, Fletcher CDM, Brenn T, Arends MJ, Quesada V, Newton- Bishop JA, Lopez- Otin C, Bishop DT, Harms PW, Johnson TM, Durham AB, Lombard DB, Adams DJ. Association of the POT1 germline missense variant p.I78T with familial melanoma. *JAMA Dermatol* 2019;155:604–9.
 13. Kelich J, Aramburu T, van der Vis JJ, Showe L, Kossenkova A, van der Smagt J, Massink M, Schoemaker A, Hennekam E, Veltkamp M, van Moersel CHM, Skordalakes E. Telomere dysfunction implicates POT1 in patients with idiopathic pulmonary fibrosis. *J Exp Med* 2022;219:e20211681.
 14. Potrony M, Puig- Butille JA, Ribera- Sola M, Iyer V, Robles- Espinoza CD, Aguilera P, Carrera C, Malveyh J, Badenas C, Landi MT, Adams DJ, Puig S. Pot1 germline mutations but not TERT promoter mutations are implicated in melanoma susceptibility in a large cohort of Spanish melanoma families. *Br J Dermatol* 2019;181:105–13.
 15. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, Collins RL, Laricchia KM, Ganna A, Birnbaum DP, Gauthier LD, Brand H, Solomonson M, Watts NA, Rhodes D, Singer- Berk M, England EM, Seaby EG, Kosmicki JA, Walters RK, Tashman K, Farjoun Y, Banks E, Poterba T, Wang A, Seed C, Whiffin N, Chong JX, Samocha KE, Pierce- Hoffman E, Zappala Z, O'Donnell- Luria AH, Minikel EV, Weisburd B, Lek M, Ware JS, Vittal C, Armean IM, Bergelson L, Cibulskis K, Connolly KM, Covarrubias M, Donnelly S, Ferriera S, Gabriel S, Gentry J, Gupta N, Jeandet T, Kaplan D, Llanwarne C, Munshi R, Novod S, Petrillo N, Roazen D, Ruano- Rubio V, Saltzman A, Schleicher M, Soto J, Tibbetts K, Tolonen C, Wade G, Talkowski ME, Neale BM, Daly MJ, MacArthur DG, Genome Aggregation Database Consortium. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* 2020;581:434–43.
 16. Calvete O, Martinez P, Garcia- Pavia P, Benitez- Buelga C, Paumard- Hernández B, Fernandez V, Dominguez F, Salas C, Romero- Laorden N, Garcia- Donas J, Carrillo J, Perona R, Triviño JC, Andrés R, Cano JM, Rivera B, Alonso- Pulpon L, Setien F, Esteller M, Rodriguez- Perales S, Bougeard G, Frebourg T, Urioste M, Blasco MA, Benítez J. A mutation in the POT1 gene is responsible for cardiac angiosarcoma in TP53- negative Li- Fraumeni- like families. *Nat Commun* 2015;6:8383.
 17. Rice C, Shastrula PK, Kossenkova AV, Hills R, Baird DM, Showe LC, Doukov T, Janicki S, Skordalakes E. Structural and functional analysis of the human POT1- TPP1 telomeric complex. *Nat Commun* 2017;8:14928.
 18. Lei M, Podell ER, Cech TR. Structure of human POT1 bound to telomeric single- stranded DNA provides a model for chromosome end- protection. *Nat Struct Mol Biol* 2004;11:1223–9.
 19. Liu J, Yu C, Hu X, Kim J- K, Bierma JC, Jun H- I, Rychnovsky SD, Huang L, Qiao F. Dissecting fission yeast shelterin interactions via MICRo- MS links disruption of shelterin bridge to tumorigenesis. *Cell Rep* 2015;12:2169–80.
 20. Harland M, Cust AE, Badenas C, Chang Y- M, Holland EA, Aguilera P, Aitken JF, Armstrong BK, Barrett JH, Carrera C, Chan M, Gascoyne J, Giles GG, Agha- Hamilton C, Hopper JL, Jenkins MA, Kanetsky PA, Kefford RF, Kolm I, Lowery J, Malveyh J, Ogbah Z, Puig- Butille J- A, Orihuela- Segalés J, Randerson- Moor JA, Schmid H, Taylor CF, Whitaker L, Bishop DT, Mann GJ, Newton- Bishop JA, Puig S. Prevalence and predictors of germline CDKN2A mutations for melanoma cases from Australia, Spain and the United Kingdom. *Hered Cancer Clin Pract* 2014;12:20.
 21. Li S, MacInnis RJ, Lee A, Nguyen- Dumont T, Dorling L, Carvalho S, Dite GS, Shah M, Luccarini C, Wang Q, Milne RL, Jenkins MA, Giles GG, Dunning AM, Pharoah PDP, Southey MC, Easton DF, Hopper JL, Antoniou AC. Segregation analysis of 17,425 population- based breast cancer families: evidence for genetic susceptibility and risk prediction. *Am J Hum Genet* 2022;109:1777–88.
 22. Ossio R, Garcia- Salinas OI, Anaya- Mancilla DS, Garcia- Sotelo JS, Aguilar LA, Adams DJ, Robles- Espinoza CD. VCF/Plotein: visualization and prioritization of genomic variants from human exome sequencing projects. *Bioinformatics* 2019;35:4803–5.

