

Granular media: flow & agitations

Dijksman, J.A.

## Citation

Dijksman, J. A. (2009, December 1). *Granular media : flow & agitations*. Retrieved from https://hdl.handle.net/1887/14482

Version: Corrected Publisher's Version

License: License agreement concerning inclusion of doctoral thesis in the

Institutional Repository of the University of Leiden

Downloaded from: <a href="https://hdl.handle.net/1887/14482">https://hdl.handle.net/1887/14482</a>

**Note:** To cite this publication please use the final published version (if applicable).

## **Propositions**

accompanying the thesis

## **Granular Media - Flow & Agitations**

- i A simple frictional model to explain flow profiles in the split-bottom geometry, introduced earlier by Unger (*Phys. Rev. Lett.* 92, 214301 (2004)), also correctly predicts the height dependence of their rheology. *Chapter 2*
- ii Gravitational suspensions in the split-bottom geometry display Bingham behavior in both their flow structure, as well as their rheology. These observations are in agreement with the inertial number theory when it is appropriately adapted to submersed particles.

  Chapter 3 & 5
- iii Exposing granular flows to weak vibrations destroys their rate independent flow behavior. This highlights the importance of the role of even the smallest fluctuations in the flow of granular materials.

  Chapter 6
- iv The compaction of vibrated granular media is not governed uniquely by the dimensionless peak acceleration; the tap duration plays an equally important role.

Chapter 7

- v The study of gravitational suspensions is unduly underrepresented in the scientific literature, given its everyday relevance.
- vi Non-local extensions to the inertial number theory are the most promising candidates to study the "flows of jammed materials".
  - J. Goyon et. al., Nature 454, 84-87 (2008)
- vii The permeability of static porous media can be studied by applying the inertial number theory to flowing gravitational suspensions.
  - C. Cassar et. al., Phys. Fluids 17, 103301 (2005)
- viii Three-dimensional imaging of granular suspensions is faster, easier and cheaper than such imaging of colloidal suspensions.