



**Universiteit
Leiden**
The Netherlands

Implicating genes, pleiotropy, and sexual dimorphism at blood lipid loci through multi-ancestry meta-analysis

Kanoni, S.; Graham, S.E.; Wang, Y.X.; Surakka, I.; Ramdas, S.; Zhu, X.; ... ; Lee,

Citation

Kanoni, S., Graham, S. E., Wang, Y. X., Surakka, I., Ramdas, S., Zhu, X., ... Lee,. (2022). Implicating genes, pleiotropy, and sexual dimorphism at blood lipid loci through multi-ancestry meta-analysis. *Genome Biology*, 23(1). doi:10.1186/s13059-022-02837-1

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3577291>

Note: To cite this publication please use the final published version (if applicable).

RESEARCH

Open Access



Implicating genes, pleiotropy, and sexual dimorphism at blood lipid loci through multi-ancestry meta-analysis

Stavroula Kanoni^{1†}, Sarah E. Graham^{2†}, Yuxuan Wang^{3†}, Ida Surakka^{2†}, Shweta Ramdas^{4†}, Xiang Zhu^{5,6,7,8†}, Shoa L. Clarke^{7,9}, Konain Fatima Bhatti¹, Sailaja Vedantam^{10,11}, Thomas W. Winkler¹², Adam E. Locke¹³, Eirini Marouli¹, Greg J. M. Zajak¹⁴, Kuan-Han H. Wu¹⁵, Ioanna Ntalla¹⁶, Qin Hui^{17,18}, Derek Klarin^{7,19}, Austin T. Hilliard⁷, Zeyuan Wang^{17,18}, Chao Xue², Gudmar Thorleifsson²⁰, Anna Helgadóttir²⁰, Daniel F. Gudbjartsson^{20,21}, Hilma Holm²⁰, Isleifur Olafsson²², Mi Yeong Hwang²³, Sohee Han²³, Masato Akiyama^{24,25}, Saori Sakaue^{24,26,27}, Chikashi Terao²⁸, Masahiro Kanai^{11,24,29}, Wei Zhou^{11,15,30}, Ben M. Brumpton^{31,32,33}, Humaira Rasheed^{31,32,34}, Aki S. Havulinna^{35,36}, Yogasudha Veturi³⁷,

[†]Stavroula Kanoni, Sarah E. Graham, Yuxuan Wang, Ida Surakka, Shweta Ramdas, and Xiang Zhu contributed equally to this work.

[†]Michael Boehnke, Christopher D. Brown, Pradeep Natarajan, Panos Deloukas, Cristen J. Willer, Themistocles L. Assimes, and Gina M. Peloso jointly supervised this work.

*Correspondence: cristen@umich.edu; gpeloso@bu.edu

² Department of Internal Medicine, Division of Cardiology, University of Michigan, Ann Arbor, MI 48109, USA

³ Department of Biostatistics, Boston University School of Public Health, 801 Massachusetts Ave, Boston, MA 02118, USA

Full list of author information is available at the end of the article

Jennifer Allen Pacheco³⁸, Elisabeth A. Rosenthal³⁹, Todd Lingren⁴⁰, QiPing Feng⁴¹, Iftikhar J. Kullo⁴², Akira Narita⁴³, Jun Takayama⁴³, Hilary C. Martin⁴⁴, Karen A. Hunt⁴⁵, Bhavi Trivedi⁴⁵, Jeffrey Haessler⁴⁶, Franco Giulianini⁴⁷, Yuki Bradford³⁷, Jason E. Miller³⁷, Archie Campbell^{48,49}, Kuang Lin⁵⁰, Iona Y. Millwood^{50,51}, Asif Rasheed⁵², George Hindy⁵³, Jessica D. Faul⁵⁴, Wei Zhao^{54,55}, David R. Weir⁵⁴, Constance Turman⁵⁶, Hongyan Huang⁵⁶, Mariaelisa Graff⁵⁷, Ananyo Choudhury⁵⁸, Dhriti Sengupta⁵⁸, Anubha Mahajan⁵⁹, Michael R. Brown⁶⁰, Weihua Zhang^{61,62}, Ketian Yu¹⁴, Ellen M. Schmidt¹⁴, Anita Pandit¹⁴, Stefan Gustafsson⁶³, Xianying Yin¹⁴, Jian'an Luan⁶⁴, Jing-Hua Zhao⁶⁵, Fumihiko Matsuda⁶⁶, Hye-Mi Jang²³, Kyunghoon Yoon²³, Carolina Medina-Gomez⁶⁷, Achilleas Pitsillides³, Jouke Jan Hottenga^{68,69}, Andrew R. Wood⁷⁰, Yingji Ji⁷⁰, Zishan Gao^{71,72,73}, Simon Haworth^{32,74}, Noha A. Yousri^{75,76}, Ruth E. Mitchell^{32,77}, Jin Fang Chai⁷⁸, Mette Aadahl^{79,80}, Anne A. Bjerregaard⁷⁹, Jie Yao⁸¹, Ani Manichaikul⁸², Chii-Min Hwu⁸³, Yi-Jen Hung⁸⁴, Helen R. Warren^{85,86}, Julia Ramirez^{85,87}, Jette Bork-Jensen⁸⁸, Line L. Kårhus⁷⁹, Anuj Goel^{59,89}, Maria Sabater-Lleal^{90,91}, Raymond Noordam⁹², Pala Mauro⁹³, Floris Matteo^{93,94}, Aaron F. McDaid^{95,96}, Pedro Marques-Vidal⁹⁷, Matthias Wielscher⁹⁸, Stella Trompet^{99,100}, Naveed Sattar¹⁰¹, Line T. Møllehave⁷⁹, Matthias Munz¹⁰², Lingyao Zeng^{103,104}, Jianfeng Huang¹⁰⁵, Bin Yang¹⁰⁵, Alaitz Poveda¹⁰⁶, Azra Kurbasic¹⁰⁶, Claudia Lamina^{107,108}, Lukas Forer^{107,108}, Markus Scholz^{109,110}, Tessel E. Galesloot¹¹¹, Jonathan P. Bradfield¹¹², Sanni E. Ruotsalainen³⁵, EWarwick Daw¹¹³,



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Joseph M. Zmuda¹¹⁴, Jonathan S. Mitchell¹¹⁵, Christian Fuchsberger¹¹⁵, Henry Christensen¹¹⁶, Jennifer A. Brody¹¹⁷, Miguel Vazquez-Moreno¹¹⁸, Mary F. Feitosa¹¹³, Mary K. Wojczynski¹¹³, Zhe Wang¹¹⁹, Michael H. Preuss¹¹⁹, Massimo Mangino^{120,121}, Paraskevi Christofidou¹²⁰, Niek Verweij¹²², Jan W. Benjamins¹²², Jorgen Engmann^{123,124}, Noah L. Tsao¹²⁵, Anurag Verma⁴, Roderick C. Sliker^{126,127}, Ken Sin Lo¹²⁸, Nuno R. Zilhao¹²⁹, Phuong Le¹³⁰, Marcus E. Kleber^{131,132}, Graciela E. Delgado¹³¹, Shaofeng Huo¹³³, Daisuke D. Ikeda¹³⁴, Hiroyuki Iha¹³⁴, Jian Yang^{135,136,137}, Jun Liu^{138,139}, Ayşe Demirkan^{139,140}, Hampton L. Leonard^{141,142}, Jonathan Marten¹⁴³, Mirjam Frank¹⁴⁴, Borge Schmidt¹⁴⁴, Laura J. Smyth¹⁴⁵, Marisa Cañadas-Garre^{145,146,147,148}, Chaolong Wang^{149,150}, Masahiro Nakatochi¹⁵¹, Andrew Wong¹⁵², Nina Hutri-Kähönen¹⁵³, Xueling Sim⁷⁸, Rui Xia¹⁵⁴, Alicia Huerta-Chagoya^{155,156,157}, Juan Carlos Fernandez-Lopez¹⁵⁸, Valeriya Lyssenko^{106,159}, Suraj S. Nongmaithem¹⁶¹, Swati Bayyana^{160,161}, Heather M. Stringham¹⁴, Marguerite R. Irvin¹⁶², Christopher Oldmeadow¹⁶³, Han-Na Kim^{164,165}, Seunggho Ryu^{166,167}, Paul R. H. J. Timmers^{143,168}, Liubov Arbeeva¹⁶⁹, Rajkumar Dorajoo^{150,170}, Leslie A. Lange¹⁷¹, Gauri Prasad^{172,173}, Laura Lorés-Motta¹⁷⁴, Marc Pauper¹⁷⁴, Jirong Long¹⁷⁵, Xiaohui Li⁸¹, Elizabeth Theusch¹⁷⁶, Fumihiko Takeuchi¹⁷⁷, Cassandra N. Spracklen^{178,179}, Anu Loukola³⁵, Sailalitha Bollepalli³⁵, Sophie C. Warner^{180,181}, Ya Xing Wang¹⁸², Wen B. Wei¹⁸³, Teresa Nutile¹⁸⁴, Daniela Ruggiero^{184,185}, Yun Ju Sung¹⁸⁶, Shufeng Chen¹⁰⁵, Fangchao Liu¹⁰⁵, Jingyun Yang^{187,188}, Katherine A. Kentistou¹⁶⁸, Bernhard Banas¹⁸⁹, Giuseppe Giovanni Nardone¹⁹⁰, Karina Meidtner^{191,192}, Lawrence F. Bielak⁵⁵, Jennifer A. Smith^{55,54}, Prashantha Hebbar¹⁹³, Aliki-Eleni Farmaki^{194,195}, Edith Hofer^{196,197}, Maoxuan Lin¹⁹⁸, Maria Pina Concas¹⁹⁰, Simona Vaccargiu¹⁹⁹, Peter J. van der Most²⁰⁰, Niina Pitkänen^{201,202}, Brian E. Cade^{203,204}, Sander W. van der Laan²⁰⁵, Kumaraswamy Naidu Chitralla^{206,207}, Stefan Weiss²⁰⁸, Amy R. Bentley²⁰⁹, Ayo P. Doumatey²⁰⁹, Adebawale A. Adeyemo²⁰⁹, Jong Young Lee²¹⁰, Eva R. B. Petersen²¹¹, Aneta A. Nielsen²¹², Hyeok Sun Choi²¹³, Maria Nethander^{214,215}, Sandra Freitag-Wolf²¹⁶, Lorraine Southam^{44,217}, Nigel W. Rayner^{44,59,217,218}, Carol A. Wang²¹⁹, Shih-Yi Lin^{220,221,222}, Jun-Sing Wang^{223,224}, Christian Couture²²⁵, Leo-Pekka Lyytikäinen^{226,227}, Kjell Nikus^{228,229}, Gabriel Cuellar-Partida²³⁰, Henrik Vestergaard^{188,231}, Bertha Hidalgo²³², Olga Giannakopoulou¹, Qiuyin Cai¹⁷⁵, Morgan O. Obura¹²⁶, Jessica van Setten²³³, Xiaoyin Li²³⁴, Jingjing Liang²³⁴, Hua Tang²³⁵, Natalie Terzikhan¹³⁹, Jae Hun Shin²¹³, Rebecca D. Jackson²³⁶, Alexander P. Reiner²³⁷, Lisa Warsinger Martin²³⁸, Zhengming Chen^{50,51}, Liming Li²³⁹, Takahisa Kawaguchi⁶⁶, Joachim Thiery^{109,240}, Joshua C. Bis¹¹⁷, Lenore J. Launer²⁴¹, Huaixing Li¹³³, Mike A. Nalls^{242,243}, Olli T. Raitakari^{201,202,244}, Sahoko Ichihara²⁴⁵, Sarah H. Wild²⁴⁶, Christopher P. Nelson^{180,181}, Harry Campbell¹⁶⁸, Susanne Jäger^{191,192}, Toru Nabika²⁴⁷, Fahd Al-Mulla¹⁹³, Harri Niinikoski^{248,249}, Peter S. Braund^{180,181}, Ivana Kolcic²⁵⁰, Peter Kovacs²⁵¹, Tota Giardoglou²⁵², Tomohiro Katsuya^{253,254}, Dominique de Kleijn²⁵⁵, Gert J. de Borst²⁵⁵, Eung Kweon Kim²⁵⁶, Hieab H. H. Adams^{139,257,258}, M. Arfan Ikram¹³⁹, Xiaofeng Zhu²³⁴, Folkert W. Asselbergs²³³, Adriaan O. Kraaijeveld²³³, Joline W. J. Beulens^{126,259},

Xiao-Ou Shu¹⁷⁵, Loukianos S. Rallidis²⁶⁰, Oluf Pedersen⁸⁸, Torben Hansen⁸⁸,
Paul Mitchell²⁶¹, Alex W. Hewitt^{262,263}, Mika Kähönen^{264,265}, Louis Pérusse^{225,266},
Claude Bouchard²⁶⁷, Anke Tönjes²⁵¹, Yii-Der Ida Chen⁸¹, Craig E. Pennell²¹⁹,
Trevor A. Mori²⁶⁸, Wolfgang Lieb²⁶⁹, Andre Franke²⁷⁰, Claes Ohlsson^{214,271},
Dan Mellström^{214,272}, Yoon Shin Cho²¹³, Hyejin Lee²⁷³, Jian-Min Yuan^{274,275},
Woon-Puay Koh^{276,277}, Sang Youl Rhee²⁷⁸, Jeong-Taek Woo²⁷⁸, Iris M. Heid¹²,
Klaus J. Stark¹², Martina E. Zimmermann¹², Henry Völzke²⁷⁹, Georg Homuth²⁰⁸,
Michele K. Evans²⁸⁰, Alan B. Zonderman²⁸⁰, Ozren Polasek^{250,281},
Gerard Pasterkamp²⁰⁵, Imo E. Hofer²⁰⁵, Susan Redline^{203,204}, Katja Pahkala^{201,202,282},
Albertine J. Oldehinkel²⁸³, Harold Snieder²⁰⁰, Ginevra Biino²⁸⁴, Reinhold Schmidt¹⁹⁶,
Helena Schmidt²⁸⁵, Stefania Bandinelli²⁸⁶, George Dedoussis¹⁹⁴,
Thangavel Alphonse Thanaraj¹⁹³, Sharon L. R. Kardia⁵⁵, Patricia A. Peyser⁵⁵,
Norihiro Kato¹⁷⁷, Matthias B. Schulze^{191,192,287}, Giorgia Grotto^{190,288},
Carsten A. Böger^{189,289,290}, Bettina Jung^{189,289,290}, Peter K. Joshi¹⁶⁸,
David A. Bennett^{187,188}, Philip L. De Jager^{11,291}, Xiangfeng Lu¹⁰⁵,
Vasiliki Mamakou^{292,293}, Morris Brown^{86,294}, Mark J. Caulfield^{85,86},
Patricia B. Munroe^{85,86}, Xiuqing Guo⁸¹, Marina Ciullo^{184,185}, Jost B. Jonas^{182,295,296,297},
Nilesh J. Samani^{180,181}, Jaakko Kaprio³⁵, Päivi Pajukanta²⁹⁸, Teresa Tusié-Luna²⁹⁹,
Carlos A. Aguilar-Salinas³⁰⁰, Linda S. Adair^{301,302}, Sonny Augustin Bechayda^{303,304},
H. Janaka de Silva³⁰⁵, Ananda R. Wickremasinghe³⁰⁶, Ronald M. Krauss³⁰⁷,
Jer-Yuarn Wu³⁰⁸, Wei Zheng¹⁷⁵, Anneke Iden Hollander¹⁷⁴,
Dwaipayan Bhargava^{173,309}, Adolfo Correa³¹⁰, James G. Wilson³¹¹, Lars Lind³¹²,
Chew-Kiat Heng³¹³, Amanda E. Nelson^{169,314}, Yvonne M. Golightly^{57,169,315,316},
James F. Wilson^{143,168}, Brenda Penninx^{69,317}, Hyung-Lae Kim³¹⁸, John Attia^{163,319},
Rodney J. Scott^{163,319}, D. C. Rao³²⁰, Donna K. Arnett³²¹, Steven C. Hunt^{75,322},
Mark Walker³²³, Heikki A. Koistinen^{36,324,325}, Giriraj R. Chandak^{161,326},
Josep M. Mercader^{156,327,328}, Maria C. Costanzo¹⁵⁶, Dongkeun Jang¹⁵⁶,
Noël P. Burt¹⁵⁶, Clicerio Gonzalez Villalpando³²⁹, Lorena Orozco³³⁰,
Myriam Fornage^{154,331}, EShyong Tai^{78,332}, Rob M. van Dam^{78,333},
Terho Lehtimäki^{226,227}, Nish Chaturvedi¹⁵², Mitsuhiro Yokota³³⁴, Jianjun Liu¹⁵⁰,
Dermot F. Reilly³³⁵, Amy Jayne McKnight¹⁴⁵, Frank Kee¹⁴⁵, Karl-Heinz Jöckel¹⁴⁴,
Mark I. McCarthy^{59,218}, Colin N. A. Palmer³³⁶, Veronique Vitart¹⁴³,
Caroline Hayward¹⁴³, Eleanor Simonsick³³⁷, Cornelia M. van Duijn^{138,139},
Zi-Bing Jin^{338,339}, Jia Qu³³⁹, Haretsugu Hishigaki¹³⁴, Xu Lin¹³³,
Winfried März^{131,340,341}, Vilmondur Gudnason^{129,342}, Jean-Claude Tardif¹²⁸,
Guillaume Lettre¹²⁸, Leen M't Hart^{127,343,344,345}, Petra J. M. Elders^{346,347},
Scott M. Damrauer^{4,125,348}, Meena Kumari³⁴⁹, Mika Kivimäki¹²⁴,
Pim van der Harst^{122,233}, Tim D. Spector¹²⁰, Ruth J. F. Loos^{88,119,350},
Michael A. Province¹¹³, Esteban J. Parra¹³⁰, Miguel Cruz¹¹⁸, Bruce M. Psaty^{117,351,352},
Ivan Brandslund^{116,353}, Peter P. Pramstaller¹¹⁵, Charles N. Rotimi²⁰⁹,
Kaare Christensen³⁵⁴, Samuli Ripatti^{35,355,356}, Elisabeth Widén³⁵,
Hakon Hakonarson^{357,358}, Struan F. A. Grant^{4,358,359}, Lambertus A. L. M. Kiemeny¹¹¹,
Jacqueline de Graaf¹¹¹, Markus Loeffler^{109,110}, Florian Kronenberg^{107,108},
Dongfeng Gu^{105,360}, Jeanette Erdmann³⁶¹, Heribert Schunkert^{103,104},
Paul W. Franks¹⁰⁶, Allan Linneberg^{79,80}, J. Wouter Jukema^{99,362}, Amit V. Khe

ra^{11,363,364,365,366}, Minna Männikkö³⁶⁷, Marjo-Riitta Jarvelin^{98,368,369}, Zoltan Kutalik^{95,96},
 Cucca Francesco^{370,371}, Dennis O. Mook-Kanamori^{372,373},
 Ko Willems van Dijk^{374,375,376}, Hugh Watkins^{89,59}, David P. Strachan³⁷⁷,
 Niels Grarup⁸⁸, Peter Sever³⁷⁸, Neil Poulter³⁷⁹, Lee-Ming Chuang³⁸⁰,
 Jerome I. Rotter⁸¹, Thomas M. Dantoft⁷⁹, Fredrik Karpe^{381,382}, Matt J. Neville^{381,382},
 Nicholas J. Timpson^{32,77}, Ching-Yu Cheng^{383,384}, Tien-Yin Wong^{383,384},
 Chiea Chuen Khor¹⁵⁰, Hengtong Li³⁸⁵, Charumathi Sabanayagam^{383,384},
 Annette Peters^{73,104,192}, Christian Gieger^{72,73,192}, Andrew T. Hattersley³⁸⁶,
 Nancy L. Pedersen³⁸⁷, Patrik K. E. Magnusson³⁸⁷, Dorret I. Boomsma^{68,69},
 Allegonda H. M. Willemsen^{68,69}, LAdrienne Cupples^{3,388}, Joyce B. J. van Meurs^{67,139},
 Mohsen Ghanbari^{139,389}, Penny Gordon-Larsen^{301,302}, Wei Huang³⁹⁰,
 Young Jin Kim²³, Yasuharu Tabara⁶⁶, Nicholas J. Wareham⁶⁴,
 Claudia Langenberg^{64,391}, Eleftheria Zeggini^{44,392,217}, Johanna Kuusisto³⁹³,
 Markku Laakso³⁹³, Erik Ingelsson^{9,63,394,395}, Goncalo Abecasis^{14,396},
 John C. Chambers^{61,62,397,398}, Jaspal S. Kooner^{62,379,399,400}, Paul S. de Vries⁶⁰,
 Alanna C. Morrison⁶⁰, Scott Hazelhurst^{58,401}, Michèle Ramsay⁵⁸, Kari E. North⁵⁷,
 Martha Daviglus⁴⁰², Peter Kraft^{56,403}, Nicholas G. Martin⁴⁰⁴, John B. Whitfield⁴⁰⁴,
 Shahid Abbas^{52,405}, Danish Saleheen^{52,406,407}, Robin G. Walters^{50,51,408},
 Michael V. Holmes^{50,51,409}, Corri Black⁴¹⁰, Blair H. Smith⁴¹¹, Aris Baras³⁹⁶,
 Anne E. Justice⁴¹², Julie E. Buring^{47,328}, Paul M. Ridker^{47,328}, Daniel I. Chasman^{47,328},
 Charles Kooperberg⁴⁶, Gen Tamiya⁴³, Masayuki Yamamoto⁴³, David A. van Heel⁴⁵,
 Richard C. Trembath⁴¹³, Wei-Qi Wei⁴¹⁴, Gail P. Jarvik⁴¹⁵, Bahram Namjou⁴¹⁶,
 M. Geoffrey Hayes^{417,418,38}, Marylyn D. Ritchie³⁷, Pekka Jousilahti³⁶,
 Veikko Salomaa³⁶, Kristian Hveem^{31,419,420}, Bjørn Olav Åsvold^{31,419,421},
 Michiaki Kubo⁴²², Yoichiro Kamatani^{423,424}, Yukinori Okada^{26,423,425,426},
 Yoshinori Murakami⁴²⁷, Bong-Jo Kim²³, Unnur Thorsteinsdottir^{20,428},
 Kari Stefansson^{20,438}, Jifeng Zhang², YEugene Chen², Yuk-Lam Ho⁴²⁹,
 Julie A. Lynch^{430,431}, Daniel J. Rader⁴, Philip S. Tsao^{7,9,432}, Kyong-Mi Chang^{433,434},
 Kelly Cho^{429,435}, Christopher J. O'Donnell^{429,435}, John M. Gaziano^{429,435},
 Peter W. F. Wilson^{18,436}, Timothy M. Frayling⁷⁰, Joel N. Hirschhorn^{10,11,437},
 Sekar Kathiresan^{364,11,365}, Karen L. Mohlke¹⁷⁸, Yan V. Sun^{17,18}, Andrew P. Morris⁴³⁸,
 Michael Boehnke^{14†}, Christopher D. Brown^{4†}, Pradeep Natarajan^{11,439,440,441†},
 Panos Deloukas^{1,442†}, Cristen J. Willer^{2,15,443*†}, Themistocles L. Assimes^{7,9,432†} and
 Gina M. Peloso^{3*†} 

Abstract

Background: Genetic variants within nearly 1000 loci are known to contribute to modulation of blood lipid levels. However, the biological pathways underlying these associations are frequently unknown, limiting understanding of these findings and hindering downstream translational efforts such as drug target discovery.

Results: To expand our understanding of the underlying biological pathways and mechanisms controlling blood lipid levels, we leverage a large multi-ancestry meta-analysis ($N = 1,654,960$) of blood lipids to prioritize putative causal genes for 2286 lipid associations using six gene prediction approaches. Using phenome-wide association (PheWAS) scans, we identify relationships of genetically predicted lipid levels to other diseases and conditions. We confirm known pleiotropic associations with cardiovascular phenotypes and determine novel associations, notably with cholelithiasis risk. We perform sex-stratified GWAS meta-analysis of lipid levels and show that 3–5% of autosomal lipid-associated loci demonstrate sex-biased effects. Finally, we report 21 novel lipid loci identified on the X chromosome. Many of the sex-biased autosomal and X chromosome lipid loci show pleiotropic associations with sex hormones, emphasizing the role of hormone regulation in lipid metabolism.

Conclusions: Taken together, our findings provide insights into the biological mechanisms through which associated variants lead to altered lipid levels and potentially cardiovascular disease risk.

Keywords: Cholesterol, Lipids, Genetics, Genome-wide association study, GWAS

Background

Abnormal blood lipid levels are a major cause of cardiovascular disease [1], the leading cause of morbidity and mortality worldwide [2]. Conventional blood lipid measures, low-density lipoprotein cholesterol (LDL-C), total cholesterol (TC), triglyceride (TG), high-density lipoprotein cholesterol (HDL-C), and nonHDL-C (TC – HDL-C), are commonly used in clinical practice to identify individuals at high risk for cardiovascular events. Several treatments for reducing LDL-C, including statins, ezetimibe, and PCSK9 inhibitors [3], also reduce the risk of developing cardiovascular disease.

Genome-wide association studies (GWAS) for blood lipids have identified nearly 1000 associated genetic loci to date [4–23], including our recent multi-ancestry GWAS meta-analysis in 1.65 M individuals [24]. The latter focused on the gains from the multi-ancestry meta-analysis relative to the single-ancestry results, in terms of number of loci, fine-mapping, and polygenic score (PGS) transferability. However, a challenge in the field is that the underlying gene and biological pathways is often unknown for GWAS loci. Within lipid GWAS, prior fine-mapping studies combined with functional follow-up have successfully identified causal genes with high confidence for only a handful of associated GWAS loci, including *SORT1* [25], *TM6SF2* [12], and *ANGPTL3* [26], among others. Highly sophisticated methods are emerging to prioritize causal genes in well-powered GWAS studies, such as the Data-driven Expression-Prioritized Integration for Complex Traits [27] (DEPICT) and the Polygenic Priority Score [28] (PoPS), that take into account genome-wide properties of associated loci and larger sets of associated loci are beneficial. These methods can be combined with algorithms that integrate expression data such as transcriptome-wide association studies (TWAS) and comprehensive experimental data sets such as mouse gene knockouts. Gene sets enriched for

causal genes will enhance our ability to unravel the biological pathways underlying these associations and there is growing interest in using a combination of gene prioritization methods to provide compelling evidence for putative causal genes [29].

In parallel, the linkage of electronic health records with genetic data in large-scale population studies and patient biobanks allows for the systematic exploration of pleiotropy of lipid-associated alleles. While blood lipid levels have a well-documented causal effect on cardiovascular disease based on genetic association studies validated by randomized controlled trials [30–32], genetic pleiotropic associations might exist for other conditions. Unraveling such pleiotropy may yield new biological insights by revealing previously unrecognized connections between blood lipids and both cardiovascular and non-cardiovascular diseases. Phenome-wide association scans (PheWAS) adopt an agnostic approach to test for pleiotropic associations between genetic factors and a wide range of phenotypes [33]. Such knowledge may allow for the identification of lipid levels as novel diagnostic biomarkers, the repurposing of drugs, and the prevention of adverse drug events [34].

Finally, given empirical sex differences in blood lipid distributions, sex-specific genetic associations may yield novel biological insights. Pre-menopausal females have lower levels of LDL-C than same-age males, and HDL-C levels are higher among females of all ages compared to males [35]. Lipid levels also show a greater estimated heritability in females compared with males [36], especially for LDL-C and TC (> 1.3-fold difference). Sexual dimorphism in lipid levels may be partly explained by X chromosome variants. Evidence from human X-linked abnormalities (like Turner or Klinefelter syndromes) suggests an important role of this chromosome in lipid metabolism [37]. This is further corroborated by the lipid and atherosclerosis profiles in the Four Core Genotypes mouse model [38], which comprises XX and XY gonadal males and XX and XY gonadal females. GWAS studies have traditionally understudied the X chromosome due to technical and analytical difficulties. A recent high coverage whole X chromosome sequencing study [39] prioritized *CHRD1* as a candidate causal lipid gene, suggesting with larger sample sizes we may be able to discover additional variation on the X chromosome associated with lipid levels.

In this study, we first prioritize genes at GWAS lipid loci through multiple in silico gene prediction algorithms and experimental data sources using the latest Global Lipids Genetics Consortium multi-ancestry meta-analysis [24]. We then identify novel disease associations related to lipid levels through PheWAS in two large biobanks using PGs. Finally, we perform sex-stratified meta-analysis to compare the associations between males and females to identify genetic loci with sex-specific associations and GWAS meta-analysis of the X chromosome, to better understand lipid level differences between the sexes. Together, our results highlight biological mechanisms through which lipid-associated variants lead to altered lipid levels.

Results

Identifying functional genes in lipid-associated loci

In a GWAS meta-analysis of blood lipid levels from 1.65 million individuals (Additional file 1: Table S1) at 91 million genotyped and imputed genetic variants, we observed a total of 2286 genome-wide significant index variants associated with lipid levels at 923

loci (± 500 kb regions). This corresponded to 416 index variants associated with LDL-C, 539 with HDL-C, 461 with TG, 487 with TC, and 383 with nonHDL-C. Uniquely, we observed 1750 variants associated with one or more lipid levels [24] (Additional file 2: Table S2).

We employed six approaches to identify candidate functional genes for all 2286 lipid associations. Our prioritization approaches include four locus-specific methods that are based on local information around the indexed variant: (1) the closest gene to the index variant, (2) genes with lipid-associated protein-altering variants, (3) colocalized expression quantitative trait loci (eQTL) genes, and (4) nearby genes prioritized by transcript-wide association studies (TWAS). We also used two genome-wide methods that leveraged genome-wide similarities of features: (1) gene-level Polygenic Priority Score (PoPS) [28] and (2) Data-driven Expression-Prioritized Integration for Complex Traits (DEPICT) [27]. We further combined the two genome-wide methods with the locus-specific methods to increase the confidence in prioritized genes: (1) PoPS intersects with any locus-specific methods (PoPS+), and (2) DEPICT intersects with any locus-specific methods (DEPICT+) (Fig. 1). Since the genome-wide gene prioritization approaches can prioritize different genes for different lipid types at the same locus, we report the gene prioritization results for all 2286 lipid-variant associations (Additional file 2: Table S2, Additional file 3: Figure S1).

We took the genes prioritized by PoPS+ and performed text mining to determine whether previous biological evidence supported these genes as playing a role in lipid levels (Additional file 4: Table S3, S4). PoPS+ leverages both locus-specific and genome-wide genetic signals to increase confidence level in prioritized genes [28]. In total, 882 out of 2286 lipid associations were assigned to one potential causal gene based on PoPS+. We identified a group of 466 unique genes among the 882 lipid associations. We determined that 31 out of the 466 PoPS+ genes have over 1000 lipid-related publications, 91 PoPS+ genes have 100–999 lipid-related publications, 321 PoPS+ genes have 1–99 lipid-related publications, and 23 PoPS+ genes had no lipid-related publications retrieved by the text mining algorithm. These 23 genes could indicate novel genes related to lipid levels for future work or be due to incorrect gene prioritization for a small fraction of index variants. (Additional file 5: Table S4). We then randomly selected 466 genes from 18,383 protein-coding genes using by the PoPS as the reference group to estimate the number of lipid-related publications we would expect to see by chance. A Mann–Whitney U test showed that there was a significant difference ($W=52,353$, $p\text{-value} < 2.2 \times 10^{-16}$) between the set of genes identified by PoPS+ compared to the reference set of 466 genes (Additional file 6: Figure S2). The median count of lipid-related publications was 19 for the PoPS+ gene set compared with 2 lipid-related publications for genes in the reference set.

We performed a comprehensive lookup of all PoPS+ prioritized lipid genes in the Therapeutic Target Database 2022 [24] and found 2092 drugs targeting at least one of our 102 PoPS+ prioritized lipid genes observed in the database (Additional file 7: Table S5). Among those 102 PoPS+ genes, we identify known drug target genes including *PCSK9* druggable as subtilisin/kexin type 9 inhibitor, *HMGCR* druggable as HMG-CoA reductase inhibitor, *PDE3A* druggable as phosphodiesterase 3A inhibitor (CILOSTAZOL), and *NRIH4* as a bile acid receptor FXR agonist (URSODIOL). We also identify several




Fig. 1 Schematic of multi-method candidate gene mapping of indexed variants associated with blood lipid levels. We defined indexed variants within the GLGC GWAS summary statistics and performed two similarity-based methods and four locus-based methods to prioritize genes for each of the indexed variants

other potential drug targets [24] such as *LIPG* (lipase G) and *NR1H3* (nuclear receptor subfamily 1 group H member 3), with relevant lipid biology. *LIPG* has phospholipase and triglyceride lipase activities and is a primary determinant of plasma HDL levels. *NR1H3* has an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR, and LRP8 that could be targeted as an LXR-alpha modulator.

Effects of protein-altering lipid alleles with protective effects on CAD, T2D, and NAFLD

Coronary artery disease (CAD), type 2 diabetes (T2D), and non-alcoholic fatty liver disease (NAFLD) are typically characterized by dyslipidemias. We examined protein-altering alleles with favorable lipid profiles for their associations with CAD, T2D, and NAFLD to identify potential cardiovascular drug targets without off-target liver or diabetes effects. Of the 2286 lipid associations, we observed 166 coding index variants. Eighteen coding variants with a protective lipid effect also had a protective effect for CAD

or T2D (p -value < 0.001) and the lipid results colocalized with the CAD or T2D results, as appropriate, with a posterior probability of a shared causal variant > 0.8 (Table 1 and Additional file 8: Table S6). Six of these twenty variants had protective effects for both CAD and T2D, while nine were protective for CAD and three were protective for T2D (Table 1). Additionally, 269 noncoding alleles with a protective lipid effect also had a protective effect for CAD or T2D ($p < 0.001$; Additional file 8: Table S6).

Driver tissues for lipid levels

We applied DESE (Driver-tissue Estimation by Selective Expression) [40] to estimate the driver tissues of five lipid traits using both gene-level and transcript-level eQTL summary statistics from GTEx v8 tissues [41]. We identified liver as the top-ranked tissue for HDL-C (gene-level p -value $= 4.5 \times 10^{-18}$, transcript-level p -value $= 3.0 \times 10^{-26}$), TC (gene-level p -value $= 1.1 \times 10^{-25}$, transcript-level p -value $= 1.4 \times 10^{-33}$), and nonHDL-C (gene-level p -value $= 2.0 \times 10^{-19}$, transcript-level p -value $= 3.9 \times 10^{-29}$) based on both gene-level and transcript-level selective expression (Additional file 9: Figure S3, Additional file 10: Table S7). For LDL-C, we identified the spleen as the top-ranked tissue using GTEx gene-level data (p -value $= 8.3 \times 10^{-20}$), while liver was ranked second (p -value $= 4.8 \times 10^{-17}$). However, when using GTEx transcript-level data, liver was the top-ranked tissue (p -value $= 4.3 \times 10^{-29}$) and second was whole blood (p -value $= 4.3 \times 10^{-20}$). The top tissue for TG according to both GTEx gene-level and transcript-level expression data was whole blood (gene-level p -value $= 6.4 \times 10^{-20}$, transcript-level p -value $= 1.4 \times 10^{-21}$). Spleen and liver were second according to GTEx gene-level and transcript-level expression data, respectively. The results were consistent with previous knowledge that the liver is a major tissue for lipid metabolism. Transcript-level selective expression provided more statistically significant results for the estimated driver tissues compared to the gene-level selective expression, as reported in the original [40].

Polygenic scores for lipid phenotypes and phenome-wide association scans

We have previously reported that a polygenic score (PGS) for LDL-C was most informative when generated from the multi-ancestry GWAS and that the multi-ancestry PGS performed equally well in European-ancestry Americans, African-ancestry Americans, and continental Africans [24]. Using a similar approach, we generated PGS for the other four lipid traits (“Methods”).

We next performed a phenome-wide association scan (PheWAS) for the multi-ancestry lipid PGS (LDL-C PGS previously reported [24]) to identify pleiotropic effects of lipids with other traits in the European subsets of the UK Biobank and the Million Veteran Program (MVP) cohorts. We compared the effect sizes from the PheWAS analysis between the UK Biobank and MVP per lipid PGS and observed a moderate correlation between the two datasets (Additional file 11: Figure S4). The correlation of the PGS effects on all phenotypes between the UK Biobank and MVP ranges from 0.12 for the HDL-C PGS to 0.39 for the TC PGS (Additional file 11: Figure S4). In general, correlations were stronger for the ICD-10-based phecodes (r^2 of 0.42–0.52) compared to the biomarkers (r^2 of 0.06–0.23) (Additional file 11: Figure S4), which may reflect differences in study populations due to varied environmental effects, prevalence of chronic health conditions, and sex distribution. Among PheWAS results with p -value ≤ 0.05 in the UK

Table 1 Protective lipid coding alleles associated with CAD and/or T2D

RSID	Trait	Coding variant	Effect allele (EAF)	Lipid effect	CAD OR (95% CI)	CAD P-value	T2D OR (95% CI)	T2D P-value	NAFLD OR (95% CI)	NAFLD P-value
Protective lipid alleles associated with reduced risk of CAD and T2D										
rs116843064	TG	ANGPTL4 p.Glu40Lys	A (0.020)	−0.238	0.87 (0.85,0.90)	1.92x10 ^{−18}	0.91 (0.86,0.95)	2.30x10 ^{−5}	0.99 (0.84,1.15)	0.85
rs1169288	TC	HNFA p.Ile27Leu	A (0.682)	−0.035	0.97 (0.96,0.98)	2.31x10 ^{−16}	0.95 (0.94,0.96)	7.30x10 ^{−13}	0.95 (0.91,1.00)	0.04
rs2307111	HDL-C	POCS p.His36Arg	C (0.440)	0.016	0.99 (0.98,0.99)	9.09x10 ^{−05}	0.95 (0.94,0.96)	3.30x10 ^{−16}	1.00 (0.96,1.05)	0.85
rs6480771	HDL-C	DUSP13 p.Ser111Gly	T (0.531)	0.008	0.99 (0.98,0.99)	1.44x10 ^{−04}	0.97 (0.96,0.99)	4.40x10 ^{−05}	0.92 (0.88,0.97)	5.21x10 ^{−04}
rs35742417	TG	RREB1 p.Ser155Tyr	A (0.173)	−0.012	0.98 (0.97,0.99)	5.06x10 ^{−04}	0.96 (0.95,0.98)	3.70x10 ^{−06}	0.98 (0.93,1.04)	0.60
rs72681869	TG	SOS2 p.Pro191Arg	C (0.008)	−0.053	0.93 (0.89,0.98)	3.71x10 ^{−03}	0.88 (0.82,0.94)	3.90x10 ^{−04}	0.87 (0.68,1.12)	0.29
Protective lipid alleles associated with reduced risk of CAD										
rs7412	LDL-C	APOE p.Arg176Cys	T (0.076)	−0.517	0.90 (0.88,0.91)	9.94x10 ^{−52}	1.01 (0.98,1.03)	0.55	1.01 (0.93,1.10)	0.84
rs11591147	LDL-C	PCSK9 p.Arg46Leu	T (0.015)	−0.434	0.80 (0.77,0.83)	5.97x10 ^{−36}	1.04 (0.99,1.09)	0.16	1.05 (0.88,1.26)	0.58
rs11601507	LDL-C	TRIM5 p.Val112Phe	C (0.926)	−0.042	0.95 (0.94,0.96)	2.80x10 ^{−12}	0.99 (0.96,1.01)	0.26	1.02 (0.93,1.11)	0.72
rs1132274	HDL-C	RREB1 p.Arg891Leu	C (0.827)	0.017	0.97 (0.96,0.98)	3.57x10 ^{−08}	1.01 (0.99,1.02)	0.43	1.00 (0.94,1.07)	0.91
rs4760	HDL-C	PLAUR p.Leu317Pro	A (0.860)	0.016	0.97 (0.96,0.98)	7.31x10 ^{−07}	0.99 (0.97,1.01)	0.3	0.96 (0.91,1.03)	0.26
rs855791	LDL-C	TM6PSS6 p.Val736Ala	G (0.578)	−0.009	0.98 (0.97,0.99)	1.08x10 ^{−06}	1.00 (0.99,1.01)	0.75	0.94 (0.90,0.98)	4.83x10 ^{−03}
rs58542926	TC	TM6SF2 p.Glu167Lys	T (0.073)	−0.124	0.97 (0.95,0.98)	4.02x10 ^{−06}	1.10 (1.07,1.12)	2.00x10 ^{−14}	1.45 (1.33,1.58)	1.05x10 ^{−16}
rs56196860	HDL-C	FKBP4 p.Asn197Lys	A (0.027)	0.031	0.95 (0.92,0.97)	1.05x10 ^{−05}	0.98 (0.94,1.02)	0.33	1.03 (0.87,1.21)	0.73

Table 1 (continued)

RSID	Trait	Coding variant	Effect allele (EAF)	Lipid effect	CAD OR (95% CI)	CAD P-value	T2D OR (95% CI)	T2D P-value	NAFLD OR (95% CI)	NAFLD P-value
rs72836561	HDL-C	CD300LG p.Arg82Cys	C (0.971)	0.187	0.95 (0.93,0.98)	1.34x10 ⁻⁰⁴	0.98 (0.95,1.02)	0.4	0.98 (0.86,1.12)	0.77
Protective lipid alleles associated with reduced risk of T2D										
rs1800961	HDL-C	HNF4A p.Thr139Ile	C (0.969)	0.134	0.99 (0.97,1.01)	0.3751	0.85 (0.82,0.88)	3.20x10 ⁻²⁰	1.02 (0.90,1.17)	0.74
rs1801253	TG	ADRB1 p.Gly389Arg	C (0.732)	-0.011	1.01 (1.00,1.02)	8.76x10 ⁻⁰³	0.97 (0.96,0.98)	1.9010x ⁻⁰⁵	1.00 (0.95,1.06)	0.85
rs13107325	HDL-C	SLC39A8 p.Ala391Thr	C (0.941)	0.082	1.00 (0.98,1.01)	0.8876	0.95 (0.93,0.98)	3.30x10 ⁻⁰⁴	0.85 (0.78,0.93)	1.85x10 ⁻⁰⁴

Biobank, the correlation was even higher for ICD-10-based phecodes (r^2 of 0.52–0.76) but remained the same for the biomarkers (r^2 of 0.07–0.22).

We meta-analyzed the results from the two cohorts to increase the power of the PheWAS, by matching ICD10-mapped phecodes and biomarkers. In the combined the UK Biobank-MVP PheWAS results, we detected 58 phenotypes associated with the LDL-C PGS at phenome-wide significance level (p -value $\leq 6.5 \times 10^{-5}$, corrected for 773 phenotypes), 165 with the HDL-C PGS, 59 with the TC PGS, 166 with the TG PGS, and 78 with the nonHDL-C PGS (Fig. 2, Additional file 12: Table S8, Additional file 13: Figure S5, Additional file 14: Figure S6, Additional file 15: Figure S7, Additional file 16: Figure S8). As expected, multiple cardiovascular phenotypes related to atherosclerosis, including the expected coronary artery disease as well as aortic aneurysm and essential hypertension, were phenome-wide significantly associated with all five lipid PGSs, indicating increased risk of these diseases for individuals with genetically predicted increased LDL-C, TG, TC, or nonHDL-C or genetically predicted decreased HDL-C. A recent wide-ranging Mendelian randomization analysis confirmed the causal effect of circulating lipids, not only for coronary artery disease, but other cardiovascular outcomes [42]. Additionally, all lipid PGSs were also significantly associated with decreased




Fig. 2 PheWAS meta-analysis results for multi-ancestry LDL-C PGS in the UK Biobank and MVP. The blue horizontal line denotes phenome-wide significance (p -value $\leq 6.5 \times 10^{-5}$, to account for multiple testing of 773 phenotypes) and the red line is genome-wide significance (p -value $\leq 5 \times 10^{-8}$). Phenotypes have been pruned, so that the most significant one per correlated phenotype group (correlation coefficient > 0.2) is retained. Pairwise correlations were estimated with chi-square test and Cramer's V for the dichotomous phenotypes and Pearson's correlation for the continuous phenotypes. AAA: abdominal aortic aneurysm, AD: Alzheimer's disease, AST: aspartate aminotransferase, Atherosclerosis*: atherosclerosis of native arteries of the extremities with intermittent claudication, Hb_con: hemoglobin concentration, IBD: irritable bowel disease, MCH: mean corpuscular hemoglobin, MCV: mean corpuscular volume, PVD: peripheral vascular disease

levels of direct bilirubin (Additional file 12: Table S8, Fig. 2, Additional file 13: Figure S5, Additional file 14: Figure S6, Additional file 15: Figure S7, Additional file 16: Figure S8), indicating genetically predicted lower LDL-C increased levels of bilirubin (Fig. 2). Correspondingly, lipid PGSs were associated with lower risk for cholelithiasis (gallstones) with the opposite direction for TG PGS, indicating that extreme lowering of LDL-C may impact rates of cholelithiasis (Additional file 12: Table S8, Fig. 2, Additional file 13: Figure S5, Additional file 14: Figure S6, Additional file 15: Figure S7, Additional file 16: Figure S8). To further clarify whether this association might be driven by the *ABCG8* gene alone, we excluded from the LDL-PGS all variants within the locus and tested the association between LDL-PGS and cholelithiasis in the UK Biobank. There was no attenuation of the observed association ($OR=0.94$ and $p\text{-value}=7.94 \times 10^{-17}$ without the *ABCG8* locus vs. $OR=0.93$ and $p\text{-value}=1.96 \times 10^{-21}$).

In the PheWAS analysis, we found that the TC and LDL-C PGS were significantly associated with increased levels of HbA1c (beta=0.101 mmol/mol per SD PGS increase, $p\text{-value}=1.21 \times 10^{-23}$ and beta=0.095 mmol/mol per SD PGS increase, $p\text{-value}=4.37 \times 10^{-21}$, respectively), while the HDL-C PGS was associated with decreased levels of HbA1c (beta=−0.257 mmol/mol per SD PGS increase, $p\text{-value}=2.84 \times 10^{-143}$) (Additional file 12: Table S8). Furthermore, genetically predicted increased LDL-C was significantly associated with decreased hemoglobin concentration ($p\text{-value}=1.92 \times 10^{-45}$, similar significant associations for all other lipid PGSs with a reverse direction of effect for TG, Additional file 12: Table S8). As expected, genetically predicted increased LDL-C and TC were both associated with increased risk for Alzheimer's disease [43] ($OR=1.33$ per SD PGS increase, $p\text{-value}=1.74 \times 10^{-44}$ and $OR=1.26$ per SD PGS increase, $p\text{-value}=1.48 \times 10^{-30}$, respectively; Additional file 12: Table S8). To further investigate how this association might be driven by the ApoE locus, we excluded all genetic variants overlapping this gene from the LDL-PGS and repeated the analysis in the UK Biobank. While the association between the LDL-PGS and the risk for Alzheimer's disease was slightly attenuated after removing the ApoE locus ($OR=1.23$ vs. 1.36 per SD PGS increase), the association remained significant ($p\text{-value}=2.51 \times 10^{-21}$). Recent Mendelian randomization studies also provide evidence for the causal effect of lipids on risk for dementia [44] and Alzheimer's disease [45]. The LDL-C and TC PGSs were also associated with increased aspartate aminotransferase levels (a liver enzyme), in accordance with other studies [46]. We also observed inverse associations between LDL-PGS ($p\text{-value}=1.43 \times 10^{-14}$) and TC PGS ($p\text{-value}=8.34 \times 10^{-14}$) with the risk of iron metabolism disorders (Additional file 12: Table S8).

To better understand the loci that drive the association between each of the lipid PGSs and cholelithiasis and cholecystitis, we interrogated the results from the single-variant PheWAS meta-analysis in the UK Biobank and MVP with all lipid multi-ancestry index variants ($N=1750$ unique). We identified 22 genetic variants associated with cholelithiasis and/or cholecystitis at genome-wide significance. Genes prioritized for these index variants included genes already reported to be associated with gallstone disease [47] (*CYP7A1*, *ABCG5/8*), as well as additional genes (*ABCB4*, *LRBA*, *HNF4A*, *NUCB1*, *GATA4*), that may play also a role. Importantly, we found there was overlap (same index variant) between the previously published index variants for gallstone disease and our lipid index variants for these two loci (Additional file 17: Table S9).

Lipid loci show sex-specific effects

Sex-stratified analyses have the potential to identify loci missed by sex-combined analyses [48] as well as to detect loci exhibiting differential effects on lipids between sexes. First, we performed GWAS meta-analysis separately in each sex ($N_{\text{males}} = 749,391$; $N_{\text{females}} = 562,410$), excluding loci discovered in the sex-combined analysis [24]. We identified twelve loci in females and four in males that reached genome-wide significance in the sex-stratified analysis ($p\text{-value} < 5 \times 10^{-8}$; Additional file 18: Table S10, Additional file 19: Table S11, Additional file 20: Table S12) but not in the sex-combined meta-analysis. As variants may show association to a single sex for reasons unrelated to biological sex differences, including differences in sample sizes between groups, we additionally tested for heterogeneity by sex for these variants in GLGC participating cohorts with close to equal number of males and females. Of the sixteen loci, eight showed significant sex-heterogeneity ($p\text{-value} < 0.0031$, Bonferroni-corrected threshold for sixteen tests). For example, the non-synonymous variant rs34372369 (*EPHA1*, p.Pro582Leu) is associated with nonHDL-C only in females (male $p\text{-value} > 0.05$) and shows significant sex-heterogeneity ($p\text{-value} = 0.0012$). This variant has been previously found to be linked with expression levels of the sex hormone-binding globulin gene (*SHBG*) more strongly in males than females [49], suggesting a possible reason for the difference in observed associations. We additionally sought to replicate the sex-heterogeneity results of these variants in 8 independent multi-ancestry cohorts ($N = 311,639$, 77% non-European ancestry, Additional file 21: Table S13). However, we did not detect significant differences in effect sizes between sexes for these variants after accounting for the number of tests ($p\text{-value} > 0.0031$, Additional file 22: Table S14), potentially due to the limited sample size or difference in ancestry makeup.

Second, we tested for a difference in the male- and female-specific effect sizes for each of the index variants identified from the sex-combined multi-ancestry meta-analysis. Of the 1750 unique index variants, 64 showed a significant difference in effect size by sex for one or more traits (Bonferroni correction for the number of index variants in each trait, Additional file 23: Table S15). These were evenly distributed across traits and more often had stronger effects in females than males (67%, Additional file 24: Figure S9). We tested for replication of the sex-specific differences in up to 311,120 participants from eight independent multi-ancestry cohorts not included in the original meta-analysis (Additional file 21: Table S13). Fifty-four of the 64 (84%) variants had effect size differences that were directionally consistent with the original meta-analysis (Additional file 25: Table S16). Of these, 10 had significantly different effect sizes ($p\text{-value} < 7.8 \times 10^{-4}$, Bonferroni correction for 64 variants) and 22 were nominally significant ($p\text{-value} < 0.05$). We attribute the low rate of replication to the small sample size and the differing proportions of ancestry groups within our replication samples, but we cannot dismiss the potential of false positives in the sex-specific discovery results.

We tested whether the observed sex differences could be caused by a higher frequency of cholesterol-lowering medications in males, potentially indicating an insufficient correction for pre-medication cholesterol levels. Among white British individuals in the UK Biobank, variants with significant sex differences had significantly higher effect size estimates on average after excluding individuals on medication (Additional file 26: Figure S10, Additional file 27: Table S17). However, of the 17 variants that exhibited a

significant difference in effect size by sex in the UK Biobank alone, 15 remained significant after excluding individuals taking medications. Based on this observation, the observed differences did not appear to be driven solely, or even primarily, by differences in medication use between sexes. Furthermore, none of the identified sex-specific variants were associated with sex-participation bias [50] (Additional file 28: Table S18), indicating that differential study enrollment between sexes was unlikely to be the cause of the observed sex-specific lipid associations. We next investigated differences in environmental factors between sexes for these variants in the UK Biobank (Additional file 29: Table S19), including alcohol use [48], smoking status [48], body mass index (BMI) [51], and waist-hip ratio adjusted for BMI (WHRadjBMI) [51]. Twenty-two of the variants (34%) with differential effects on lipids by sex also exhibited a significant difference by sex for WHRadjBMI and one variant had a significant difference by sex for alcohol use (*ADH1B* p.His48Arg). The observed sex differences may therefore be partially attributed to pleiotropic associations with other traits.

Finally, we annotated each locus that showed significant sex differences with regulatory information to identify biological mechanisms that could underlie this difference. Of the 64 lipid variants with significant sex-stratified associations, 14 colocalized (posterior probability of $H_4 > 0.8$) with expression of 20 genes in lipid-related tissues (liver, adipose, or skeletal muscle; Additional file 30: Table S20). Eight of these loci also show a sex-biased eQTL effect in at least one tissue in the direction concordant with the observed sex specificity of the GWAS effect (Additional file 30: Table S20). Among these ten is *CETP*, a gene with strong prior evidence for association with lipids, and *UGT2B17* [20] (Additional file 31: Supplementary Note, Fig. 3). The lead variant of *UGT2B17*, rs4860987, shows a significantly stronger effect of LDL-C in males ($\text{Beta}_{\text{male}} = 0.042$, $\text{SE}_{\text{male}} = 0.002$, $\text{Beta}_{\text{female}} = 0.016$, $\text{SE}_{\text{female}} = 0.003$, $p\text{-value}_{\text{difference}} = 4.2 \times 10^{-15}$) and colocalizes with a male-specific liver eQTL associated with increased expression of *UGT2B17*. Common variants at this locus are in moderate LD ($R^2 = 0.51$) with a common copy number variation (CNV), which may mediate the causal effect (Additional file 31: Supplementary Note). *UGT2B17* plays a role in the metabolism of androgens [52], including testosterone, which is consistent with the observed pleiotropic relationship of this locus with testosterone in males (Additional file 30: Table S20). We note that the index variant in *UGT2B17*, rs4860987, did not show significant sex-specific effects in the replication cohorts, but this could be due to varying frequencies for the index variant between ancestry groups and the moderate LD to the causal CNV in the region. We observed that the combined frequency of rs4860987 across the replication studies was much lower (8%) compared with our combined frequency in the discovery (24%) due to differing proportions of ancestry groups and, along with the lower number of individuals ($N = 218,437$), led to a much-reduced power to replicate this sex-specific effect.

Lipid-associated loci on the X chromosome

Lastly, we meta-analyzed association statistics for 3.1 million X chromosomal variants, including PAR regions, across 1,238,180 individuals from multiple ancestry groups. We identified 28 index variants significantly associated with lipid levels (Additional file 32: Table S21), of which 21 have not been previously reported [20, 39, 53] (15 for HDL-C, 4 for LDL-C, 4 for TC, 5 for TG and 4 for nonHDL-C, Table 2). Among these 28 loci, two




Fig. 3 Sex specificity at the *UGT2B17* Locus. **A** The association signal for LDL-C (top panel) colocalizes with the *UGT2B17* eQTL signal in the liver (bottom panel). **B** The effect sizes of this variant on LDL-C and *UGT2B17* expression are both significantly higher for males (red) compared to females (blue)

Table 2 Novel X chromosome lipid-associated loci

RSID	Position in chromosome X (hg19)	EA/NEA	Annotation (closest gene)	Associated trait	EAF	N	Effect size (SE) from METAL	Ancestry and GC corrected p-value from MR-MEGA	Sex difference p-value
rs35143646	2,856,155	T/C	Missense (ARSL)	LDL-C	0.6352	1,038,070	0.0115 (0.0014)	1.83×10^{-16}	0.1791
				TC	0.6449	1,100,310	0.0091 (0.0014)	1.47×10^{-14}	0.08484
				nonHDL-C	0.6822	712,983	0.0112 (0.0017)	2.08×10^{-11}	0.5294
rs5934507	8,917,206	G/A	Intergenic (FAM98)	HDL-C	0.274	1,158,000	−0.0076 (0.0012)	9.99×10^{-10}	5.83×10^{-4}
rs191084933	14,133,208	G/A	Intergenic (GEMIN8)	HDL-C	0.0055	1,052,630	0.0694 (0.0101)	5.20×10^{-10}	0.3272
rs7888119	16,813,128	T/C	Intronic (TXLNG)	HDL-C	0.546	1,158,000	0.0114 (0.0012)	1.47×10^{-20}	0.2212
rs2230488	20,204,461	T/G	Synonymous (RPS6KA3)	LDL-C	0.1772	1,135,110	−0.0143 (0.0015)	6.90×10^{-18}	0.9297
				TC	0.1745	1,237,380	−0.0114 (0.0014)	4.68×10^{-12}	0.8906
rs6527977	20,322,238	A/C	Intergenic (RPS6KA3)	nonHDL-C	0.1602	789,200	−0.0147 (0.0020)	3.06×10^{-10}	0.7000
rs12012576	21,813,178	G/A	Intergenic (SMPX)	TG	0.3103	1,160,340	−0.0072 (0.0012)	4.88×10^{-8}	0.4558
rs6609434	46,636,767	C/A	Intergenic (SLC9A7)	HDL-C	0.4959	767,051	0.0060 (0.0014)	4.25×10^{-8}	0.9035
rs113957181	49,848,600	A/C	Intronic (CLCN5)	HDL-C	0.0566	452,268	−0.0297 (0.0047)	1.39×10^{-9}	0.1896
rs782397956	53,993,589	T/C	Intronic (PHF8)	HDL-C	0.2050	621,342	−0.0009 (0.0021)	2.65×10^{-15}	0.2585
rs72305711	55,981,911	T/TTA	Intronic (RPI3-188A5.1)	HDL-C	0.1111	534,967	−0.0167 (0.0033)	2.22×10^{-22}	0.03202
rs5914559	56,139,739	G/T	Intergenic (KLF8)	TG	0.6697	738,155	−0.0131 (0.0016)	4.44×10^{-16}	0.3224
rs5964416	64,368,487	A/C	Intergenic (ZC4H2)	HDL-C	0.0924	614,283	−0.0316 (0.0016)	2.68×10^{-11}	0.1478
rs5965342	66,204,144	T/C	Intergenic (EDA2R)	HDL-C	0.2037	745,721	−0.0212 (0.0020)	1.20×10^{-29}	0.02050
rs505520	66,258,914	C/A	Intergenic (EDA2R)	TG	0.2191	750,415	0.0217 (0.0019)	2.91×10^{-30}	0.004371
rs771540123	67,967,645	A/G	Intergenic (STARD8)	nonHDL-C	0.0027	15,311	−0.0202 (0.115)	1.61×10^{-9}	NA
rs5937000	70,047,788	C/T	Intronic (TEX11)	HDL-C	0.4515	796,971	0.0075 (0.0013)	9.55×10^{-9}	0.02603

Table 2 (continued)

RSID	Position in chromosome X (hg19)	EA/NEA	Annotation (closest gene)	Associated trait	EAF	N	Effect size (SE) from METAL	Ancestry and GC corrected <i>p</i> -value from MR-MEGA	Sex difference <i>p</i> -value
rs5938008	74,496,225	C/T	Intronic (<i>UPR1</i>)	HDL-C	0.9167	1,072,550	0.0174 (0.003)	1.24x10 ⁻⁸	0.7235
rs1802288	99,890,204	T/C	Missense (<i>TSPAN6</i>)	LDL-C	0.1665	714,113	0.0159 (0.0019)	1.42x10 ⁻¹⁶	0.9642
				TC	0.1684	753,479	0.0128 (0.0018)	2.49x10 ⁻¹⁰	0.8646
rs139144471	117,829,694	G/T	Intergenic (<i>DOCK11</i>)	HDL-C	0.0876	1,098,540	− 0.0121 (0.0022)	1.16x10 ⁻⁸	0.1111
rs6648533	122,804,678	C/T	Intronic (<i>THOC2</i>)	HDL-C	0.299	1,106,090	− 0.0100 (0.0013)	2.24x10 ⁻¹²	0.4516
rs5929738	135,265,287	C/A	Intronic (<i>FHL1</i>)	HDL-C	0.4679	970,330	0.0071 (0.0011)	3.21x10 ⁻¹⁰	0.4197
rs5975692	135,266,089	G/A	Intronic (<i>FHL1</i>)	TG	0.465	1,138,740	− 0.0084 (0.0011)	7.87x10 ⁻¹²	0.9215
rs2070826	153,582,198	C/T	Intronic (<i>FLNA</i>)	HDL-C	0.1427	1,141,020	0.0170 (0.0017)	6.65x10 ⁻²⁶	0.08671
rs11593	153,627,145	C/A	Intronic (<i>RPL10</i>)	TG	0.1586	1,143,360	− 0.0166 (0.0017)	5.76x10 ⁻²¹	0.8695
rs7886627	153,679,609	G/A	Intergenic (<i>FAM50A</i>)	nonHDL-C	0.1214	771,706	− 0.0147 (0.0022)	9.26x10 ⁻⁹	0.06819
rs1050828	153,764,217	T/C	Missense (<i>G6PD</i>)	LDL-C	0.0113	744,968	− 0.0514 (0.0061)	2.54x10 ⁻¹⁵	0.6533
rs762517	153,764,734	A/G	Intronic (<i>G6PD</i>)	TC	0.0142	798,600	− 0.0480 (0.0057)	5.26x10 ⁻¹⁶	0.7594

EA Effect allele, NEA Non-effect allele, EAF Effect allele frequency, N Number of samples, SE Standard error of the effect size, GC Genomic control, LDL-C Low-density lipoprotein cholesterol, TC Total cholesterol, HDL-C High-density lipoprotein cholesterol, TG Triglycerides

have index variants with a minor allele frequency (MAF) < 1% and three index variants are missense mutations (in genes *ARSL*, *TSPAN6*, and *G6PD*), all of which are novel. We validated the identified X chromosomal associations in up to 255,475 individuals from seven multi-ancestry cohorts (Additional file 21: Table S13). Twenty index variants were at least nominally associated (p -value < 0.05), with five reaching genome-wide significance in the replication cohorts alone (p -value < 5×10^{-8} , Additional file 32: Table S21).

We additionally considered potential sex differences for the X chromosome variants. A missense variant in *RENB*P with MAF = 2.5% reached genome-wide significance only in males but was not significant in the sex-combined meta-analysis or in the female-only analysis (p -value = 4.59×10^{-8} , 0.003 and 0.2, respectively). We also observe three X chromosome loci with significant heterogeneity in effects between sexes; however, these were not significant in the replication cohorts alone, possibly due to the lower sample size (Bonferroni correction for the number of index variants in each trait, Additional file 32: Table S21).

Using a PheWAS approach in the UK Biobank, we found four of the novel loci to have pleiotropic associations with body composition traits (*FAM9B* [HDL-C], *EDA2R* [HDL, TG], *TSPAN6* [LDL-C, TC], and *DOCK11* [HDL-C]), four variants with coronary atherosclerosis and ischemic heart disease, three with immune-related biomarkers (*SLC9A7* [HDL-C], *CLCN5* [HDL-C], *THOC2* [HDL-C]), and two with blood clotting-related biomarkers (*KLF8* [TG], *TEX11* [HDL-C]) (Additional file 32: Table S21). Interestingly, two of the three sex-biased X chromosome variants demonstrate the most significant association with testosterone of all lipid X chromosome variants tested in the PheWAS (rs505520: beta/SE = $-0.089/0.007$ nmol/L per TG-increasing allele and rs5934507: beta/SE = $0.237/0.006$ nmol/L per HDL-increasing allele).

Discussion

In this study, we identify and prioritize likely candidate genes at lipid-associated loci discovered through a variety of approaches including multi-ancestry meta-analysis of autosomes [24] (~ 91 million variants) and the X chromosome (~ 3 million variants), as well as sex-specific meta-analyses using sample sizes ranging from 1.35 to 1.65 million individuals. We previously reported a comparison of multi-ancestry vs single-ancestry lipid findings using autosomal chromosomes and identified improvements in fine-mapping of credible sets and PGS performance, with slight differences in the number of identified loci by ancestry group [24]. Here, we add X chromosome and sex-specific discovery results. We also focus on lipid biology by prioritizing implicated genes, identifying novel phenotypes and diseases associated with genetically predicted lipid levels, and predicting candidate drug target genes.

Our results from this effort translate our GWAS findings for three complementary research areas, helping us further elucidate the biological mechanisms underlying the lipid-associated genetic variants. We first sought to identify methods for prioritization of functional genes at GWAS loci by performing six gene prioritization methods. Lipids are an excellent exemplar phenotype for gene prioritization algorithms because of a wealth of GWAS loci (~ 1000), Mendelian dyslipidemia genes (21), and mouse dyslipidemia phenotypes observed in gene knockouts (740). While the gene prioritization approaches are not independent of each other, integrating several prioritization predictors provides

higher confidence when attempting to characterize causal genes. Others have also highlighted the importance of such frameworks in different diseases [29, 54, 55].

We identify 466 unique genes by combining evidence from a global approach (PoPS) with local gene prioritization approaches. The vast majority of these genes had many lipid-related publications, suggesting the accuracy of our combined prioritization approach. Twenty-three PoPS + identified genes had no lipid-related publications, indicating they could be truly novel or possibly were incorrectly prioritized. Functional validation of the larger pool of prioritized genes, which will require highly parallel experimental methods, will help to further optimize bioinformatics algorithms to prioritize genes and is beyond the scope of this manuscript.

Our prioritization approach also indicates several genes as potential drug targets including *PDE3A* and *NR1H4*. *PDE3A* encodes the phosphodiesterase 3A gene and is predicted to be druggable as phosphodiesterase 3A inhibitor (CILOSTAZOL). Cilostazol has antiplatelet, anti-proliferative, vasodilatory, and ischemic-reperfusion protective properties [56] and has been previously suggested for the primary or secondary prevention of CAD [22]. *NR1H4* encodes a bile acid receptor and regulates the expression of genes involved in bile acid synthesis and transport. The target gene is predicted to be druggable as a bile acid receptor FXR agonist (URSODIOL). Ursodiol is used to treat primary biliary cirrhosis and cholelithiasis and could be a potential candidate for drug repurposing.

We also identify eighteen coding variants where the protective lipid allele is also protective for CAD or T2D. Among these, *PCSK9* is a well-documented drug target, not only for lipids but also for cardiovascular events [57–59]. In comparison to published studies [60], others find a non-significant increased risk for T2D [61] and an arguably stronger protective effect for CAD [62], for *PCSK9* variant carriers. Our observation is consistent with the lack of excess T2D risk observed in *PCSK9* inhibitor clinical trials [57–59, 63] and with strong protective effects for coronary heart disease [64]. Furthermore, these variants are potential therapeutic targets for protective lipid profiles and lowering risk of disease.

Our second goal was to identify diseases that may benefit from lipid-lowering as well as diseases or traits that may become problematic due to very low lipids. To accomplish this, we examined the association of genetically predicted lipid traits (using PGS) with 773 phenotypes in 478,556 individuals. We observed that genetically predicted increased LDL-C, TC, and HDL-C levels, or decreased TG levels, decrease the risk of cholelithiasis. Prior epidemiological studies have not consistently reported an association between lipid levels and risk of gallstones, with some studies showing that increased levels of LDL-C, TC, and TG and decreased levels of HDL-C predispose to the risk for cholelithiasis [65, 66], but others showing no association [67, 68]. Our results are corroborated by a recent Mendelian randomization meta-analysis study in the FinGen and UK Biobank cohorts [69]. The prioritized genes for the individual index lipid variants significantly associated with cholelithiasis in the PheWAS analysis include *ABCG8*, a hepatic cholesterol transporter, responsible for the efflux of cholesterol from the enterocytes to the lumen and from the hepatocytes into bile [70]. The lipid-decreasing allele of index variant in *ABCG8*, rs4245791, has been previously associated with high risk for gallstone disease [47] and high intestinal cholesterol absorption [71], possibly

mediated by an increased expression of *ABCG8* [72]. Furthermore, even after excluding *ABCG5/8* variants from the LDL-PGS, the association with the risk of cholelithiasis was not attenuated. These PGS-PheWAS results suggest the existence of many other cholesterol transporters like *ABCG8* that modify blood cholesterol levels perhaps in large part by facilitating an increased secretion of cholesterol into the biliary system, which in turn increases the risk of the formation of gallstones through the supersaturation of bile. We also observed that HbA1c levels were elevated among subjects with genetically predicted increased LDL-C and TC and with genetically predicted decreased HDL-C. Previous epidemiological studies have established associations between dyslipidemia (increased LDL-C, TC, TG, and decreased HDL-C levels) and increased HbA1c levels among subjects with T2D, as well as insulin-resistant subjects without diabetes [73, 74]. Our observations support a strong genetic basis to these associations and are in accordance with previous studies showing shared pathways between lipid biology, T2D, and HbA1c [75], as well as pleiotropic effects of blood red cell variants with lipid levels [76]. Mendelian randomization studies have shown that hemoglobin and LDL show bidirectional inverse relationships and hemoglobin effects on LDL are also mediated through Hb1Ac, implying that genetic variation influencing erythrocytic factors could also determine lipid levels and the opposite [77]. While most of our significant PheWAS findings could be confirmed via Mendelian randomization studies, we cannot exclude the possibility of spurious associations due to pleiotropy.

Lastly, we sought to expand the coverage of the genome and performed the most comprehensive GWAS of lipid levels to date by including assessment of 3 million variants on the X chromosome as well as explicitly testing for sex-specific effects across 23 chromosomes in 1.35 million individuals of diverse ancestries. We report 21 novel X chromosome loci, including an LDL-lowering locus involving a missense variant in *G6PD*, known to cause G6PD deficiency (p.V68M) [78]. The proposed mechanism is via the inhibition of the NADPH-dependent hydroxymethylglutaryl-CoA (HMG-CoA) reductase, resulting in decreased cholesterol biosynthesis, even though the protective effect of the G6PD deficiency on cardiovascular risk is debatable [79].

We also observed that approximately 3–5% of the genome-wide lipid index variants exhibited differential effects between sexes, which did not seem to be due to differential prevalence in the use of lipid medications or study selection bias. These findings may have important implications in the interpretation of lipid biology, the identification of novel drug targets, and possibly for more accurate prediction of blood cholesterol-related conditions. For example, the *UGT2B17* locus, one of the ten sex-biased loci with corresponding sex-biased eQTL effect, is known to be implicated in androgen and drug metabolism [52]. A common CNV in the region, partially tagged by the lipid index variant, is associated with significant variations in expression levels between ethnic groups [80], which would explain lack of replication in the set of independent studies, and the deletion has been linked to testosterone-related decreased BMI levels [81], as well as decreased risk for osteoporosis in men [82].

Several of the reported sex-biased and X chromosome loci showed significant pleiotropic effects with sex hormone levels, including testosterone and SHBG, highlighting the role of hormone regulation in lipid metabolism [83]. In particular, we

observe an overall inverse effect between the X chromosome lipid index variants and the sex hormone levels. Observational studies have long suggested a potential influence of the sex hormones on the risk for cardiovascular risk [84] but this hypothesis has not been consistently supported by recent Mendelian randomization studies, raising the issues of reverse causality [85, 86].

Conclusions

In conclusion, we leverage the power of a large multi-ancestry GWAS study to further our understanding of lipid metabolism and the impact on chronic diseases. We identify novel lipid loci on the X chromosome and autosomal loci with evident sex-biased lipid effects. We compare a range of gene prioritizing methods to identify causal genes, an approach applicable to studying other complex traits. We additionally further our understanding of lipid metabolism through a phenome-wide study that implicates a relationship between genetically predicted low cholesterol with risk of cholelithiasis.

Methods

Meta-analysis

Summary statistics for sex-combined autosomal analyses were previously published [24]. Following the same procedure, we carried out meta-analyses stratified by sex for 5 lipid traits (HDL-C, LDL-C, TG, nonHDL-C, and TC) for both the autosomes and chromosome X. The sample size for chromosome X (Total $N = 1,238,180$; males = 749,391; females = 562,410) was lower than available for autosomes as not all participating biobanks submitted results for chromosome X. Quality control of summary statistics from contributing cohorts was performed using EasyQC [87]. Prior to meta-analysis, we removed variants with low imputation info scores ($r^2 < 0.3$), those with minor allele count < 3 , and those with Hardy–Weinberg equilibrium p -value $< 1 \times 10^{-8}$. Variants on the X chromosome were filtered using the female imputation info scores and Hardy–Weinberg equilibrium p -values. Summary statistics were corrected by the genomic-control factor calculated from the median p -value of variants with minor allele frequency $> 0.5\%$. For cohorts that contributed summary statistics imputed both on the Haplotype Reference Consortium (HRC) and 1000 Genomes Population v3 (1KGP3) panels, we generated a single file containing all possible variants, favoring those imputed from the HRC imputation panel due to generally higher imputation quality of these variants. Multi-ancestry meta-analysis was performed with MR-MEGA [88] with 5 principal components and using the inverse-variance weighted method in METAL to estimate effect sizes [89]. Independent loci were defined with physical distance > 500 kb or genetic distance > 0.25 cM, whichever one would result in a larger window, followed by a conditional analysis using rareGWAMA [90] as previously described [24], to identify index variants that were shadows of nearby, more-significant associations. Conditional analysis for chromosome X used a female-only UK Biobank LD reference ($N = 21,510$). In line with the analysis in the autosomes, a locus was identified as dependent if the effect size after conditioning

on the most significant variant in the area was more than 1.43 times smaller than the original (95th percentile of the effect size ratios for chromosome X).

Differences in effect size between males and females were tested within each cohort using [91]:

$$Z = \frac{B_m - B_f}{\sqrt{se_m^2 + se_f^2 - 2 * r * se_f * se_m}}$$

and were then meta-analyzed across studies using METAL, to account for cohort-specific ascertainment (e.g., enrichment of cases for type 2 diabetes), or demographics, such as age.

Replication

We collected summary statistics from 8 cohorts across 6 ancestry groups, including African or African American, East Asian, European, Hispanic, Middle Eastern, and South Asian. Each cohort provided sex-stratified and X chromosome association results for the tested traits, as available. The difference in effect sizes between males and females was calculated within each cohort as described above and then meta-analyzed across studies using METAL. X chromosome association results were meta-analyzed using METAL with weighting by sample size.

Gene prioritization methods

Closest gene

We annotated the closest gene to the lipid multi-ancestry index variants [24] by identifying the closest gene transcript on either side (500 kb) of the index variant [92].

Colocalization with GTEx eQTLs

For each of the five lipid phenotypes, we first lifted over GWAS summary statistics from the multi-ancestry meta-analysis [24] to GRCh38 using the UCSC liftOver tool. Then, we defined a set of approximately independent windows across the genome within which colocalization with eQTLs was run. To define these, we first identified all genome-wide significant variants (p -value $< 5e-08$) from the meta-analysis for each lipid trait and sorted them by significance, from most significant to least. Starting with the most significant variant, we aimed to define a window defining independent genetic signals; we define a variant's window as a region within the greater of 500 kb or 0.25 cM on either side of this "sentinel variant." Genetic distances were defined using reference maps from HapMap 3. All other genome-wide significant variants within this window were discarded from the list of sentinel variants, and similar windows were defined for the remaining genome-wide significant variants.

We ran an eQTL colocalization using GTEx v8 eQTL summary statistics within each of our defined windows for all lipid traits. For each of the 49 GTEx tissues, we first identified all genes within 1 Mb of the sentinel variant, and then restricted analysis to those genes with eQTLs ("eGenes") in that tissue (FDR < 0.05). We used the R package "coloc"

(run on R version 3.4.3, coloc version 3.2.1) [93] with default parameters to run colocalization between the GWAS signal and the eQTL signal for each of these cis-eGenes, using as input those variants in the defined window, i.e., all variants present in both datasets. A colocalization posterior probability of $(PP3 + PP4) > 0.8$ was used to identify loci with enough colocalization power, and $PP4/PP3 > 0.9$ was used to define those loci that show significant colocalization [94].

Transcriptome-wide association studies (TWAS)

For our transcriptome-wide association analysis (TWAS), we integrated the results of our GWAS with eQTL summary statistics from GTEx v8. The S-PrediXcan software [95] allows us to integrate these two datasets using only summary statistics from GWAS without needing individual-level genotype data. We used the multi-ancestry lipid GWAS summary statistics [24] and harmonized them with the GTEx summary statistics. Then we performed the TWAS using the eQTL models estimated on GTEx v8 expression data. For each of the 49 GTEx tissues, we identified “significant genes” those genes with p -values more significant than an FDR threshold of 0.05.

Genes with coding variants

We determine the coding variants within 99% credible sets and the coding variants in $LD > 0.8$ with variants in the 99% credible sets with the credible sets as defined here [24]. We define regions for construction of the credible sets as ± 500 kb around each index variant. We used Bayes factors (BFs) for each variant from the MR-MEGA output and generated the credible sets within each region by ranking all variants by BF and calculating the number of variants required to reach a cumulative probability of at least 99%. Additionally, we used previously established gene-based associations [96] to determine whether rare coding variation in a gene were associated with blood lipid levels ($p < 0.001$). We labeled a gene as having coding variants if any of these criteria were met.

DEPICT

We used Data-driven Expression-Prioritized Integration for Complex Traits (DEPICT, v1 beta version rel194 for 1 KG imputed GWAS) to prioritize genes at our index variants, on the assumption that truly associated genes share functional annotations [27]. Index variants [24] with p -value $< 5 \times 10^{-8}$ were retained as input. We implemented the DEPICT analysis with the default settings of 500 permutations for bias adjustment and 20 replications for FDR estimation. DEPICT prioritizes genes by calculating the similarity of a given gene to genes from other associated loci across 14,461 reconstituted gene sets and estimates the nominal gene prioritization p -value and the estimated false discovery rate of each gene. The prioritized genes at $FDR < 0.05$ were considered significant.

PoPS

We used the PoPS method to prioritize genes in the previously reported [24] multi-ancestry index variants for all lipid traits. The PoPS method [28] is a new gene prioritization method that identifies the causal genes by integrating GWAS summary statistics with gene expression, biological pathway, and predicted protein–protein interaction

data. First, as part of the PoPS analysis, we used MAGMA to compute gene association statistics (z -scores) and gene–gene correlations from GWAS summary statistics and LD information from a multi-ancestry reference panel (1000 Genomes). Next, PoPS performs marginal feature selection by using MAGMA to perform enrichment analysis for each gene feature separately. The model is fitted by generalized least squares (GLS), and MAGMA results are used to perform marginal feature selection, retaining only features that pass a nominal significance threshold ($p < 0.05$). Then PoPS computes a joint enrichment of all selected features simultaneously in a leave one chromosome out (LOCO) framework. The gene features employed by PoPS are listed here: https://github.com/FinucaneLab/gene_features. Finally, PoPS computes polygenic priority scores for each gene by fitting a joint model for the enrichment of all selected features. The PoPS score for a gene is independent of the GWAS data on the chromosome where the gene is located. The PoPS analysis returned scores for a total of 18,383 genes per lipid trait. We only kept the top 20% genes among all 18,383 genes. We then annotated our index variants with the nearest ENSEMBL genes in a 500-kb window (either side) and selected the highest PoPS score gene in the locus as the prioritized one.

We performed the PoPS analysis on our lipid-specific multi-ancestry meta-analysis results, using all populations from 1000G as the reference for the LD information in MAGMA. As a sensitivity step, we also repeated the same analysis using only the European population from 1000G as the reference. We observed high concordance in the top two PoPS prioritized genes from both reference panels. In detail, the same 2119 genes (89%) were prioritized as the top genes from both panels, a further 203 genes were prioritized as a top gene with one panel and as the second top with the other and only 7 genes were completely mismatched between the two reference panels.

Monogenic genes

We annotated genes known to cause Mendelian lipid disorders based on proximity with identified GWAS loci [97, 98]. GWAS index variants within ± 500 kb of the transcription start and end positions from the USCS genome browser annotations were annotated as nearby known monogenic dyslipidemia genes.

Mouse knockout lipid phenotype silver set genes

Human gene symbols (9557 unique genes) were mapped to gene identifiers (HGNC) and their corresponding mouse ortholog genes were obtained using Ensembl (www.ensembl.org). Phenotype data for single-gene knockout mouse models were obtained from the International Mouse Phenotyping Consortium (IMPC) (www.mousephenotype.org) latest data release 12.0 (www.mousephenotype.org/data/release). The knockout mouse models were primarily produced by IMPC but also include some models that have been reported from the relevant literature and were curated by Mouse Genome Informatics (MGI) data release 6.16 (www.informatics.jax.org). For each mouse model, reported phenotypes were grouped using the mammalian phenotype ontology hierarchy into broad categories relevant to lipids: growth and body weight (MP:0001259), lipid homeostasis (MP:0002118), cholesterol homeostasis (MP:0005278), and lipid metabolism (MP:0013245). This resulted in mapping of human genes to significant phenotypes in animals.

For each of the multi-ancestry lipid index variant [24], we mapped the closest gene to the knockout mouse phenotypes and curated the set to only include mouse phenotypes strictly relating to lipid metabolism. That resulted in our silver set of 740 genes with mouse lipid phenotypes (Additional file 33: Table S22).

Overlap between methods

We standardized the gene names across different methods using the R/geneSynonym package, a wrapper to gene synonym information in ftp://ftp.ncbi.nlm.nih.gov/gene/DATA/gene_info.gz. We also quantified how often the same gene was prioritized by multiple methods for each index variant and determined scores that ranged from 1 to 6 (S1-S6), based on the number of methods that prioritized the gene.

We integrated multiple gene prioritization methods to identify likely causal genes in the latest global lipid genetics consortium GWAS results. In total, we have implemented the 6 individual gene prioritization methods above that utilize the GWAS summary statistics from meta-analysis. Our gene prioritization methods can be placed into two broad categories, the locus-specific methods and the genome-wide methods. The locus-specific methods leverage local GWAS data by connecting the causal variants to the causal gene(s) using genomic distance, eQTLs, or protein-coding variants.

More specifically, there are four locus-specific methods that have been implemented including: (1) The closest protein-coding gene around the index variants based on the genomic distance, (2) eQTL colocalization using r COLOC package, (3) TWAS using S-PrediXcan, (4) coding variants which have been identified in 99% credible sets OR in $LD > 0.8$ with coding variants OR from gene-based tests ($p < 0.001$) of rare coding variants. For the eQTL and TWAS, we first used all the 49 GTEx tissues and then restricted to only 5 lipid-specific tissues: liver, adipose subcutaneous, adipose visceral, whole blood, and small intestine. In addition, two genome-wide methods were employed: (1) DEPICT ($FDR < 0.05$), (2) PoPS (Top 1 gene). It is reasonable to combine similarity-based methods with locus-based methods since they use two different sources of information.

To determine the relative performance of each prioritization method and their combined scores for lipid loci, we used 21 genes known to cause Mendelian dyslipidemias as a gold standard set (*ABCA1*, *ABCG5*, *ANGPTL3*, *APOA5*, *APOB*, *APOE*, *CETP*, *CYP27A1*, *GPD1*, *GPIHBP1*, *LCAT*, *LDLR*, *LDLRAP1*, *LIPA*, *LIPC*, *LMF1*, *LPL*, *MTTP*, *PCSK9*, *SAR1B*, *SCARB1*), and 740 mouse knockout genes causing lipid phenotypes as a silver standard set (Additional file 33: Table S22). We examined two metrics for each gene prioritization approach: (1) the proportion of prioritized genes in the gold/silver standard set, and (2) the proportion of correctly identified genes among all prioritized genes (Additional file 3: Figure S1). Note that out of the 2286 lipid associations, 97 fell within 500 kb of a Mendelian gene and 1280 within 500 kb of a mouse knockout gene with a lipid phenotype. We observed that the TWAS results yielded a high number of prioritized genes, but lead to a low proportion correctly identified. The TWAS approach had a much smaller proportion of genes correctly prioritized among all the prioritized genes, given it prioritized a total of 3511 genes, which was 3.5-fold greater than the other methods (~1000 genes). Notably, PoPS provided a similar proportion of correctly identified genes (78%) as of TWAS, while retained a high proportion of prioritized genes in the gold standard set (67%). Compared with PoPS, PoPS+ (PoPS plus one of the local

methods) slightly sacrificed the proportion of correctly identified genes from 78 to 71%, but improved the proportion of prioritized genes in the gold standard set from 67% to 79%. Overall, PoPS/PoPS+ outperform other gene prioritization methods on both metrics for our gold (Additional file 3: Figure S1A) and silver (Additional file 3: Figure S1B) standard gene sets. We also assessed lipid-relevant tissue (liver, subcutaneous and visceral adipose, whole blood, and small intestine) expression QTLs (lipid eQTLs) and transcriptome-wide association (lipid TWAS) and found that the expression results from all tissues performed slightly better at recovering the reference gene sets compared with limiting to the lipid-relevant tissues (Additional file 3: Figure S1).

Text mining analysis

We retrieved the whole MEDLINE/PubMed titles and abstracts as of March 06, 2022, from National Library of Medicine (<https://ftp.ncbi.nlm.nih.gov/pubmed/baseline/>; <https://ftp.ncbi.nlm.nih.gov/pubmed/updatefiles/>). We then examined whether a list of genes prioritized by PoPS+ and any one of the lipid-related keywords (lipid, lipids, triglyceride, triglycerides, fatty acid, cholesterol, dyslipidemias, hyperlipidemia, hypercholesteremia, diabetes, type 2 diabetes, type II diabetes, heart, cardiovascular, artery, coronary, coronary artery, coronary heart, atherosclerosis, peripheral vascular, PAD, stroke) occurred in the same abstract. We counted how many lipid-related publications that have a specific gene co-occurred with at least one lipid-related keyword. The same text mining approach was also implemented to a set of randomly selected genes from the 18,383 protein-coding genes used by the PoPS. We estimated the number of lipid-related publications we would expect to see by chance. A Mann–Whitney *U* test was performed to show whether there was a significant difference between the number of lipid-related publications of the PoPS+ gene set and reference gene set.

Drug target mining analysis

To gain therapeutic insights from our gene prioritization results, we performed a lookup in Therapeutic Target Database (TTD) 2022 [99] (<http://db.idrblab.net/ttd/>). Specifically, we cross-referenced 466 unique lipid-associated genes prioritized by PoPS+ (Additional file 2: Table S2) with 1563 genes corresponding to at least one drug (either under development or approved) with known clinical indication in TTD 2022. As a quality control for this lookup, we excluded all TTD entries related to drugs that were discontinued, terminated, or withdrawn from the market. The full lookup results are available in Additional file 8: Table S6.

Driver tissues for lipid levels

We performed phenotype-tissue association analysis using DESE (driver-tissue estimation by selective expression) [40]. DESE estimates the causal tissues by selective expression of phenotype-associated genes in GWAS. We used the GWAS summary statistics from the five lipid traits and the GTEx v8 normalized gene-level and transcript-level expression datasets as input. SNPs inside a gene and its ± 5 kb adjacent regions were first mapped to the gene, and then DESE ran iteratively to produce a list of driver tissues and the corresponding *p*-values of the associations. We used a Bonferroni-corrected significance threshold of $0.05/54 = 9.3 \times 10^{-4}$.

PheWAS analysis

Construction of lipid PGSs

We had previously developed a multi-ancestry PGS for LDL-C that was demonstrated to perform well across multiple ancestry groups [24]. In a similar manner, we also generated PGS for HDL-C, nonHDL-C, TC, and triglycerides. First, multi-ancestry meta-analysis results were generated with METAL [89] after excluding individuals from the Michigan Genomics Initiative and the UK Biobank. The set of variants used to construct the PGS was limited to those that were well-imputed ($R^2 > 0.3$) in MGI, UK Biobank, and MVP. Risk scores based on PRS-CS [100] or pruning and thresholding with Plink [101] across several r^2 (0.1, 0.2), distance (250 kb, 500 kb), and p -value thresholds (5×10^{-10} , 5×10^{-9} , 5×10^{-8} , 5×10^{-7} , 5×10^{-6} , 5×10^{-5} , 5×10^{-4} , 5×10^{-3} , 0.05) were developed. For each trait, the single best score was selected based on the adjusted r^2 calculated in the UK Biobank of the linear model for the lipid trait with the risk score and age, sex, batch, and PC1-4 as covariates. This corresponded to PRS-CS for HDL-C and nonHDL-C and pruning and thresholding for LDL-C ($r^2 = 0.1$, p -value = 5×10^{-4} , 500 kb), TG ($r^2 = 0.1$, p -value = 5×10^{-3} , 500 kb), and TC ($r^2 = 0.1$, p -value = 5×10^{-4} , 500 kb). The variance explained by the risk score among the UK Biobank participants was similar across traits (adjusted r^2 of the full model-adjusted r^2 of covariates: HDL-C = 0.13; LDL-C = 0.15; nonHDL-C = 0.14; TC = 0.14; TG = 0.10) and validated the ability of the risk score to predict genetically increased lipid levels.

PheWAS of lipid PGSs and index lipid variants in the UK Biobank and MVP

We used the European ancestry subset of individuals from the UK Biobank (408,886 samples) and the European samples from MVP (69,670 samples) to perform the PheWAS analysis.

We constructed a weighted PGS for each of the lipid traits, based on the corresponding genome-wide significant multi-ancestry index variants. We used the PheWAS package in R [102] to map ICD-10 codes from hospital records into clinically relevant phenotypes (phecodes) and to implement these association analyses, while adjusting for sex, age, 10 genetic principal components, and genotyping array (for the UK Biobank only) in each cohort. For the lipid-PGS PheWAS, each PGS was inverse normalized prior to analysis and lipid levels were corrected for statin use. The MVP samples used for the PheWAS analysis were not included in the GWAS meta-analysis [24].

Similarly, we extracted all multi-ancestry autosomal index variants for all lipid traits from the same European ancestry subset of the UK Biobank and MVP and performed a single-variant PheWAS association analysis per cohort. Additionally, we performed a single-variant PheWAS association analysis in the UK Biobank only with the sex-stratified and X chromosome index variants from the multi-ancestry analysis.

Meta-analysis of MVP and the UK Biobank PheWAS results

We combined, via meta-analysis, PheWAS lipid-specific PGS results for all intersecting phecodes and biomarkers between the UK Biobank and MVP (Europeans only) per lipid trait. We used ICD10-based phecodes and manually matched biomarkers to identify intersecting phenotypes between the two datasets. We restricted our meta-analysis to phenotypes that had at least 100 samples (total number for continuous traits or

number of cases for binary traits) in each cohort. After the meta-analysis, we excluded phenotypes that had less than 500 combined samples (total number for continuous traits or number of cases for binary traits), to avoid reporting spurious results [103]. That resulted in a total of 773 phenotypes (739 phecodes and 34 biomarkers/measurements). We used both fixed and random effects model for the meta-analysis. We assessed heterogeneity using the p -value for Cochran's q and set the level for significant heterogeneity at a Bonferroni threshold (p -value $\leq 6.5 \times 10^{-5}$, to account for multiple testing of 773 phenotypes). We report the results from the fixed-effects model for the phenotypes with non-significant heterogeneity and the results from the random effects model for all others. Similarly, we meta-analyzed all index-variant PheWAS results between the UK Biobank and MVP and obtained results for 811 phenotypes and 1750 lipid multi-ancestry index variants, after excluding instances with a combined sample size < 500 .

Lipid index variants with CAD, T2D, and NAFLD datasets

The GWAS meta-analysis results of CAD and T2D were acquired from MVP [62] and DIAGRAM Consortium [61], respectively. For variant rs1229984, the CAD result is from CARDIoGRAMPlusC4D meta-analysis [104], as it was not present in the MVP results. The NAFLD GWAS and meta-analysis was performed in the UK Biobank and Michigan Genomics Initiative (MGI). We determined the association of the lipid index variants with CAD, T2D, and NAFLD and aligned the alleles across all the traits to the LDL-lowering allele. We then highlighted the protective lipid coding alleles associated with CAD.

GWAS and meta-analysis of NAFLD in the UK Biobank and Michigan Genomics Initiative (MGI)

Individuals with NAFLD were identified using ICD-9 571.8 and ICD-10 K76.0. Individuals with hepatitis, liver cirrhosis, liver abscess, ascites, a liver transplant, hepatomegaly, jaundice, or with abnormal result of serum enzyme levels or a function study of the liver were excluded (exclusion phecodes 70.2, 70.3, 571.51, 571.6, 571.8, 571.81, 572, 573, 573.2, 573.3, 573.5, 573.7, 573.9) [105]. Analysis was performed using SAIGE v43.3 [106]. Analysis in the UK Biobank included white British individuals with batch, sex, birth year, and the first 4 genetic principal components as covariates. A total of 1122 cases and 399,900 controls were included in the analysis. Analysis in MGI included only European-ancestry participants with array version, sex, birth year, and the first 4 genetic principal components as covariates. A total of 2901 cases and 49,098 controls were analyzed. Meta-analysis was performed using METAL with weighting based on the effective sample size calculated as $4/((1/N_{\text{cases}}) + (1/N_{\text{controls}}))$.

CAD/T2D colocalization analysis with lipid traits

We used R package coloc v3.2.1 [93] to perform summary statistics-based colocalization via a Bayesian approach and test whether the 5 lipid traits share common genetic causal variants with CAD or T2D. We first defined a window of ± 100 kb around each index variant [24]. Then for each window of the 10 pairs of traits, we ran colocalization with default parameters using those SNPs present in both datasets. A colocalization posterior probability of $PP4 > 0.8$ was used to define those loci that show significant colocalization.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13059-022-02837-1>.

- Additional file 1: Table S1.** Characteristics of contributing cohorts (as provided by each participating cohort).
- Additional file 2: Table S2.** Association results for the multi-ancestry index SNPs with the gene prioritization.
- Additional file 3: Figure S1.** Summary of prioritizing genes for A. Mendelian and B. mouse model genes separately by trait.
- Additional file 4: Table S3.** Text mining results for the PoPS+ prioritized genes.
- Additional file 5: Table S4.** Frequency of lipid-related publications for the PoPS+ prioritized genes.
- Additional file 6: Figure S2.** Frequency distribution of the lipid-related publications for both high confidence genes and the baseline genes.
- Additional file 7: Table S5.** Lookup of all prioritized lipid genes in the Therapeutic Target Database 2022.
- Additional file 8: Table S6.** Association of lipid index variants with CAD, T2D and NAFLD.
- Additional file 9: Figure S3.** Lipid traits – tissue/cell type associations estimated by DESE according to GTEx gene-level and GTEx transcript-level selective expression.
- Additional file 10: Table S7.** DESE phenotype-tissue association results using both GTEx gene-level and transcript-level selective expression.
- Additional file 11: Figure S4.** Comparison of PheWAS results in UKB and MVP for the LDL-C PGS, HDL-C PGS, TC PGS, TG PGS and nonHDL-C PGS.
- Additional file 12: Table S8.** PheWAS UKB-MVP meta-analysis results for each lipid PGS.
- Additional file 13: Figure S5.** PheWAS meta-analysis results for the trans-ethnic HDL-C PGS in UK Biobank and MVP.
- Additional file 14: Figure S6.** PheWAS meta-analysis results for the trans-ethnic TC PGS in UK Biobank and MVP.
- Additional file 15: Figure S7.** PheWAS meta-analysis results for the trans-ethnic TG PGS in UK Biobank and MVP.
- Additional file 16: Figure S8.** PheWAS meta-analysis results for the trans-ethnic nonHDL-C PGS in UK Biobank and MVP.
- Additional file 17: Table S9.** PheWAS UKB-MVP meta-analysis results for each index lipid variant at Bonferroni threshold for multiple testing $p \leq 3.5e-8$
- Additional file 18: Table S10.** Lambda GC values across minor allele frequency bins for sex-specific meta-analyses.
- Additional file 19: Table S11.** Significant female-specific multi-ancestry meta-analysis results.
- Additional file 20: Table S12.** Significant male-specific multi-ancestry meta-analysis results.
- Additional file 21: Table S13.** Characteristics of replication cohorts (as provided by each participating cohort).
- Additional file 22: Table S14.** Test for difference in effects for index variants from sex-stratified meta-analysis.
- Additional file 23: Table S15.** Comparison of the sex-specific effects.
- Additional file 24: Figure S9.** Comparison of effect size estimates between males and females for index variants showing a significant difference in effect size between sexes.
- Additional file 25: Table S16.** Comparison of effect size estimates for sex-stratified analysis in the replication cohorts.
- Additional file 26: Figure S10.** Comparison of effect sizes for trans-ancestry index variants excluding cholesterol-lowering medication.
- Additional file 27: Table S17.** Sex-stratified effect sizes in UK Biobank considering all individuals or only those not on cholesterol lowering medications.
- Additional file 28: Table S18.** Sex-participation association of the variants with significant sex-specific lipid results.
- Additional file 29: Table S19.** Comparison of sex-stratified effect sizes in the UK Biobank for BMI, waist hip ratio adjusted for BMI, alcohol use, and smoking status.
- Additional file 30: Table S20.** Colocalization results for the sex-specific loci.
- Additional file 31: Supplementary Note.** Supplementary Note and Cohort Acknowledgments.
- Additional file 32: Table S21.** Significant X chromosome results in the sex-combined and sex-stratified analysis and replication.
- Additional file 33: Table S22.** Mouse genes with lipid phenotypes (silver set).
- Additional file 34. Review history.

Acknowledgements

We thank Bethany Klunder for her administrative support of the Global Lipids Genetics Consortium. Study-specific acknowledgements are available in the Additional file 32: Supplementary Note.

Peer review information

Wenjing She was the primary editor of this article and managed its editorial process and peer review in collaboration with the rest of the editorial team.

Review history

The review history is available as Additional file 34.

Authors' contributions

S.Kanoni, S.E.G. Y.W., I.S., S.Ramdas, and Xiang.Zhu contributed equally to this work as co-first authors. All authors reviewed the manuscript. Consortium management: G.M.P., P.N., T.L.A., M.Boehnke, and C.J.W. Study design, interpretation of results, and drafting of the manuscript: S.Kanoni, S.E.G. Y.W., I.S., S.Ramdas, Xiang.Zhu, S.L.C., K.F.B., S.Vedantam, T.W.W., A.E.L., E.M., G.J.M.Z., K-H.H.W., I.N., Y.V.S., A.P.M., M.Boehnke, C.D.B., P.N., P.D., C.J.W., T.L.A., and G.M.P. Primary analyses: S.Kanoni, S.E.G. Y.W., I.S., S.Ramdas, Xiang.Zhu, S.L.C., K.F.B., S.Vedantam, T.W.W., A.E.L. Individual study design, analysis, and oversight: S.Kanoni, S.E.G. Y.W., I.S., S.Ramdas, Xiang.Zhu, S.L.C., K.F.B., S.Vedantam, T.W.W., A.E.L., E.M., G.J.M.Z., K-H.H.W., I.N., Q.H., D.K., A.T.Hilliard, Zeyuan.Wang, C.X., G.Thorleifsson, A.H., D.F.G., H.Holm, I.O., M.Y.H., S.Han, M.Akiyama, S.S., C.Terao, M.Kanai, W.Zhou, B.M.B., H.R., A.S.H., Y.V., J.A.P., E.A.R., T.Lingren, Q.P.F., I.J.K., A.N., J.Takayama, H.C.M., K.A.H., B.T., J.Haessler, F.G., Y.B., J.E.M., A.Campbell, K.L., I.Y.M., A.R., G.Hindy, J.D.F., W.Zhao, D.R.W., C.Turman, H.Huang, M.Graff, A.Choudhury, D.Sengupta, A.Mahajan, M.R.B., W.Zhang, K.Yu, E.M.S., A.Pandit, S.G., X.Y., J.Luan, J-H.Z., F.Matsuda, H-M.J., Yoon, C.M-G., A.Pitsillides, J.J.H., A.R.Wood, Y.J., Z.G., S.Haworth, R.E.M., J.F.C., M.Aadahl, A.A.B., J.Yao, A.Manichaikul, C-M.H., Y-J.H., H.R.W., J.R., J-B-J., L.L.K., A.G., M.S-L., R.N., P.Mauro, F.Matteo, A.F.McD., P.M-V., M.Wielscher, S.T., N.S., L.T.M., M.Munz, L.Z., J.Huang, B.Y., A.Poveda, A.K., C.Lamina, L.F., M.S., T.E.G., J.P.B., S.E.R., E.W.D., J.M.Z., J.S.M., C.Fuchsberger, H.Christensen, J.A.B., M.V-M., M.F.F., M.K.W., Zhe.Wang, M.H.P., M.Mangino, P.C., N.V., J.W.B., J.Engmann, N.L.T., A.V., R.C.S., K.S.L., N.R.Z., P.L., M.E.K., G.E.D., S.Huo, D.D.I., H.I., Jian.Yang, Jun.Liu, A.D., H.L.L., J.M., M.Frank, B.S., L.J.S., M.C.G., C.W., M.Nakatochi, A.W., N.H-K., X.S., R.X., A.H-C., J.C.F-L., V.L., S.S.N., S.Bayyana, H.M.S., M.R.I., C.Oldmeadow, H-N.K., S.Ryu, P.R.H.T., L.A., R.D., L.A.L., G.Prasad, L-L-M., M.P., J.Long, X.Li, E.T., F.T., C.N.S., A.Loukola, S.Bollepalli, S.C.W., Y.X.W., W.B.W., T.Nutile, D.R., Y.J.S., S.C., F.L., Jingyun.Yang, K.A.K., B.B., G.G.N., K.M., L.F.B., J.A.S., P.H., A-E.F., E.H., M.Lin, M.P.C., S.Vaccargiu, P.J.van der M., N.Pitkänen, B.E.C., S.W.van der L., K.N.C., S.W., A.R.B., A.P.D., A.A.A., J.Y.L., E.R.B.P., A.N., H.S.C., M.Nethander, S.F-W., L.S., N.W.R., C.A.W., S-Y.L., J-S.W., C.C., L-P.L., K.N., G.C-P, H.Vestergaard, B.H., O.G., Q.C., M.O.O., J.van S., J.Liang, H.T., N.T., J.H.S., R.D.J., A.P.R., L.W.M., Z.C., L.Li, T.Kawaguchi, J.Thiery, J.C.B., L.J.L., Huaixing.Li, M.A.N., O.T.R., S.I., S.H.W., C.P.N., H.Campbell, S.J., T.Nabika, F.A-M., H.N., P.S.B., I.K., P.K., T.G., T.Katsuya, D.de K., Gert J.de B., E.K.K., H.H.H.A., M.A.I., Xiaofeng.Zhu, F.W.A., A.O.K., J.W.J.B., X-O.S., L.S.R., O.Pedersen, T.H., P.Mitchell, A.W.H., M.Kähönen, L.P., C.Bouchard, A.T., Y-D. I.C., C.E.P., T.A.M., W.L., A.F., C.Ohlsson, D.M., Y.S.C., H.Lee, J-M.Y., W-P.K., S.Y.R., J-T.W., I.M.H., K.J.S., M.E.Z., H.Völzke, G.Homuth, M.K.E., A.B.Z., O.Polasek, G.Pasterkamp, I.E.H., S.Redline, K.P., A.J.O., H.Snieider, G.B., R.S., H.Schmidt, S.Bandinelli, G.D., T.A.T., S.L.R.K., P.A.P., N.K., M.B.S., G.G., C.A.B., B.J., P.K.J., D.A.B., P.L.De J., X.Lu, V.M., M.Brown, M.J.C., P.B.M., X.G., M.Ciullo, J.B.J., N.J.S., J.Kaprio, P.P., T.T-L., C.A.A-S., L.S.A., S.A.B., H. J.de S., A.R.Wickremasinghe, R.M.K., J-Y.W., W.Zeng, A.I.den H., D.B., A.Correa, J.G.W., L.Lind, C-K.H., A.E.N., Y.M.G., J.F.W., B.P., H-L.K., J.A., R.J.S., D.C.R., D.K.A., M.Walker, H.A.K., G.R.C., J.M.M., M.C.C., D.J., N.P.B., C.G.V., L.O., M.Fornage, E S.T., R.M.van D., T.Lehtimäki, N.C., M.Yokota, Jianjun.Liu, D.F.R., A.J.McK., J.Kee, K-H.J., M.I.McC., C.N.A.P., V.V., C.H., E.S., C.M.van D., Z-B.J., J.Q., H.Hishigaki, X.Lin, W.M., V.G., J-C.T., G.L., L.M.t H., P.J.M.E., S.M.D., M.Kumari, M.Kivimäki, P.van der H., T.D.S., R.J.F.L., M.A.P., E.J.P., M.Cruz, B.M.P., I.B., P.P.P., C.N.R., K.Christensen, S.Ripatti, E.W., H.Hakonarson, S.F.A.G., L.A.L.M.K., J.de G., M.Loeffler, F.Kronenberg, D.G., J.Erdmann, H.Schunkert, P.W.F., A.Linneberg, J.W.J., A.V.K., M.Männikkö, M-R.J., Z.K., C.Francesco, D.O.M-K., K.W.van D., H.W., D.P.S., N.G., P.S., N.Poulter, L-M.C., J.I.R., T.M.D., F.Karpe, M.J.N., N.J.T., C-Y.C., T-Y.W., C.C.K., Hengtong.Li, C.S., A.Peters, C.G., A.T.Hattersley, N.L.P., P.K.E.M., D.I.B., A.H.M.W., L.A.C., J.B.J.van M., M.Ghanbari, P.G-L., W.H., Y.J.K., Y.T., N.J.W., Langenberg, E.Z., J.Kuusisto, M.Laakso, E.I., G.A., J.C.C., J.S.K., P.S.de V., A.C.M., S.Hazellhurst, M.R., K.E.N., M.D., P.K., N.G.M., J.B.W., S.A., D.Saleheen, R.G.W., M.V.H., C.Black, B.H.S., A.B., A.E.J., J.E.B., P.M.R., D.I.C., C.K., G.Tamiya, M.Yamamoto, D.A.van H., R.C.T., W-Q.W., G.P.J., B.N., M.G.H., M.D.R., P.J., V.S., K.H., B.O.A., M.Kubo, Y.K., Y.O., Y.M., B-J.K., U.T., K.S., J.Z., Y.E.C., Y-L.H., J.A.L., D.J.R., P.S.T., K-M.C., K.Cho, C.J.O'D., J.M.G., P.W.F.W., T.M.F., J.N.H., S.Kathiresan, K.L.M., Y.V.S., A.P.M., M.Boehnke, C.D.B., P.N., P.D., C.J.W., T.L.A., and G.M.P. All author(s) read and approved the final manuscript.

Funding

GMP, PN, and CW are supported by NHLBI R01HL127564. GMP and PN are supported by R01HL142711. AG acknowledges support from the Wellcome Trust (201543/B/16/Z), European Union Seventh Framework Programme FP7/2007–2013 under grant agreement no. HEALTH-F2-2013–601456 (CVGenes@Target) & the TriPartite Immunometabolism Consortium [TriC]-Novo Nordisk Foundation's Grant number NNF15CC0018486. JMM is supported by American Diabetes Association Innovative and Clinical Translational Award 1–19-ICTS-068. SR was supported by the Academy of Finland Center of Excellence in Complex Disease Genetics (Grant No 312062), the Finnish Foundation for Cardiovascular Research, the Sigrid Juselius Foundation, and University of Helsinki HiLIFE Fellow and Grand Challenge grants. EW was supported by the Finnish innovation fund Sitra (EW) and Finska Läkaresällskapet. CNS was supported by American Heart Association Postdoctoral Fellowships 15POST24470131 and 17POST33650016. Charles N Rotimi is supported by 201HG200362. Zhe Wang, Michael H Preuss, and Ruth JF Loos are supported by R01HL142302. NJT is a Wellcome Trust Investigator (202802/Z/16/Z), is the PI of the Avon Longitudinal Study of Parents and Children (MRC & WT 217065/Z/19/Z), is supported by the University of Bristol NIHR Biomedical Research Centre (BRC-1215–2001) and the MRC Integrative Epidemiology Unit (MC_UU_00011), and works within the CRUK Integrative Cancer Epidemiology Programme (C18281/A19169). Ruth E Mitchell is a member of the MRC Integrative Epidemiology Unit at the University of Bristol funded by the MRC (MC_UU_00011/1). Simon Haworth is supported by the UK National Institute for Health Research Academic Clinical Fellowship. Paul S. de Vries was supported by American Heart Association grant number 18CDA34110116. Julia Ramierz acknowledges support by the People Programme of the European Union's Seventh Framework Programme grant n° 608765 and Marie Skłodowska-Curie grant n° 786833. Maria Sabater-Lleal is supported by a Miguel Servet contract from the ISCIII Spanish Health Institute (CP17/00142) and co-financed by the European Social Fund. Jian Yang is funded by the Westlake Education Foundation. Olga Giannakopoulou has received funding from the British Heart Foundation (BHF) (FS/14/66/3129). CHARGE Consortium cohorts were supported by R01HL105756. Study-specific acknowledgements are available in the Additional file 32: Supplementary Note. The views expressed in this manuscript are those of the authors

and do not necessarily represent the views of the National Heart, Lung, and Blood Institute; the National Institutes of Health; or the U.S. Department of Health and Human Services.

Availability of data and materials

The GWAS meta-analysis results (including both ancestry-specific and trans-ancestry analyses) and risk score weights are available at: <http://csg.sph.umich.edu/willer/public/glgc-lipids2021> [107]. A web browser displaying the gene prioritization and PheWAS results is available at https://hugeamp.org:8000/research.html?pageid=GLGC_149 [108]. The optimized trans-ancestry polygenic score weights are deposited within the PGS Catalog (<https://www.pgscatalog.org/publication/PGP000230/> [109] and <https://www.pgscatalog.org/publication/PGP000366/> [110]). Scripts used for analysis and summary of results are available under the MIT license on this GitHub repository: <https://github.com/Global-Lipids-Genetics> [111]. The version of source code used in the manuscript is deposited in Zenodo: <https://doi.org/10.5281/zenodo.7130299> [112].

Declarations

Ethics approval and consent to participate

The overall study was approved by the IRB of the Boston University Medical Center. Individual studies were approved by the appropriate institutional review boards (IRB) and informed consent was obtained from all participants.

Competing interests

Ioanna Ntalla is an employee and stock owner of Gilead Sciences since August 2019. Derek Klarin accepts consulting fees from Regeneron Pharmaceuticals. All deCODE-affiliated authors (Gudmar Thorleifsson, Anna Helgadóttir, Daniel F Gudbjartsson, Hilma Holm, Unnur Thorsteinsdóttir, Kari Stefánsson) are employees of deCODE/Amgen Inc. As of January 2020, Anubha Mahajan is an employee of Genentech, and a holder of Roche stock. Markus Scholz receives funding from Pfizer Inc. for a project not related to this research. Marcus E Kleber is employed by SYNLAB MVZ Mannheim GmbH. Gabriel Cuellar-Partida contributed to this work while employed at The University of Queensland, but he is now an employee of 23andMe Inc. Mark J Caulfield is Chief Scientist for Genomics England, a UK Government company. The views expressed in this article are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health. Mark I McCarthy has served on advisory panels for Pfizer, NovoNordisk, and Zoe Global and has received honoraria from Merck, Pfizer, Novo Nordisk, and Eli Lilly and research funding from Abbvie, Astra Zeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Merck, NovoNordisk, Pfizer, Roche, Sanofi Aventis, Servier, and Takeda. As of June 2019, Mark I McCarthy is an employee of Genentech, and a holder of Roche stock. Winfried März has received grants from Siemens Healthineers, grants and personal fees from Aegerion Pharmaceuticals, grants and personal fees from AMGEN, grants from Astrazeneca, grants and personal fees from Sanofi, grants and personal fees from Alexion Pharmaceuticals, grants and personal fees from BASF, grants and personal fees from Abbott Diagnostics, grants and personal fees from Numares AG, grants and personal fees from Berlin-Chemie, grants and personal fees from Akzea Therapeutics, grants from Bayer Vital GmbH, grants from bestbion dx GmbH, grants from Boehringer Ingelheim Pharma GmbH Co KG, grants from Immundiagnostik GmbH, grants from Merck Chemicals GmbH, grants from MSD Sharp and Dohme GmbH, grants from Novartis Pharma GmbH, grants from Olink Proteomics, other from Synlab Holding Deutschland GmbH, all outside the submitted work. Bruce M Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson. Amit V Khera has served as a consultant to Sanofi, Medicines Company, Maze Pharmaceuticals, Navitor Pharmaceuticals, Verve Therapeutics, Amgen, and Color Genomics; received speaking fees from Illumina, the Novartis Institute for Biomedical Research; received sponsored research agreements from the Novartis Institute for Biomedical Research and IBM Research, and reports a patent related to a genetic risk predictor (20190017119). Dennis O Mook-Kanamori is a part-time clinical research consultant for Metabolon, Inc. Danish Saleheen has received support from the British Heart Foundation, Pfizer, Regeneron, Genentech, and Eli Lilly pharmaceuticals. Veikko Salomaa has received honoraria for consultations from Novo Nordisk and Sanofi and has ongoing research collaboration with Bayer Ltd, all unrelated to the present study. Sekar Kathiresan is an employee of Verve Therapeutics, and holds equity in Verve Therapeutics, Maze Therapeutics, Catabasis, and San Therapeutics. He is a member of the scientific advisory boards for Regeneron Genetics Center and Corvidia Therapeutics; he has served as a consultant for Acceleron, Eli Lilly, Novartis, Merck, Novo Nordisk, Novo Ventures, Ionis, Alnylam, Aegerion, Haug Partners, Noble Insights, Leerink Partners, Bayer Healthcare, Illumina, Color Genomics, MedGenome, Quest, and Medscape; he reports patents related to a method of identifying and treating a person having a predisposition to or afflicted with cardiometabolic disease (20180010185) and a genetics risk predictor (20190017119). Cristen J Willer's spouse is employed by Regeneron.

Author details

¹William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, UK. ²Department of Internal Medicine, Division of Cardiology, University of Michigan, Ann Arbor, MI 48109, USA. ³Department of Biostatistics, Boston University School of Public Health, 801 Massachusetts Ave, Boston, MA 02118, USA. ⁴Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA. ⁵Department of Statistics, The Pennsylvania State University, University Park, PA, USA. ⁶Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA, USA. ⁷VA Palo Alto Health Care Systems, Palo Alto, CA, USA. ⁸Department of Statistics, Stanford University, Stanford, CA, USA. ⁹Department of Medicine, Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA. ¹⁰Boston Children's Hospital, Endocrinology, Boston, MA 02115, USA. ¹¹Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA. ¹²Department of Genetic Epidemiology, University of Regensburg, Regensburg, Germany. ¹³McDonnell Genome Institute and Department of Medicine, Washington University, St. Louis, MO 63108, USA. ¹⁴Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA. ¹⁵Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA. ¹⁶Clinical Pharmacology, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ¹⁷Department of Epidemiology, Emory University Rollins

School of Public Health, Atlanta, GA, USA. ¹⁸Atlanta VA Health Care System, Decatur, GA, USA. ¹⁹Department of Surgery, Stanford University School of Medicine, Stanford, CA, USA. ²⁰deCODE Genetics/Amgen, Inc. Sturlugata 8, Reykjavik 102, Iceland. ²¹School of Engineering and Natural Sciences, University of Iceland, Sæmundargötu 2, Reykjavik 102, Iceland. ²²Department of Clinical Biochemistry, Landspítali - National University Hospital of Iceland, Hringbraut, Reykjavik 101, Iceland. ²³Division of Genome Science, Department of Precision Medicine, National Institute of Health, Chungcheongbuk-Do, South Korea. ²⁴Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. ²⁵Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan. ²⁶Department of Statistical Genetics, Osaka University Graduate School of Medicine, Osaka, Japan. ²⁷Department of Allergy and Rheumatology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan. ²⁸Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. ²⁹Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA. ³⁰Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA. ³¹K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim, Norway. ³²MRC Integrative Epidemiology Unit (IEU), Bristol Medical School, University of Bristol, Oakfield House, Oakfield Grove, Bristol BS8 2BN, UK. ³³Clinic of Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. ³⁴Division of Medicine and Laboratory Sciences, University of Oslo, Oslo, Norway. ³⁵Institute for Molecular Medicine Finland (FIMM), HILIFE, University of Helsinki, Tukholmankatu 8, 00014 Helsinki, Finland. ³⁶Department of Public Health and Welfare, Finnish Institute for Health and Welfare, Helsinki, Finland. ³⁷Department of Genetics, Institute for Biomedical Informatics, University of Pennsylvania, Perelman School of Medicine, Philadelphia, PA 19104, USA. ³⁸Center for Genetic Medicine, Northwestern University, Feinberg School of Medicine, Chicago, IL 60618, USA. ³⁹Department of Medicine (Medical Genetics), University of Washington, Seattle, WA, USA. ⁴⁰Division of Biomedical Informatics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA. ⁴¹Division of Clinical Pharmacology, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA. ⁴²Department of Cardiovascular Medicine and the Gonda Vascular Center, Mayo Clinic, Rochester, MN, USA. ⁴³Tohoku Medical Megabank Organization, Tohoku University, Sendai 980-8573, Japan. ⁴⁴Wellcome Trust Sanger Institute, Hinxton CB10 1SA, UK. ⁴⁵Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK. ⁴⁶Fred Hutchinson Cancer Center, Division of Public Health Sciences, Seattle, WA 9810, USA. ⁴⁷Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA 02215, USA. ⁴⁸Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, UK. ⁴⁹Usher Institute, The University of Edinburgh, Nine, Edinburgh Bioquarter, 9 Little France Road, Edinburgh EH16 4UX, UK. ⁵⁰Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Oxford OX3 7LF, UK. ⁵¹Medical Research Council Population Health Research Unit, Nuffield Department of Population Health, University of Oxford, Oxford OX3 7LF, UK. ⁵²Center for Non-Communicable Diseases, Karachi, Sindh, Pakistan. ⁵³Department of Population Medicine, Qatar University College of Medicine, QU Health, Doha, Qatar. ⁵⁴Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI 48104, USA. ⁵⁵Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI 48109, USA. ⁵⁶Program in Genetic Epidemiology and Statistical Genetics, Department of Epidemiology, Harvard T.H. Chan School of Public Health, 677 Huntington Avenue, Boston, MA 02115, USA. ⁵⁷Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA. ⁵⁸Sydney Brenner Institute for Molecular Bioscience, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa. ⁵⁹Wellcome Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK. ⁶⁰Human Genetics Center, Department of Epidemiology, School of Public Health, Human Genetics, and Environmental Sciences, The University of Texas Health Science Center at Houston, Houston, TX 77030, USA. ⁶¹Department of Epidemiology and Biostatistics, Imperial College London, London W2 1PG, UK. ⁶²Department of Cardiology, Ealing Hospital, London North West University Healthcare NHS Trust, Middlesex UB1 3HW, UK. ⁶³Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden. ⁶⁴MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge CB2 0QQ, UK. ⁶⁵Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge CB1 8RN, UK. ⁶⁶Center for Genomic Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan. ⁶⁷Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands. ⁶⁸Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands. ⁶⁹Amsterdam Public Health Research Institute, Amsterdam UMC, the Netherlands. ⁷⁰Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, Exeter EX2 5DW, UK. ⁷¹Nanjing University of Chinese Medicine, Nanjing 210029, Jiangsu, China. ⁷²Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany. ⁷³Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany. ⁷⁴Bristol Dental School, University of Bristol, Lower Maudlin Street, Bristol BS1 2LY, UK. ⁷⁵Department of Genetic Medicine, Weill Cornell Medicine-Qatar, Doha, Qatar. ⁷⁶Department of Computer and Systems Engineering, Alexandria University, Alexandria, Egypt. ⁷⁷Population Health Sciences, Bristol Medical School, University of Bristol, Oakfield Grove, Bristol BS8 2BN, UK. ⁷⁸Saw Swee Hock School of Public Health, National University of Singapore and National University Health System, Singapore 117549, Singapore. ⁷⁹Center for Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark. ⁸⁰Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. ⁸¹The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, Lundquist Institute for Biomedical Innovations (Formerly LABioMed) at Harbor-UCLA Medical Center, Torrance, CA 90502, USA. ⁸²Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22903, USA. ⁸³Section of Endocrinology and Metabolism, Department of Medicine, Taipei Veterans General Hospital, Taipei, Taiwan. ⁸⁴Institute of Preventive Medicine, National Defense Medical Center, Postbox 90048~700, Sanhsia Dist, New Taipei City 237101, Taiwan. ⁸⁵William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, John Vane Science Centre, Charterhouse Square, London EC1M 6BQ, UK. ⁸⁶NIHR Barts Cardiovascular Biomedical Research Centre, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ⁸⁷Aragon Institute of Engineering Research, University of Zaragoza and Centro de Investigación Biomédica en Red - Bioingeniería, Biomateriales Y Nanomedicina, Spain. ⁸⁸Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health

and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. ⁸⁹Division of Cardiovascular Medicine, Radcliffe Department of Medicine, John Radcliffe Hospital, University of Oxford, Oxford OX3 9DU, UK. ⁹⁰Unit of Genomics of Complex Diseases, Institut d'Investigació Biomèdica Sant Pau (IIB SANT PAU), Barcelona, Spain. ⁹¹Cardiovascular Medicine Unit, Department of Medicine, Karolinska Institutet, Center for Molecular Medicine, Karolinska University Hospital, Stockholm, Sweden. ⁹²Department of Internal Medicine, Section of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands. ⁹³Istituto Di Ricerca Genetica E Biomedica, Consiglio Nazionale Delle Ricerche, Rome, Italy. ⁹⁴Dipartimento Di Scienze Biomediche, Università Degli Studi Di Sassari, Sardinia, Italy. ⁹⁵University Center for Primary Care and Public Health, University of Lausanne, Rte de Berne 113, 1010 Lausanne, Switzerland. ⁹⁶Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland. ⁹⁷Department of Medicine, Internal Medicine, Lausanne University Hospital and University of Lausanne, Rue du Bugnon 46, 1011 Lausanne, Switzerland. ⁹⁸Department of Epidemiology and Biostatistics, MRC-PHE Centre for Environment and Health, School of Public Health, Imperial College London, London, UK. ⁹⁹Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands. ¹⁰⁰Dept of Internal Medicine, Section of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands. ¹⁰¹BHF Glasgow Cardiovascular Research Centre, Faculty of Medicine, Glasgow, UK. ¹⁰²Institute for Cardiogenetics, University of Lübeck, DZHK (German Research Centre for Cardiovascular Research), Partner Site Hamburg/Lübeck/Kiel, University Heart Center Lübeck, Lübeck and Charité – University Medicine Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität Zu Berlin and Berlin Institute of Health, Institute for Dental and Craniofacial Sciences, Department of Periodontology and Synoptic Dentistry, Berlin, Germany. ¹⁰³Deutsches Herzzentrum München, Klinik Für Herz- Und Kreislauferkrankungen, Technische Universität München, Munich, Germany. ¹⁰⁴Deutsches Zentrum Für Herz-Kreislauf-Forschung (DZHK) E.V., Partner Site Munich Heart Alliance, Munich, Germany. ¹⁰⁵Key Laboratory of Cardiovascular Epidemiology and Department of Epidemiology, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100037, China. ¹⁰⁶Lund University Diabetes Center, Lund University, Malmö, Sweden. ¹⁰⁷Institute of Genetic Epidemiology, Department of Genetics, Medical University of Innsbruck, Innsbruck, Austria. ¹⁰⁸German Chronic Kidney Disease Study, Berlin, Germany. ¹⁰⁹Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Haertelstrasse 16-18, 04107 Leipzig, Germany. ¹¹⁰LIFE Research Centre for Civilization Diseases, University of Leipzig, Philipp-Rosenthal-Straße 27, 04103 Leipzig, Germany. ¹¹¹Radboud University Medical Center, Radboud Institute for Health Sciences, Nijmegen, The Netherlands. ¹¹²Quantinuum Research LLC, Wayne, PA 19087, USA. ¹¹³Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine, St. Louis, MO, USA. ¹¹⁴Department of Epidemiology, University of Pittsburgh, Pittsburgh, PA, USA. ¹¹⁵Institute for Biomedicine, Eurac Research, Affiliated Institute of the University of Lübeck, Via Galvani 31, 39100 Bolzano, Italy. ¹¹⁶Department of Clinical Biochemistry, Lillebaelt Hospital, Vejle, Denmark. ¹¹⁷Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle 98101, USA. ¹¹⁸Unidad de Investigación Médica en Bioquímica, Hospital de Especialidades, Centro Médico Nacional Siglo XXI, Instituto Mexicano del Seguro Social, Mexico City, Mexico. ¹¹⁹The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ¹²⁰Department of Twin Research and Genetic Epidemiology, King's College London, London SE1 7EH, UK. ¹²¹NIHR Biomedical Research Centre at Guy's and St Thomas' Foundation Trust, London SE1 9RT, UK. ¹²²Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. ¹²³Institute of Cardiovascular Sciences, University College London, Gower Street, London WC1E 6BT, UK. ¹²⁴Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London WC1E 6BT, UK. ¹²⁵Department of Surgery, University of Pennsylvania, Philadelphia, PA, USA. ¹²⁶Amsterdam UMC, Department of Epidemiology and Data Science, Amsterdam Public Health Research Institute, Amsterdam 1081HV, the Netherlands. ¹²⁷Department of Cell and Chemical Biology, Leiden University Medical Center, Leiden 2333ZA, The Netherlands. ¹²⁸Montreal Heart Institute, Université de Montréal, 5000 Belanger Street, Montreal, PQ H1T1C8, Canada. ¹²⁹Icelandic Heart Association, 201 Kopavogur, Iceland. ¹³⁰Department of Anthropology, University of Toronto at Mississauga, Mississauga, ON L5L 1C6, Canada. ¹³¹Vth Department of Medicine, Medical Faculty Mannheim, Heidelberg University, 68167 Mannheim, Germany. ¹³²SYNLAB MVZ Humangenetik Mannheim GmbH, 68163 Mannheim, Germany. ¹³³Shanghai Institute of Nutrition and Health, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai, China. ¹³⁴Biomedical Technology Research Center, Tokushima Research Institute, Otsuka Pharmaceutical Co., Ltd, Tokushima, Japan. ¹³⁵School of Life Sciences, Westlake University, Hangzhou 310024, Zhejiang, China. ¹³⁶Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou 310024, Zhejiang, China. ¹³⁷Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD 4072, Australia. ¹³⁸Nuffield Department of Population Health, University of Oxford, Oxford, UK. ¹³⁹Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands. ¹⁴⁰Section of Statistical Multi-Omics, Department of Clinical and Experimental Research, University of Surrey, Guildford, Surrey, UK. ¹⁴¹Laboratory of Neurogenetics, National Institute On Aging, NIH, Bethesda, MD, USA. ¹⁴²Data Tecnica International, Glen Echo, MD, USA. ¹⁴³MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh EH4 2XU, Scotland. ¹⁴⁴Institute for Medical Informatics, Biometry and Epidemiology, University of Duisburg-Essen, Essen, Germany. ¹⁴⁵Centre for Public Health, Queen's University of Belfast, Belfast, Northern Ireland. ¹⁴⁶Genomic Oncology Area, GENYO, Centre for Genomics and Oncological Research: Pfizer-University of Granada-Andalusian Regional Government, Granada, Spain. ¹⁴⁷Hematology Department, Hospital Universitario Virgen de Las Nieves, Granada, Spain. ¹⁴⁸Instituto de Investigación Biosanitaria de Granada (Ibs.GRANADA), Granada, Spain. ¹⁴⁹Department of Epidemiology and Biostatistics, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China. ¹⁵⁰Genome Institute of Singapore, Agency for Science, Technology and Research, Singapore, Singapore. ¹⁵¹Public Health Informatics Unit, Department of Integrated Health Sciences, Nagoya University Graduate School of Medicine, Nagoya 461-8673, Japan. ¹⁵²MRC Unit for Lifelong Health and Ageing at UCL, 1-19 Torrington Place, London WC1E 7HB, UK. ¹⁵³Tampere Centre for Skills Training and Simulation, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland. ¹⁵⁴Brown Foundation Institute of Molecular Medicine, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX 77030, USA. ¹⁵⁵CONACYT, Instituto Nacional de Ciencias Médicas Y Nutrición Salvador Zubirán, Ciudad de México, México. ¹⁵⁶Programs in Metabolism and Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA. ¹⁵⁷Departamento de Medicina Genómica Y Toxicología Ambiental, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México,

Coyoacán, 04510 Mexico City, Mexico. ¹⁵⁸Departamento de Genómica Computacional, Instituto Nacional de Medicina Genómica, Mexico City, Mexico. ¹⁵⁹Center for Diabetes Research, University of Bergen, Bergen, Norway. ¹⁶⁰Genomic Research On Complex Diseases (GRC Group), CSIR-Centre for Cellular and Molecular Biology, Hyderabad, Telangana, India. ¹⁶¹Academy of Scientific and Innovative Research (AcSIR), New Delhi, India. ¹⁶²Epidemiology, School of Public Health, University of Alabama at Birmingham, Birmingham, AL, USA. ¹⁶³Hunter Medical Research Institute, Newcastle, Australia. ¹⁶⁴Medical Research Institute, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul 03181, Korea. ¹⁶⁵Department of Clinical Research Design & Evaluation, SAHST, Sungkyunkwan University, Seoul 06355, Korea. ¹⁶⁶Center for Cohort Studies, Total Healthcare Center, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul 04514, Korea. ¹⁶⁷Department of Occupational and Environmental Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul 03181, Korea. ¹⁶⁸Centre for Global Health Research, Usher Institute, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, Scotland. ¹⁶⁹Thurston Arthritis Research Center, University of North Carolina, Chapel Hill, NC, USA. ¹⁷⁰Duke-NUS Medical School, Health Services and Systems Research, Singapore 169857, Singapore. ¹⁷¹Division of Biomedical Informatics and Personalized Medicine, Department of Medicine, Anschutz Medical Campus, University of Colorado, Denver, Aurora, CO 80045, USA. ¹⁷²Genomics and Molecular Medicine Unit, CSIR-Institute of Genomics and Integrative Biology, New Delhi 110020, India. ¹⁷³Academy of Scientific and Innovative Research, CSIR-Human Resource Development Centre, Ghaziabad, Uttar Pradesh, India. ¹⁷⁴Departments of Ophthalmology and Human Genetics, Radboud University Nijmegen Medical Center, Philips Van Leydenlaan 15, Nijmegen 6525 EX, the Netherlands. ¹⁷⁵Vanderbilt Epidemiology Center, Division of Epidemiology, Vanderbilt University Medical Center, Nashville, USA. ¹⁷⁶Department of Pediatrics, University of California San Francisco, Oakland, CA 94609, USA. ¹⁷⁷National Center for Global Health and Medicine, Tokyo 1628655, Japan. ¹⁷⁸Department of Genetics, University of North Carolina, Chapel Hill, NC 27599, USA. ¹⁷⁹Department of Biostatistics and Epidemiology, University of Massachusetts-Amherst, Amherst, MA 01003, USA. ¹⁸⁰Department of Cardiovascular Sciences, University of Leicester, Leicester, UK. ¹⁸¹NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester, UK. ¹⁸²Beijing Institute of Ophthalmology, Beijing Key Laboratory of Ophthalmology and Visual Sciences, Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, 17 Hougou Lane, Chong Wen Men, Beijing 100005, China. ¹⁸³Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, 1 Dong Jiao Min Xiang, Beijing 100730, Dong Cheng District, China. ¹⁸⁴Institute of Genetics and Biophysics "Adriano Buzzati-Traverso" - CNR, Naples, Italy. ¹⁸⁵IRCCS Neuromed, Pozzilli, Isernia, Italy. ¹⁸⁶Division of Biostatistics, Washington University, St. Louis, MO 63110, USA. ¹⁸⁷Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, USA. ¹⁸⁸Department of Neurological Sciences, Rush University Medical Center, Chicago, IL, USA. ¹⁸⁹Dept of Nephrology, University Hospital Regensburg, Regensburg, Germany. ¹⁹⁰Institute for Maternal and Child Health, IRCCS Burlo Garofolo, Trieste, Italy. ¹⁹¹Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany. ¹⁹²German Center for Diabetes Research (DZD), Munich-Neuherberg, Germany. ¹⁹³Department of Genetics and Bioinformatics, Dasman Diabetes Institute, Kuwait City, Kuwait. ¹⁹⁴Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University of Athens, Athens, Greece. ¹⁹⁵Department of Clinical Epidemiology, Institute of Health Informatics, University College London, London, UK. ¹⁹⁶Clinical Division of Neurogeriatrics, Department of Neurology, Medical University of Graz, Graz, Austria. ¹⁹⁷Institute for Medical Informatics, Statistics and Documentation, Medical University of Graz, Graz, Austria. ¹⁹⁸Massachusetts General Hospital Cancer Center, Charlestown, MA 02129, USA. ¹⁹⁹Institute of Genetic and Biomedical Research, National Research Council of Italy, UOS of Sassari, Sassari, Italy. ²⁰⁰Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen 9700 RB, the Netherlands. ²⁰¹Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland. ²⁰²Centre for Population Health Research, University of Turku and Turku University Hospital, Turku, Finland. ²⁰³Sleep Medicine and Circadian Disorders, Brigham and Women's Hospital, Boston, MA 02115, USA. ²⁰⁴Division of Sleep Medicine, Harvard Medical School, Boston, MA 02115, USA. ²⁰⁵Central Diagnostics Laboratory, Division Laboratories, Pharmacy, and Biomedical Genetics, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands. ²⁰⁶Laboratory of Epidemiology and Population Sciences, National Institute On Aging, NIH, Baltimore, MD 20892-9205, USA. ²⁰⁷Department of Engineering Technology, University of Houston-Sugarland, Houston, TX, USA. ²⁰⁸Interfaculty Institute for Genetics and Functional Genomics, Department of Functional Genomics, University of Greifswald and University Medicine Greifswald, Greifswald, Germany. ²⁰⁹Center for Research On Genomics and Global Health, National Human Genome Research Institute, National Institutes of Health, 12 South Drive, Room 1025, Bethesda, MD 20892, USA. ²¹⁰Oneomics, Co. Ltd. 2F, Soonchunhyang Mirai Medical Center 173, Buheuyng-Ro, Bucheon-Si Gyeonggi-Do 14585, Korea. ²¹¹Department of Clinical Biochemistry and Immunology, Hospital of Southern Jutland, Kresten Philipsens Vej 15, 6200 Aabenraa, Denmark. ²¹²Department of Clinical Biochemistry, Lillebaelt Hospital, Kolding, Denmark. ²¹³Department of Biomedical Science, Hallym University, Chuncheon 24252, Gangwon-Do, Korea. ²¹⁴Centre for Bone and Arthritis Research, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ²¹⁵Bioinformatics Core Facility, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ²¹⁶Institute of Medical Informatics and Statistics, Kiel University, Kiel, Germany. ²¹⁷Institute of Translational Genomics, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany. ²¹⁸Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford, UK. ²¹⁹School of Medicine and Public Health, College of Health, Medicine and Wellbeing, University of Newcastle, Newcastle, NSW 2308, Australia. ²²⁰Center for Geriatrics and Gerontology, Division of Endocrinology and Metabolism, Department of Internal Medicine, Taichung Veterans General Hospital, Taichung, Taiwan. ²²¹School of Medicine, National Yang-Ming University, Taipei, Taiwan. ²²²School of Medicine, National Defense Medical Center, Taipei, Taiwan. ²²³Division of Endocrinology and Metabolism, Department of Internal Medicine, Taichung Veterans General Hospital, Taichung, Taiwan. ²²⁴Department of Medicine, School of Medicine, National Yang Ming Chiao Tung University, Taipei, Taiwan. ²²⁵Department of Kinesiology, Université Laval, Québec, Canada. ²²⁶Department of Clinical Chemistry, Fimlab Laboratories, 33520 Tampere, Finland. ²²⁷Department of Clinical Chemistry, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Health Technology, Tampere University, 33014 Tampere, Finland. ²²⁸Department of Cardiology, Heart Center, Tampere University Hospital, 33521 Tampere, Finland. ²²⁹Department of Cardiology, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Health Technology, Tampere University, 33014 Tampere, Finland. ²³⁰University of Queensland Diamantina Institute, Translational Research Institute,

Kent St, Woolloongabba, Brisbane, QLD 4102, Australia. ²³¹Department of Medicine, Bornholms Hospital, Rønne, Denmark. ²³²Department of Epidemiology, Ryals School of Public Health, University of Alabama at Birmingham, Birmingham, AL, USA. ²³³Cardiology, Division Heart and Lungs, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands. ²³⁴Department of Population and Quantitative Health Sciences, Case Western Reserve University, Cleveland, OH 44106, USA. ²³⁵Department of Genetics, Stanford University School of Medicine Stanford, Palo Alto, CA 94305, USA. ²³⁶Division of Endocrinology, Ohio State University, Columbus, OH 43210, USA. ²³⁷Department of Epidemiology, University of Washington, Seattle, WA 98195, USA. ²³⁸School of Medicine and Health Sciences, George Washington University, Washington, DC 20037, USA. ²³⁹Department of Epidemiology, School of Public Health, Peking University Health Science Center, Beijing, China. ²⁴⁰Institute for Laboratory Medicine, University Hospital Leipzig, Paul-List-Strasse 13/15, 04103 Leipzig, Germany. ²⁴¹Laboratory of Epidemiology and Population Sciences, National Institute On Aging, NIH, Baltimore, MD 20892-9205, USA. ²⁴²Center for Alzheimer's and Related Dementias, NIH, Bethesda, MD, USA. ²⁴³Data Tecnica International, Washington, DC, USA. ²⁴⁴Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland. ²⁴⁵Department of Environmental and Preventive Medicine, Jichi Medical University School of Medicine, Shimotsuke 329-0498, Japan. ²⁴⁶Centre for Population Health Sciences, Usher Institute, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, Scotland. ²⁴⁷Department of Functional Pathology, Shimane University School of Medicine, Izumo 6938501, Japan. ²⁴⁸Department of Pediatrics and Adolescent Medicine, Turku University Hospital and University of Turku, Turku, Finland. ²⁴⁹Department of Physiology, University of Turku, Turku, Finland. ²⁵⁰University of Split School of Medicine, Šoltanska 2, HR-21000 Split, Croatia. ²⁵¹University of Leipzig Medical Center, Liebigstr. 18, 04103 Medical Department III – Endocrinology, Nephrology, Rheumatology Leipzig, Germany. ²⁵²Department of Nutrition-Dietetics, Harokopio University, Eleftheriou Venizelou, 17676 Athens, Greece. ²⁵³Department of Clinical Gene Therapy, Osaka University Graduate School of Medicine, Suita 5650871, Japan. ²⁵⁴Department of Geriatric and General Medicine, Osaka University Graduate School of Medicine, Suita 5650871, Japan. ²⁵⁵Department of Vascular Surgery, Division of Surgical Specialties, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands. ²⁵⁶Corneal Dystrophy Research Institute, Yonsei University College of Medicine, Saevit Eye Hospital, Seoul 03722, Korea. ²⁵⁷Dept of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands. ²⁵⁸Latin American Brain Health (BrainLat), Universidad Adolfo Ibáñez, Santiago, Chile. ²⁵⁹Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, 3584CG Utrecht, the Netherlands. ²⁶⁰Second Department of Cardiology, Medical School, National and Kapodistrian University of Athens, Attikon University Hospital, Athens, Greece. ²⁶¹Center for Vision Research, Department of Ophthalmology and The Westmead Institute, University of Sydney, Hawkesbury Rd, Sydney, NSW 2145, Australia. ²⁶²School of Medicine, Menzies Institute for Medical Research, University of Tasmania, Liverpool St, Hobart, TAS 7000, Australia. ²⁶³Centre for Eye Research Australia, University of Melbourne, Melbourne, VIC 3002, Australia. ²⁶⁴Department of Clinical Physiology, Tampere University Hospital, 33521 Tampere, Finland. ²⁶⁵Department of Clinical Physiology, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Health Technology, Tampere University, 33014 Tampere, Finland. ²⁶⁶Centre Nutrition, Santé Et Société (NUTRIS), Institute of Nutrition and Functional Foods (INAF), Québec, Canada. ²⁶⁷Pennington Biomedical Research Center, Baton Rouge, LA 70808, USA. ²⁶⁸Discipline of Internal Medicine, Medical School, The University of Western Australia, Perth, WA, Australia. ²⁶⁹Institute of Epidemiology, Kiel University, Kiel, Germany. ²⁷⁰Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany. ²⁷¹Department of Drug Treatment, Sahlgrenska University Hospital, Gothenburg, Sweden. ²⁷²Geriatric Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ²⁷³Department of Internal Medicine, EwhaWomans University School of Medicine, Seoul, Korea. ²⁷⁴Division of Cancer Control and Population Sciences, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA 15232, USA. ²⁷⁵Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA 15232, USA. ²⁷⁶Healthy Longevity Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117545, Singapore. ²⁷⁷Singapore Institute for Clinical Sciences, Agency for Science Technology and Research (A*STAR), Singapore 117609, Singapore. ²⁷⁸Department of Endocrinology and Metabolism, Kyung Hee University School of Medicine, Seoul 02447, Korea. ²⁷⁹Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany. ²⁸⁰Laboratory of Epidemiology and Population Science, National Institute On Aging Intramural Research Program, NIH Biomedical Research Center, NIH 251 Bayview Blvd, Baltimore, MD 21224, USA. ²⁸¹Algebra University College, Ilica 242, Zagreb, Croatia. ²⁸²Department of Physical Activity and Health, Paavo Nurmi Centre, Sports and Exercise Medicine Unit, University of Turku, Turku, Finland. ²⁸³Interdisciplinary Center Psychopathology and Emotion Regulation (ICPE), University of Groningen, University Medical Center Groningen, Groningen 9700 RB, the Netherlands. ²⁸⁴Institute of Molecular Genetics, National Research Council of Italy, Pavia, Italy. ²⁸⁵Gottfried Schatz Research Center for Cell Signaling, Metabolism and Aging, Medical University of Graz, Graz, Austria. ²⁸⁶Local Health Unit Toscana Centro, Florence, Italy. ²⁸⁷Institute of Nutritional Science, University of Potsdam, Nuthetal, Germany. ²⁸⁸Department of Medicine, Surgery and Health Sciences, University of Trieste, Strada Di Fiume 447, 34149 Trieste, Italy. ²⁸⁹Department of Nephrology, Traunstein Hospital, Diabetology, Rheumatology Traunstein, Germany. ²⁹⁰KfH Kidney Center Traunstein, Traunstein, Germany. ²⁹¹Department of Neurology, Center for Translational and Systems Neuroimmunology, Columbia University Medical Center, New York, NY, USA. ²⁹²Medical School, National and Kapodistrian University Athens, 75 M. Assias Street, 115 27 Athens, Greece. ²⁹³Dromokaiteio Psychiatric Hospital, 124 61 Athens, Greece. ²⁹⁴Clinical Pharmacology, William Harvey Research Institute, Queen Mary University of London, London EC1M 6BQ, UK. ²⁹⁵Department of Ophthalmology, Medical Faculty Mannheim, Heidelberg University, Kutzerufer 1, 68167 Mannheim, Germany. ²⁹⁶Institute of Molecular and Clinical Ophthalmology, Basel, Switzerland. ²⁹⁷Privatpraxis Prof Jonas Und Dr Panda-Jonas, Heidelberg, Germany. ²⁹⁸Department of Human Genetics, David Geffen School of Medicine at UCLA, University of California, Los Angeles, CA, USA. ²⁹⁹Unidad de Biología Molecular Y Medicina Genómica, Instituto de Investigaciones Biomédicas UNAM/ Instituto Nacional de Ciencias Médicas Y Nutrición Salvador Zubirán, Mexico City, Mexico. ³⁰⁰Departamento de Endocrinología Y Metabolismo, Instituto Nacional de Ciencias Médicas Y Nutrición Salvador Zubirán, 14080 Mexico, Mexico. ³⁰¹Department of Nutrition, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA. ³⁰²Carolina Population Center, University of North Carolina, Chapel Hill, NC 27516, USA. ³⁰³USC–Office of Population Studies Foundation, University of San Carlos, 6000 Cebu City, Philippines. ³⁰⁴Department of Anthropology, Sociology, and History, University of San Carlos, 6000 Cebu City, Philippines. ³⁰⁵Department of Medicine, Faculty of Medicine, University of Kelaniya, Ragama 11010, Sri Lanka. ³⁰⁶Department of Public Health, Faculty of Medicine, University

of Kelaniya, Ragama 11010, Sri Lanka. ³⁰⁷Children's Hospital Oakland Research Institute, Oakland, CA 94609, USA. ³⁰⁸Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan. ³⁰⁹Systems Genomics Laboratory, School of Biotechnology, Jawaharlal Nehru University, New Delhi 110067, India. ³¹⁰Department of Medicine, University of Mississippi Medical Center, Jackson, MS 39216, USA. ³¹¹Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS 39216, USA. ³¹²Department of Medical Sciences, Uppsala University, Uppsala, Sweden. ³¹³Department of Paediatrics, Yong Loo Lin School of Medicine, National University of Singapore and Khoo Teck Puat - National University Children's Medical Institute, National University Health System, Singapore, Singapore. ³¹⁴Department of Medicine, University of North Carolina, Chapel Hill, NC, USA. ³¹⁵Injury Prevention Research Center, University of North Carolina, Chapel Hill, NC, USA. ³¹⁶Division of Physical Therapy, University of North Carolina, Chapel Hill, NC, USA. ³¹⁷Department of Psychiatry, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands. ³¹⁸Department of Biochemistry, College of Medicine, Ewha Womans University, Seoul 07804, Korea. ³¹⁹Faculty of Health and Medicine, University of Newcastle, Callaghan, Australia. ³²⁰Division of Biostatistics, Washington University School of Medicine, St. Louis, USA. ³²¹Office of the Provost, University of South Carolina, Columbia, SC, USA. ³²²Department of Internal Medicine, University of Utah, Salt Lake City, Utah 84132, USA. ³²³Institute of Cellular Medicine (Diabetes), The Medical School, Newcastle University, Framlington Place, Newcastle Upon Tyne NE2 4HH, UK. ³²⁴Department of Medicine, Helsinki University Hospital, University of Helsinki, Haartmaninkatu 4, P.O.Box 340, 00029 Helsinki, Finland. ³²⁵Minerva Foundation Institute for Medical Research, Biomedicum 2U, Tukholmankatu 8, 00290 Helsinki, Finland. ³²⁶JSS Academy of Higher Education and Research, Mysuru, India. ³²⁷Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA. ³²⁸Harvard Medical School, Boston, MA 02115, USA. ³²⁹Instituto Nacional de Salud Publica Y Centro de Estudios en Diabetes, Cuernavaca, Mexico. ³³⁰Laboratorio de Inmunogenómica Y Enfermedades Metabólicas, Instituto Nacional de Medicina Genómica, Mexico City, Mexico. ³³¹Human Genetics Center, School of Public Health, University of Texas Health Science Center at Houston, Houston, TX 77030, USA. ³³²Yong Loo Lin School of Medicine, National University of Singapore and National University Health System, Singapore 119228, Singapore. ³³³Department of Exercise and Nutrition Sciences, Milken Institute School of Public Health, George Washington University, Washington, DC 20052, USA. ³³⁴Kurume University School of Medicine, Kurume 830-0011, Japan. ³³⁵Genetics, Merck Sharp & Dohme Corp., Kenilworth, NJ 07033, USA. ³³⁶Population Health and Genomics, University of Dundee, Ninewells Hospital and Medical School, Dundee DD1 9SY, UK. ³³⁷Intramural Research Program, National Institute On Aging, 3001 S. Hanover St., Baltimore, MD 21225, USA. ³³⁸Beijing Institute of Ophthalmology, Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing Ophthalmology and Visual Sciences Key Laboratory, Beijing 100730, China. ³³⁹The Eye Hospital, School of Ophthalmology and Optometry, Wenzhou Medical University, Wenzhou 325027, Zhejiang, China. ³⁴⁰Synlab Academy, SYNLAB Holding Deutschland GmbH, Mannheim and Augsburg, Germany. ³⁴¹Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz, Austria. ³⁴²Faculty of Medicine, University of Iceland, 101 Reykjavik, Iceland. ³⁴³Department of Biomedical Data Sciences, Section Molecular Epidemiology, Leiden University Medical Center, Leiden 2333ZA, The Netherlands. ³⁴⁴Department of Epidemiology and Data Science, Amsterdam UMC, Amsterdam 1081HV, the Netherlands. ³⁴⁵Amsterdam Public Health Research Institute, Amsterdam Cardiovascular Sciences, Amsterdam 1081HV, the Netherlands. ³⁴⁶Department of General Practice, Amsterdam UMC, Amsterdam 1081HV, the Netherlands. ³⁴⁷Amsterdam Public Health Research Institute, Health Behaviours and Chronic Diseases, Amsterdam 1081HV, the Netherlands. ³⁴⁸Corporal Michael Crescenzi VA Medical Center, Philadelphia, PA 19104, USA. ³⁴⁹Institute of Social and Economic Research, University of Essex, Wivenhoe Park, Colchester CO4 3SQ, UK. ³⁵⁰Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³⁵¹Department of Epidemiology, University of Washington, Seattle, WA, USA. ³⁵²Department of Health Systems and Population Health, University of Washington, Seattle, WA, USA. ³⁵³Institute of Regional Health Research, University of Southern Denmark, Odense, Denmark. ³⁵⁴Danish Aging Research Center, University of Southern Denmark, Odense C, Denmark. ³⁵⁵Public Health, Faculty of Medicine, University of Helsinki, Helsinki, Finland. ³⁵⁶Broad Institute of MIT and Harvard, Cambridge, MA, USA. ³⁵⁷Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA. ³⁵⁸Department of Pediatrics, The University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, USA. ³⁵⁹Division of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA. ³⁶⁰School of Medicine, Southern University of Science and Technology, Shenzhen, China. ³⁶¹Institute for Cardiogenetics, University of Lübeck, DZHK (German Research Centre for Cardiovascular Research), Partner Site Hamburg/Lübeck/Kiel, and University Heart Center Lübeck, Lübeck, Germany. ³⁶²Netherlands Heart Institute, Utrecht, the Netherlands. ³⁶³Division of Cardiology, Department of Medicine, Massachusetts General Hospital, Boston, MA, USA. ³⁶⁴Department of Medicine, Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA. ³⁶⁵Department of Medicine, Harvard Medical School, Boston, MA, USA. ³⁶⁶Verve Therapeutics, Cambridge, MA, USA. ³⁶⁷Northern Finland Birth Cohorts, Infrastructure for Population Studies, Faculty of Medicine, University of Oulu, Oulu, Finland. ³⁶⁸Center for Life Course Health Research, Faculty of Medicine, University of Oulu, Oulu, Finland. ³⁶⁹Biocenter of Oulu, University of Oulu, Oulu, Finland. ³⁷⁰Institute for Genetic and Biomedical Research, Italian National Council of Research (IRGB CNR), Cagliari, Italy. ³⁷¹University of Sassari, Sassari, Italy. ³⁷²Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands. ³⁷³Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, the Netherlands. ³⁷⁴Department of Internal Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, the Netherlands. ³⁷⁵Eindhoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, the Netherlands. ³⁷⁶Department of Human Genetics, Leiden University Medical Center, Leiden, the Netherlands. ³⁷⁷Population Health Research Institute, St George's, University of London, London SW17 0RE, UK. ³⁷⁸National Heart and Lung Institute, Imperial College London, London W12 0NN, UK. ³⁷⁹School of Public Health, Imperial College London, London W12 7RH, UK. ³⁸⁰Department of Internal Medicine, National Taiwan University Hospital, No. 7, Chung-Shan South Road, Taipei, Taiwan. ³⁸¹OCDEM, University of Oxford, Churchill Hospital, Oxford OX3 7LE, UK. ³⁸²NIHR Oxford Biomedical Research Centre, Churchill Hospital, Oxford, UK. ³⁸³Ocular Epidemiology, Singapore Eye Research Institute, Singapore National Eye Centre, Singapore 168751, Singapore. ³⁸⁴Ophthalmology and Visual Sciences Academic Clinical Program (Eye ACP), Duke-NUS Medical School, Singapore 169857, Singapore. ³⁸⁵Data Science, Singapore Eye Research Institute, Singapore National Eye Centre, Singapore 168751, Singapore. ³⁸⁶Medical School, University of Exeter, University of Exeter, Exeter EX2 5DW, UK. ³⁸⁷Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. ³⁸⁸Framingham Heart Study,

National Heart, Lung, and Blood Institute, US National Institutes of Health, Bethesda, MD, USA. ³⁸⁹Department of Genetics, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. ³⁹⁰Department of Genetics, Shanghai-MOST Key Laboratory of Health and Disease Genomics, Chinese National Human Genome Center at Shanghai, Shanghai 201203, China. ³⁹¹Computational Medicine, Berlin Institute of Health at Charité – Universitätsmedizin, Berlin, Germany. ³⁹²Technical University of Munich (TUM) and Klinikum Rechts Der Isar, TUM School of Medicine, Munich, Germany. ³⁹³Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland. ³⁹⁴Stanford Cardiovascular Institute, Stanford University, Stanford, CA 94305, USA. ³⁹⁵Stanford Diabetes Research Center, Stanford University, Stanford, CA 94305, USA. ³⁹⁶Regeneron Pharmaceuticals, Tarrytown, NY, USA. ³⁹⁷Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore 308232, Singapore. ³⁹⁸Imperial College Healthcare NHS Trust, Imperial College London, London W12 0HS, UK. ³⁹⁹Imperial College Healthcare NHS Trust, London W12 0HS, UK. ⁴⁰⁰MRC-PHE Centre for Environment and Health, Imperial College London, London W2 1PG, UK. ⁴⁰¹School of Electrical and Information Engineering, University of the Witwatersrand, Johannesburg, South Africa. ⁴⁰²Institute for Minority Health Research, University of Illinois College of Medicine, Chicago, IL, USA. ⁴⁰³Department of Biostatistics, Harvard T.H. Chan School of Public Health, 677 Huntington Avenue, Boston, MA 02115, USA. ⁴⁰⁴QIMR Berghofer Medical Research Institute, 300 Herston Road, Brisbane, QLD 4006, Australia. ⁴⁰⁵Faisalabad Institute of Cardiology, Faisalabad, Pakistan. ⁴⁰⁶Department of Medicine, Columbia University Irving Medical Center, New York, NY, USA. ⁴⁰⁷Department of Cardiology, Columbia University Irving Medical Center, New York, NY, USA. ⁴⁰⁸Big Data Institute, University of Oxford, Oxford OX3 7LF, UK. ⁴⁰⁹National Institute for Health Research Oxford Biomedical Research Centre, Oxford University Hospitals, Oxford, UK. ⁴¹⁰Aberdeen Centre for Health Data Science, 1:042 Polwarth Building School of Medicine, Medical Science and Nutrition University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK. ⁴¹¹Division of Population Health and Genomics, Ninewells Hospital and Medical School, University of Dundee, Dundee DD1 9SY, UK. ⁴¹²Department of Population Health Sciences, Geisinger Health, Danville, PA 17822, USA. ⁴¹³School of Basic and Medical Biosciences, Faculty of Life Sciences and Medicine, King's College London, London, UK. ⁴¹⁴Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN, USA. ⁴¹⁵Departments of Medicine (Medical Genetics) and Genome Sciences, University of Washington, Washington, USA. ⁴¹⁶Center for Autoimmune Genomics and Etiology, Cincinnati Children's Hospital Medical Center (CCHMC), Cincinnati, OH, USA. ⁴¹⁷Division of Endocrinology, Metabolism, and Molecular Medicine, Department of Medicine, Northwestern University, Feinberg School of Medicine, Chicago, IL 60618, USA. ⁴¹⁸Department of Anthropology, Northwestern University, Evanston, IL 60208, USA. ⁴¹⁹Department of Public Health and Nursing, HUNT Research Centre, NTNU, Norwegian University of Science and Technology, 7600 Levanger, Norway. ⁴²⁰Department of Research, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. ⁴²¹Department of Endocrinology, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. ⁴²²RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan. ⁴²³Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan. ⁴²⁴Laboratory of Complex Trait Genomics, Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan. ⁴²⁵Laboratory for Systems Genetics, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan. ⁴²⁶Department of Genome Informatics, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan. ⁴²⁷Division of Molecular Pathology, Institute of Medical Science, The University of Tokyo, Tokyo, Japan. ⁴²⁸Faculty of Medicine, University of Iceland, Sæmundargötu 2, Reykjavík 102, Iceland. ⁴²⁹VA Boston Healthcare System, Boston, MA, USA. ⁴³⁰VA Informatics and Computing Infrastructure, VA Salt Lake City Health Care System, Salt Lake City, UT, USA. ⁴³¹University of Massachusetts, Boston, MA, USA. ⁴³²Cardiovascular Institute, Stanford University School of Medicine, Stanford, CA, USA. ⁴³³Corporal Michael J. Crescenz VA Medical Center, Philadelphia, PA, USA. ⁴³⁴Department of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ⁴³⁵Department of Medicine, Brigham Women's Hospital, Boston, MA, USA. ⁴³⁶Division of Cardiology, Emory University School of Medicine, Atlanta, GA, USA. ⁴³⁷Departments of Pediatrics and Genetics, Harvard Medical School, Boston, MA, USA. ⁴³⁸Centre for Genetics and Genomics Versus Arthritis, Centre for Musculoskeletal Research, Division of Musculoskeletal and Dermatological Sciences, The University of Manchester, Manchester, UK. ⁴³⁹Cardiology Division, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA. ⁴⁴⁰Department of Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA. ⁴⁴¹Cardiovascular Research Center and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA. ⁴⁴²Princess Al-Jawhara Al-Brahim Centre for Excellence in Research of Hereditary Disorders (PACER-HD), King Abdulaziz University, Jeddah, Saudi Arabia. ⁴⁴³Department of Human Genetics, University of Michigan, Ann Arbor, MI 48109, USA.

Received: 5 April 2022 Accepted: 6 December 2022

Published online: 27 December 2022

References

- Castelli WP, Anderson K, Wilson PW, Levy D. Lipids and risk of coronary heart disease. The Framingham Study. *Ann Epidemiol.* 1992;2:23–8.
- GBD. Diseases and Injuries Collaborators: Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet.* 2019;2020(396):1204–22.
- Grundy SM, Stone NJ, Bailey AL, Beam C, Birtcher KK, Blumenthal RS, Braun LT, de Ferranti S, Faiella-Tommasino J, Forman DE, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation.* 2019;139:e1082–143.
- Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research, Saxena R, Voight BF, Lyssenko V, Burt NP, de Bakker PI, Chen H, Roix JJ, et al. Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science.* 2007;316:1331–6.

5. Kathiresan S, Manning AK, Demissie S, D'Agostino RB, Surti A, Guiducci C, Gianniny L, Burt NP, Melander O, Orho-Melander M, et al. A genome-wide association study for blood lipid phenotypes in the Framingham Heart Study. *BMC Med Genet*. 2007;8(Suppl 1):S17.
6. Kathiresan S, Melander O, Anevski D, Guiducci C, Burt NP, Roos C, Hirschhorn JN, Berglund G, Hedblad B, Groop L, et al. Polymorphisms associated with cholesterol and risk of cardiovascular events. *N Engl J Med*. 2008;358:1240–9.
7. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, Pirruccello JP, Ripatti S, Chasman DI, Willer CJ, et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*. 2010;466:707–13.
8. Asselbergs FW, Guo Y, van Iperen EP, Sivapalaratnam S, Tragante V, Lanktree MB, Lange LA, Almoguera B, Appelman YE, Barnard J, et al. Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. *Am J Hum Genet*. 2012;91:823–38.
9. Albrechtsen A, Grarup N, Li Y, Sparso T, Tian G, Cao H, Jiang T, Kim SY, Korneliusen T, Li Q, et al. Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. *Diabetologia*. 2013;56:298–310.
10. Tachmazidou I, Dedoussis G, Southam L, Farmaki AE, Ritchie GR, Xifara DK, Matchan A, Hatzikotoulas K, Rayner NW, Chen Y, et al. A rare functional cardioprotective APOC3 variant has risen in frequency in distinct population isolates. *Nat Commun*. 2013;4:2872.
11. Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, Ganna A, Chen J, Buchkovich ML, Mora S, et al. Discovery and refinement of loci associated with lipid levels. *Nat Genet*. 2013;45:1274–83.
12. Holmen OL, Zhang H, Fan Y, Hovelson DH, Schmidt EM, Zhou W, Guo Y, Zhang J, Langhammer A, Lochen ML, et al. Systematic evaluation of coding variation identifies a candidate causal variant in TM6SF2 influencing total cholesterol and myocardial infarction risk. *Nat Genet*. 2014;46:345–51.
13. Peloso GM, Auer PL, Bis JC, Voorman A, Morrison AC, Stitzel NO, Brody JA, Khetarpal SA, Crosby JR, Fornage M, et al. Association of low-frequency and rare coding-sequence variants with blood lipids and coronary heart disease in 56,000 whites and blacks. *Am J Hum Genet*. 2014;94:223–32.
14. Surakka I, Horikoshi M, Magi R, Sarin AP, Mahajan A, Lagou V, Marullo L, Ferreira T, Miraglio B, Timonen S, et al. The impact of low-frequency and rare variants on lipid levels. *Nat Genet*. 2015;47:589–97.
15. Tang CS, Zhang H, Cheung CY, Xu M, Ho JC, Zhou W, Cherny SS, Zhang Y, Holmen O, Au KW, et al. Exome-wide association analysis reveals novel coding sequence variants associated with lipid traits in Chinese. *Nat Commun*. 2015;6:10206.
16. van Leeuwen EM, Karssen LC, Deelen J, Isaacs A, Medina-Gomez C, Mbarek H, Kanterakis A, Trompet S, Postmus I, Verweij N, et al. Genome of The Netherlands population-specific imputations identify an ABCA6 variant associated with cholesterol levels. *Nat Commun*. 2015;6:6065.
17. Iotchkova V, Huang J, Morris JA, Jain D, Barbieri C, Walter K, Min JL, Chen L, Astle W, Cocca M, et al. Discovery and refinement of genetic loci associated with cardiometabolic risk using dense imputation maps. *Nat Genet*. 2016;48:1303–12.
18. Liu DJ, Peloso GM, Yu H, Butterworth AS, Wang X, Mahajan A, Saleheen D, Emdin C, Alam D, Alves AC, et al. Exome-wide association study of plasma lipids in >300,000 individuals. *Nat Genet*. 2017;49:1758–66.
19. Lu X, Peloso GM, Liu DJ, Wu Y, Zhang H, Zhou W, Li J, Tang CS, Dorajoo R, Li H, et al. Exome chip meta-analysis identifies novel loci and East Asian-specific coding variants that contribute to lipid levels and coronary artery disease. *Nat Genet*. 2017;49:1722–30.
20. Hoffmann TJ, Theusch E, Haldar T, Ranatunga DK, Jorgenson E, Medina MW, Kvale MN, Kwok PY, Schaefer C, Krauss RM, et al. A large electronic-health-record-based genome-wide study of serum lipids. *Nat Genet*. 2018;50:401–13.
21. Kanai M, Akiyama M, Takahashi A, Matoba N, Momozawa Y, Ikeda M, Iwata N, Ikegawa S, Hirata M, Matsuda K, et al. Genetic analysis of quantitative traits in the Japanese population links cell types to complex human diseases. *Nat Genet*. 2018;50:390–400.
22. Klarin D, Damrauer SM, Cho K, Sun YV, Teslovich TM, Honerlaw J, Gagnon DR, DuVall SL, Li J, Peloso GM, et al. Genetics of blood lipids among ~300,000 multi-ethnic participants of the Million Veteran Program. *Nat Genet*. 2018;50:1514–23.
23. Spracklen CN, Chen P, Kim YJ, Wang X, Cai H, Li S, Long J, Wu Y, Wang YX, Takeuchi F. Association analyses of East Asian individuals and trans-ancestry analyses with European individuals reveal new loci associated with cholesterol and triglyceride levels. *Hum Mol Genet*. 2018;27:1122.
24. Graham SE, Clarke SL, Wu KH, Kanoni S, Zajac GJM, Ramdas S, Surakka I, Ntalla I, Vedantam S, Winkler TW, et al. The power of genetic diversity in genome-wide association studies of lipids. *Nature*. 2021;600:675–9.
25. Musunuru K, Strong A, Frank-Kamenetsky M, Lee NE, Ahfeldt T, Sachs KV, Li X, Li H, Kuperwasser N, Ruda VM, et al. From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature*. 2010;466:714–9.
26. Musunuru K, Pirruccello JP, Do R, Peloso GM, Guiducci C, Sougnez C, Garimella KV, Fisher S, Abreu J, Barry AJ, et al. Exome sequencing, ANGPTL3 mutations, and familial combined hypolipidemia. *N Engl J Med*. 2010;363:2220–7.
27. Pers TH, Karjalainen JM, Chan Y, Westra HJ, Wood AR, Yang J, Lui JC, Vedantam S, Gustafsson S, Esko T, et al. Biological interpretation of genome-wide association studies using predicted gene functions. *Nat Commun*. 2015;6:5890.
28. Weeks EM, Ulirsch JC, Cheng NY, Tripple BL, Fine RS, Miao J, Patwardhan TA, Kanai M, Nasser J, Fulco CP, et al. Leveraging polygenic enrichments of gene features to predict genes underlying complex traits and diseases. *medRxiv* 2020:2020.2009.2008.20190561.
29. Stanzick KJ, Li Y, Schlosser P, Gorski M, Wuttke M, Thomas LF, Rasheed H, Rowan BX, Graham SE, Vanderweff BR, et al. Discovery and prioritization of variants and genes for kidney function in >1.2 million individuals. *Nature Communications*. 2021;12:4350.
30. The Emerging Risk Factors Collaboration, Di Angelantonio E, Sarwar N, Perry P, Kaptoge S, Ray KK, Thompson A, Wood AM, Lewington S, Sattar N, et al. Major lipids, apolipoproteins, and risk of vascular disease. *JAMA*. 2009;302:1993–2000.
31. Richardson TG, Sanderson E, Palmer TM, Ala-Korpela M, Ference BA, Davey Smith G, Holmes MV. Evaluating the relationship between circulating lipoprotein lipids and apolipoproteins with risk of coronary heart disease: a multivariable Mendelian randomisation analysis. *PLoS Med*. 2020;17:e1003062.
32. Allara E, Morani G, Carter P, Gkatzionis A, Zuber V, Foley CN, Rees JMB, Mason AM, Bell S, Gill D, et al. Genetic determinants of lipids and cardiovascular disease outcomes. *Circulation: Genomic Precision Med*. 2019;12:e002711.

33. Veturi Y, Lucas A, Bradford Y, Hui D, Dudek S, Theusch E, Verma A, Miller JE, Kullo I, Hakonarson H, et al. A unified framework identifies new links between plasma lipids and diseases from electronic medical records across large-scale cohorts. *Nat Genet.* 2021;53:972–81.
34. Bush WS, Oetjens MT, Crawford DC. Unravelling the human genome-phenome relationship using phenome-wide association studies. *Nat Rev Genet.* 2016;17:129–45.
35. Abbott RD, Garrison RJ, Wilson PW, Epstein FH, Castelli WP, Feinleib M, LaRue C. Joint distribution of lipoprotein cholesterol classes. The Framingham study. *Arteriosclerosis.* 1983;3:260–72.
36. Flynn E, Tanigawa Y, Rodriguez F, Altman RB, Sinnott-Armstrong N, Rivas MA. Sex-specific genetic effects across biomarkers. *Eur J Hum Genet.* 2021;29:154–63.
37. Zore T, Palafox M, Reue K. Sex differences in obesity, lipid metabolism, and inflammation—A role for the sex chromosomes? *Mol Metab.* 2018;15:35–44.
38. AlSiraj Y, Chen X, Thatcher SE, Temel RE, Cai L, Blalock E, Katz W, Ali HM, Petriello M, Deng P, et al. XX sex chromosome complement promotes atherosclerosis in mice. *Nat Commun.* 2019;10:2631.
39. Natarajan P, Pampana A, Graham SE, Ruotsalainen SE, Perry JA, de Vries PS, Broome JG, Pirruccello JP, Honigberg MC, Aragam K, et al. Chromosome Xq23 is associated with lower atherogenic lipid concentrations and favorable cardiometabolic indices. *Nat Commun.* 2021;12:2182.
40. Jiang L, Xue C, Dai S, Chen S, Chen P, Sham PC, Wang H, Li M. DESE: estimating driver tissues by selective expression of genes associated with complex diseases or traits. *Genome Biol.* 2019;20:233.
41. The GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science.* 2020;369:1318–30.
42. Allara E, Morani G, Carter P, Gkatzionis A, Zuber V, Foley CN, Rees JMB, Mason AM, Bell S, Gill D, et al. Genetic determinants of lipids and cardiovascular disease outcomes: a wide-angled Mendelian randomization investigation. *Circ Genom Precis Med.* 2019;12:e002711.
43. Saiz-Vazquez O, Puente-Martinez A, Ubillos-Landa S, Pacheco-Bonrostro J, Santabarbara J. Cholesterol and Alzheimer's disease risk: a meta-meta-analysis. *Brain Sci.* 2020;10:386.
44. Zhang X, Tian Q, Liu D, Geng T, Xu X, Ge S, Zheng D, Wu L, Song M, Hou H, et al. Causal association of circulating cholesterol levels with dementia: a mendelian randomization meta-analysis. *Transl Psychiatry.* 2020;10:145.
45. Tan JS, Hu MJ, Yang YM, Yang YJ. Genetic predisposition to low-density lipoprotein cholesterol may increase risks of both individual and familial Alzheimer's disease. *Front Med (Lausanne).* 2021;8:798334.
46. Deb S, Puthanveetil P, Sakthar P. A population-based cross-sectional study of the association between liver enzymes and lipid levels. *Int J Hepatol.* 2018;2018:1286170.
47. Joshi AD, Andersson C, Buch S, Stender S, Noordam R, Weng LC, Weeke PE, Auer PL, Boehm B, Chen C, et al. Four susceptibility loci for gallstone disease identified in a meta-analysis of genome-wide association studies. *Gastroenterology.* 2016;151(351–363):e328.
48. Bernabeu E, Canela-Xandri O, Rawlik K, Talenti A, Prendergast J, Tenesa A. Sex differences in genetic architecture in the UK Biobank. *Nat Genet.* 2021;53:1283–9.
49. Ruth KS, Day FR, Tyrrell J, Thompson DJ, Wood AR, Mahajan A, Beaumont RN, Wittemans L, Martin S, Busch AS, et al. Using human genetics to understand the disease impacts of testosterone in men and women. *Nat Med.* 2020;26:252–8.
50. Pirastu N, Cordioli M, Nandakumar P, Mignogna G, Abdellaoui A, Hollis B, Kanai M, Rajagopal VM, Parolo PDB, Baya N, et al. Genetic analyses identify widespread sex-differential participation bias. *Nat Genet.* 2021;53:663–71.
51. Yengo L, Sidorenko J, Kempner KE, Zheng Z, Wood AR, Weedon MN, Frayling TM, Hirschhorn J, Yang J, Visscher PM, Consortium G. Meta-analysis of genome-wide association studies for height and body mass index in approximately 700,000 individuals of European ancestry. *Hum Mol Genet.* 2018;27:3641–9.
52. Bhatt DK, Basit A, Zhang H, Gaedigk A, Lee SB, Claw KG, Mehrotra A, Chaudhry AS, Pearce RE, Gaedigk R, et al. Hepatic abundance and activity of androgen- and drug-metabolizing enzyme UGT2B17 are associated with genotype, age, and sex. *Drug Metab Dispos.* 2018;46:888–96.
53. Nielsen JB, Rom O, Surakka I, Graham SE, Zhou W, Roychowdhury T, Fritsche LG, Gagliano Taliun SA, Sidore C, Liu Y, et al. Loss-of-function genomic variants highlight potential therapeutic targets for cardiovascular disease. *Nat Commun.* 2020;11:6417.
54. Aragam KG, Jiang T, Goel A, Kanoni S, Wolford BN, Weeks EM, Wang M, Hindy G, Zhou W, Grace C, et al. Discovery and systematic characterization of risk variants and genes for coronary artery disease in over a million participants. *medRxiv* 2021:2021.2005.2024.21257377.
55. Votava JA, Parks BW. Cross-species data integration to prioritize causal genes in lipid metabolism. *Curr Opin Lipidol.* 2021;32:141–6.
56. Kherallah RY, Khawaja M, Olson M, Angiolillo D, Birnbaum Y. Cilostazol: a review of basic mechanisms and clinical uses. *Cardiovasc Drugs Ther.* 2022;36:777–92.
57. Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, Kuder JF, Wang H, Liu T, Wasserman SM, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med.* 2017;376:1713–22.
58. Schwartz GG, Steg PG, Szarek M, Bhatt DL, Bittner VA, Diaz R, Edelberg JM, Goodman SG, Hanotin C, Harrington RA, et al. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med.* 2018;379:2097–107.
59. Ray KK, Wright RS, Kallend D, Koenig W, Leiter LA, Raal FJ, Bisch JA, Richardson T, Jaros M, Wijnngaard PLJ, Kastelein JJP. Two phase 3 trials of inclisiran in patients with elevated LDL cholesterol. *N Engl J Med.* 2020;382:1507–19.
60. Nelson CP, Lai FY, Nath M, Ye S, Webb TR, Schunkert H, Samani NJ. Genetic assessment of potential long-term on-target side effects of PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) inhibitors. *Circ Genom Precis Med.* 2019;12:e002196.
61. Mahajan A, Taliun D, Thurner M, Robertson NR, Torres JM, Rayner NW, Payne AJ, Steinthorsdottir V, Scott RA, Grarup N, et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat Genet.* 2018;50:1505–13.
62. Assimes T, Catherine T, Xiang Z, Austin H, Shoa C, Valerio N, Shining M, Huaying F, Bryan RG, Kyung Min L, et al. A large-scale multi-ethnic genome-wide association study of coronary artery disease. *Nat Med.* 2022;28:1679–92.

63. Ridker PM, Revkin J, Amarenco P, Brunell R, Curto M, Civeira F, Flather M, Glynn RJ, Gregoire J, Jukema JW, et al. Cardiovascular efficacy and safety of bococizumab in high-risk patients. *N Engl J Med*. 2017;376:1527–39.
64. Hopewell JC, Malik R, Valdes-Marquez E, Worrall BB, Collins R. ISGC MCot: Differential effects of PCSK9 variants on risk of coronary disease and ischaemic stroke. *Eur Heart J*. 2018;39:354–9.
65. Hayat S, Hassan Z, Changazi SH, Zahra A, Noman M, Zain UI Abidin M, Javed H, Ans AH: Comparative analysis of serum lipid profiles in patients with and without gallstones: a prospective cross-sectional study. *Ann Med Surg (Lond)*. 2019;42:11–3.
66. Wang J, Shen S, Wang B, Ni X, Liu H, Ni X, Yu R, Suo T, Liu H. Serum lipid levels are the risk factors of gallbladder stones: a population-based study in China. *Lipids Health Dis*. 2020;19:50.
67. Gustafsson U, Sahlin S, Einarsson C. Biliary lipid composition in patients with cholesterol and pigment gallstones and gallstone-free subjects: deoxycholic acid does not contribute to formation of cholesterol gallstones. *Eur J Clin Invest*. 2000;30:1099–106.
68. Weerakoon HT, Ranasinghe S, Navaratne A, Sivakanesan R, Galketiya KB, Rosairo S. Serum lipid concentrations in patients with cholesterol and pigment gallstones. *BMC Res Notes*. 2014;7:548.
69. Chen L, Yang H, Li H, He C, Yang L, Lv G. Insights into modifiable risk factors of cholelithiasis: a Mendelian randomization study. *Hepatology*. 2022;75:785–96.
70. Yu XH, Qian K, Jiang N, Zheng XL, Cayabyab FS, Tang CK. ABCG5/ABCG8 in cholesterol excretion and atherosclerosis. *Clin Chim Acta*. 2014;428:82–8.
71. Silbernagel G, Chapman MJ, Genser B, Kleber ME, Fauler G, Scharnagl H, Grammer TB, Boehm BO, Makela KM, Kahonen M, et al. High intestinal cholesterol absorption is associated with cardiovascular disease and risk alleles in ABCG8 and ABO: evidence from the LURIC and YFS cohorts and from a meta-analysis. *J Am Coll Cardiol*. 2013;62:291–9.
72. Teupser D, Baber R, Ceglarek U, Scholz M, Illig T, Gieger C, Holdt LM, Leichtle A, Greiser KH, Huster D, et al. Genetic regulation of serum phytosterol levels and risk of coronary artery disease. *Circulation Cardiovasc Genet*. 2010;3:331–9.
73. Artha I, Bhargava A, Dharmawan NK, Pande UW, Triyana KA, Mahariski PA, Yuwono J, Bhargava V, Prabawa IPY, Manubaba I, Rina IK. High level of individual lipid profile and lipid ratio as a predictive marker of poor glycemic control in type-2 diabetes mellitus. *Vasc Health Risk Manag*. 2019;15:149–57.
74. Hussain A, Ali I, Ijaz M, Rahim A. Correlation between hemoglobin A1c and serum lipid profile in Afghani patients with type 2 diabetes: hemoglobin A1c prognosticates dyslipidemia. *Ther Adv Endocrinol Metab*. 2017;8:51–7.
75. Chen J, Spracklen CN, Marenne G, Varshney A, Corbin LJ, Luan J, Willems SM, Wu Y, Zhang X, Horikoshi M, et al. The trans-ancestral genomic architecture of glycemic traits. *Nat Genet*. 2021;53:840–60.
76. Chami N, Chen MH, Slater AJ, Eicher JD, Evangelou E, Tajuddin SM, Love-Gregory L, Kacprowski T, Schick UM, Nomura A, et al. Exome genotyping identifies pleiotropic variants associated with red blood cell traits. *Am J Hum Genet*. 2016;99:8–21.
77. Leong A, Chen J, Wheeler E, Hivert MF, Liu CT, Merino J, Dupuis J, Tai ES, Rotter JI, Florez JC, et al. Mendelian randomization analysis of hemoglobin A1c as a risk factor for coronary artery disease. *Diabetes Care*. 2019;42:1202–8.
78. McDonagh EM, Thorn CF, Bautista JM, Youngster I, Altman RB, Klein TE. PharmGKB summary: very important pharmacogenetic information for G6PD. *Pharmacogenet Genomics*. 2012;22:219–28.
79. Dore MP, Parodi G, Portoghese M, Pes GM. The controversial role of glucose-6-phosphate dehydrogenase deficiency on cardiovascular disease: a narrative review. *Oxid Med Cell Longev*. 2021;2021:5529256.
80. Spielman RS, Bastone LA, Burdick JT, Morley M, Ewens WJ, Cheung VG. Common genetic variants account for differences in gene expression among ethnic groups. *Nat Genet*. 2007;39:226–31.
81. Zhu AZ, Cox LS, Ahluwalia JS, Renner CC, Hatsukami DK, Benowitz NL, Tyndale RF. Genetic and phenotypic variation in UGT2B17, a testosterone-metabolizing enzyme, is associated with BMI in males. *Pharmacogenet Genomics*. 2015;25:263–9.
82. Yang TL, Chen XD, Guo Y, Lei SF, Wang JT, Zhou Q, Pan F, Chen Y, Zhang ZX, Dong SS, et al. Genome-wide copy-number-variation study identified a susceptibility gene, UGT2B17, for osteoporosis. *Am J Hum Genet*. 2008;83:663–74.
83. Gencer B, Bonomi M, Adorni MP, Sirtori CR, Mach F, Ruscica M. Cardiovascular risk and testosterone - from subclinical atherosclerosis to lipoprotein function to heart failure. *Rev Endocr Metab Disord*. 2021;22:257–74.
84. Firtser S, Juonala M, Magnussen CG, Julia A, Loo BM, Marniemi J, Viikari JS, Toppari J, Perheentupa A, Hutri-Kahonen N, Raitakari OT. Relation of total and free testosterone and sex hormone-binding globulin with cardiovascular risk factors in men aged 24–45 years. The Cardiovascular Risk in Young Finns Study. *Atherosclerosis*. 2012;222:257–62.
85. Schooling CM, Luo S, Au Yeung SL, Thompson DJ, Karthikeyan S, Bolton TR, Mason AM, Ingelsson E, Burgess S. Genetic predictors of testosterone and their associations with cardiovascular disease and risk factors: a Mendelian randomization investigation. *Int J Cardiol*. 2018;267:171–6.
86. Au Yeung SL, Cheng KK, Zhao J, Zhang W, Jiang C, Lam TH, Leung GM, Schooling CM. Genetically predicted 17beta-estradiol and cardiovascular risk factors in women: a Mendelian randomization analysis using young women in Hong Kong and older women in the Guangzhou Biobank Cohort Study. *Ann Epidemiol*. 2016;26:171–5.
87. Winkler TW, Day FR, Croteau-Chonka DC, Wood AR, Locke AE, Magi R, Ferreira T, Fall T, Graff M, Justice AE, et al. Quality control and conduct of genome-wide association meta-analyses. *Nat Protoc*. 2014;9:1192–212.
88. Magi R, Horikoshi M, Sofer T, Mahajan A, Kitajima H, Franceschini N, McCarthy MI, Cogent-Kidney Consortium TDCG, Morris AP. Trans-ethnic meta-regression of genome-wide association studies accounting for ancestry increases power for discovery and improves fine-mapping resolution. *Hum Mol Genet*. 2017;26:3639–50.
89. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26:2190–1.
90. Liu DJ, Peloso GM, Zhan X, Holmen OL, Zawistowski M, Feng S, Nikpay M, Auer PL, Goel A, Zhang H, et al. Meta-analysis of gene-level tests for rare variant association. *Nat Genet*. 2014;46:200–4.
91. Winkler TW, Justice AE, Cupples LA, Kronenberg F, Kutalik Z, Heid IM. consortium G: Approaches to detect genetic effects that differ between two strata in genome-wide meta-analyses: recommendations based on a systematic evaluation. *PLoS ONE*. 2017;12:e0181038.
92. Fauman EB, Hyde C: An optimal variant to gene distance window derived from an empirical definition of cis and trans protein QTLs. *bioRxiv* 2022:2022.2003.2007.483314.

93. Giambartolomei C, Vukcevic D, Schadt EE, Franke L, Hingorani AD, Wallace C, Plagnol V. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* 2014;10:e1004383.
94. Caliskan M, Manduchi E, Rao HS, Segert JA, Beltrame MH, Trizzino M, Park Y, Baker SW, Chesí A, Johnson ME, et al. Genetic and epigenetic fine mapping of complex trait associated loci in the human liver. *Am J Hum Genet.* 2019;105:89–107.
95. Barbeira AN, Dickinson SP, Bonazzola R, Zheng J, Wheeler HE, Torres JM, Torstenson ES, Shah KP, Garcia T, Edwards TL, et al. Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. *Nat Commun.* 2018;9:1825.
96. Hindy G, Dornbos P, Chaffin MD, Liu DJ, Wang M, Selvaraj MS, Zhang D, Park J, Aguilar-Salinas CA, Antonacci-Fulton L, et al. Rare coding variants in 35 genes associate with circulating lipid levels—a multi-ancestry analysis of 170,000 exomes. *Am J Hum Genet.* 2022;109:81–96.
97. Brown EE, Sturm AC, Cuchel M, Braun LT, Duell PB, Underberg JA, Jacobson TA, Hegele RA. Genetic testing in dyslipidemia: a scientific statement from the National Lipid Association. *J Clin Lipidol.* 2020;14:398–413.
98. Hegele RA, Boren J, Ginsberg HN, Arca M, Aversa M, Binder CJ, Calabresi L, Chapman MJ, Cuchel M, von Eckardstein A, et al. Rare dyslipidaemias, from phenotype to genotype to management: a European Atherosclerosis Society task force consensus statement. *Lancet Diabetes Endocrinol.* 2020;8:50–67.
99. Zhou Y, Zhang Y, Lian X, Li F, Wang C, Zhu F, Qiu Y, Chen Y. Therapeutic target database update 2022: facilitating drug discovery with enriched comparative data of targeted agents. *Nucleic Acids Res.* 2022;50:D1398–407.
100. Ge T, Chen CY, Ni Y, Feng YA, Smoller JW. Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nat Commun.* 2019;10:1776.
101. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007;81:559–75.
102. Denny JC, Ritchie MD, Basford MA, Pulley JM, Bastarache L, Brown-Gentry K, Wang D, Masys DR, Roden DM, Crawford DC. PheWAS: demonstrating the feasibility of a phenome-wide scan to discover gene-disease associations. *Bioinformatics.* 2010;26:1205–10.
103. Verma A, Bradford Y, Dudek S, Lucas AM, Verma SS, Pendergrass SA, Ritchie MD. A simulation study investigating power estimates in phenome-wide association studies. *BMC Bioinformatics.* 2018;19:120.
104. Nelson CP, Goel A, Butterworth AS, Kanoni S, Webb TR, Marouli E, Zeng L, Ntalla I, Lai FY, Hopewell JC, et al. Association analyses based on false discovery rate implicate new loci for coronary artery disease. *Nat Genet.* 2017;49:1385–91.
105. Liu Z, Zhang Y, Graham SE, Wang X, Cai D, Huang M, Pique-Regi R, Dong XC, Chen YE, Willer C, Liu W. Causal relationships between NAFLD, T2D and obesity have implications for disease subphenotyping. *J Hepatol.* 2020;73:263–76.
106. Zhou W, Nielsen JB, Fritsche LG, Dey R, Gabrielsen ME, Wolford BN, LeFaive J, VandeHaar P, Gagliano SA, Gifford A, et al. Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat Genet.* 2018;50:1335–41.
107. Graham SE, Clarke SL, Wu KH, Kanoni S, Zajac GJM, Ramdas S, Surakka I, Ntalla I, Vedantam S, Winkler TW, et al: GLGC GWAS meta-analysis results and risk score weights repository: <http://csg.sph.umich.edu/willer/public/glgc-lipids2021> 2021.
108. Kanoni S, Wang Y, Surakka I, Ramdas S, Zhu X, Costanzo M, Jang D, Burt NP, Willer CJ, Assimes TL, Peloso GM: A web browser displaying the gene prioritization and PheWAS results: https://hugeamp.org/8000/research.html?pageid=GLGC_149 2021.
109. Graham SE, Clarke SL, Wu KH, Kanoni S, Zajac GJM, Ramdas S, Surakka I, Ntalla I, Vedantam S, Winkler TW, et al: Optimized trans-ancestry polygenic score weights for LDL in the PGS Catalog: <https://www.pgscatalog.org/publication/PGP000230/> 2021.
110. Kanoni S, Graham SE, Wang Y, Surakka I, Ramdas S, Zhu X, Clarke SL, Bhatti KF, Vedantam S, Winkler TW, et al: Optimized trans-ancestry polygenic score weights for HDL, TC, TG and non-HDL in the PGS Catalog: <https://www.pgscatalog.org/publication/PGP000366/> 2022.
111. Kanoni S, Graham SE, Wang Y, Surakka I, Ramdas S, Zhu X, Clarke SL, Bhatti KF, Vedantam S, Winkler TW, et al: Implicating genes, pleiotropy and sexual dimorphism at blood lipid loci through multi-ancestry meta-analysis. Github. <https://github.com/Global-Lipids-Genetics>. 2022.
112. Kanoni S, Graham SE, Wang Y, Surakka I, Ramdas S, Zhu X, Clarke SL, Bhatti KF, Vedantam S, Winkler TW, et al. Implicating genes, pleiotropy and sexual dimorphism at blood lipid loci through multi-ancestry meta-analysis. 2022. Zenodo. <https://doi.org/10.5281/zenodo.7130299>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.