



**Universiteit
Leiden**
The Netherlands

Evidence- and consensus-based guidelines for drug-drug interactions with anticancer drugs; a practical and universal tool for management

Leeuwen, R.W.F.V.; Comte, M. le; Reyners, A.K.L.; Tweel, A. van den; Vlijmen, B. van; Kwee, W.; ... ; Jansman, F.G.A.

Citation

Leeuwen, R. W. F. V., Comte, M. le, Reyners, A. K. L., Tweel, A. van den, Vlijmen, B. van, Kwee, W., ... Jansman, F. G. A. (2022). Evidence- and consensus-based guidelines for drug-drug interactions with anticancer drugs; a practical and universal tool for management. *Seminars In Oncology*, 49(2), 119-129. doi:10.1053/j.seminoncol.2022.03.002

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3458849>

Note: To cite this publication please use the final published version (if applicable).



Evidence- and consensus-based guidelines for drug-drug interactions with anticancer drugs; A practical and universal tool for management

Roelof W.F. van Leeuwen¹, Marianne le Comte², Anna K.L. Reyners³, Annemieke van den Tweel⁴, Bas van Vlijmen⁵, Wilma Kwee², Brigit Wensveen², Neeltje Steeghs⁶, Otto Visser⁷, Teun van Gelder⁸, Frank G.A. Jansman^{9,10,*}

¹ Departments of Pharmacy and Medical Oncology, Erasmus University Medical Center, Rotterdam, the Netherlands

² Drug Information Center, Royal Dutch Pharmacists Association KNMP, the Hague, the Netherlands

³ Department of Medical Oncology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

⁴ Department of Clinical Pharmacy, Jeroen Bosch Hospital, 's Hertogenbosch, the Netherlands

⁵ Department of Pharmacy, Radboud University Medical Center, Nijmegen, the Netherlands

⁶ Department of Clinical Pharmacology, Division of Medical Oncology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

⁷ Department of Hematology, Isala Oncology Center, Zwolle, The Netherlands

⁸ Department of Clinical Pharmacy and Toxicology, Leiden University Medical Center, Leiden, the Netherlands

⁹ Department of Clinical Pharmacy, Deventer Teaching Hospital, Deventer, the Netherlands

¹⁰ Unit of Pharmacotherapy, Epidemiology & Economics, Groningen Research Institute of Pharmacy (GRIP), University of Groningen, Groningen, the Netherlands

ARTICLE INFO

Article history:

Received 23 July 2021

Revised 11 March 2022

Accepted 11 March 2022

Keywords:

Drug-drug Interactions

Herbals

Chemotherapy

Cancers

Practice

ABSTRACT

Drug-drug interactions (DDIs) with anticancer drugs are common and can significantly affect efficacy and toxicity of treatment. Therefore, a Dutch Multidisciplinary Expert group is assessing the clinical significance of DDIs in oncology and provides recommendations for the management of these DDIs. We present an overview of methodology and outcome of an evidence- and consensus-based assessment of DDIs between anticancer drugs and non-anticancer drugs.

A literature search was performed through PubMed and EMA and FDA assessment reports, to identify potential DDI's involving anticancer drugs. For each potential DDI a concept report for risk analysis and practical advice for management was created. Subsequently, this risk analysis and the corresponding advice were assessed and weighed.

A total of 290 potential DDIs have been identified in the literature thus far. Of these 290 potential DDIs, the Expert Group has identified 94 (32%) DDIs as clinically relevant, with a need for an automated alert and a suggested intervention. Furthermore, 110 DDIs have been identified as clinically not relevant. For 86 potential DDIs evidence supporting a relevant DDI was insufficient and in these cases neither an alert nor advice regarding a suggested intervention were formulated.

A transparent risk analysis is presented for identification of clinically relevant DDIs with anticancer drugs. Integration of DDI guidelines into the national electronic prescribing system is essential to achieve optimal efficacy and minimal toxicity in patients receiving anticancer therapy. A clear overview of clinically relevant DDIs with anticancer therapy provides clinicians with a structured, evidence-based and consensus-built tool for anticancer therapy surveillance.

© 2022 The Author(s). Published by Elsevier Inc.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Introduction

Over the past decades the incidence of cancer has rapidly increased making cancer the leading cause of death worldwide [1].

Cancer occurs primarily in older age groups, with over two third of the patients with cancer 65 years or older at the time of diagnosis [2]. Furthermore, comorbidities and polypharmacy defined as the use of ≥ 5 medications concomitantly, are common in elderly patients [3,4]. As a consequence, patients with cancer are at significant higher risk for drug related problems, such as drug-drug interactions (DDIs) [5,6]. Additionally, many anticancer drugs are potent and have narrow therapeutic windows with the result that (minor) changes in pharmacodynamic or pharmacokinetic

* Corresponding author. Deventer Teaching Hospital, Department of Clinical Pharmacy, Nico Bolkesteinaan 75, 7416SE, Deventer, the Netherlands. Tel.: 570-536-000; Fax.: 570-501-425

E-mail address: f.g.a.jansman@rug.nl (F.G.A. Jansman).

parameters caused by DDIs may profoundly affect efficacy or toxicity. In this case, the anticancer drug is regarded as a victim. At the same time, an anticancer drug can act as the causative agent, that is, perpetrator, and compromise the efficacy or toxicity of other concomitantly used drugs.

Until a decade ago, cancer patients were predominately treated in a hospital setting with intravenously administered anticancer agents. Although anticancer therapy often involves highly potent and potentially toxic agents, medication surveillance on DDIs with anticancer agents was usually not standard of care. However, with the increased life expectancy and shifting paradigm towards personalized medicine, the need for monitoring and management of DDIs by healthcare professionals to optimize anticancer therapy became more apparent [5,6]. Furthermore, many novel orally administered anticancer drugs, often administered for prolonged periods of time have been introduced in the past decades (eg, tyrosine kinase inhibitors) [7]. Although in many cases these novel oral agents have significantly improved life expectancy, patient comfort and quality of life, many new challenges have emerged since these drugs are highly prone to DDIs that include amongst others impact on absorption or metabolism by cytochrome P450 (CYP)-enzymes as well as additional effects on QT_c-interval [8,9].

To obtain a solid medication surveillance in (hemo)oncology a novel approach was chosen by our Multidisciplinary Expert Group consisting of (hospital)pharmacists, medical oncologists, hematologists, internists and clinical pharmacologists, originally constituted in 2006 [10]. The Expert Group identified potential DDIs between anticancer drugs and non-anticancer drugs, assessed their clinical significance and provided practical recommendations for the management of these DDIs. The results of these assessments and the practical recommendation for the management of these DDIs were subsequently made available for healthcare professionals through integration into the national computerized medication surveillance system. In this system, alerts regarding DDIs pop-up during the prescribing process. The aim was to improve medication safety and optimize the efficacy-toxicity balance during anticancer therapy.

The first results of the Multidisciplinary Expert Group were published in 2011 [10]. This article represents an updated overview of the assessed DDIs, including novel insights in the evidence-based and consensus-based practical guidelines on the management of DDIs involving anticancer drugs. This article may function as a practical, universal tool to further optimize medication surveillance of DDIs in cancer care.

Methods

The Multidisciplinary Expert Group was established by the Royal Dutch Pharmacists Association and meets on a regular basis. The members represent their scientific organizations, and include the Dutch Association of Hospital Pharmacists, the Royal Dutch Pharmacists Association, the Dutch Society for Medical Oncology, the Haemato Oncology Foundation for Adults in the Netherlands, and the Dutch Society for Clinical Pharmacology & Biopharmacy. Data on potential DDIs were primarily derived from a database search in PubMed with the following RSS-Feed: ((drug-drug interaction[MeSH Terms] AND ((Clinical Trial[ptyp] OR Case Reports[ptyp] OR Controlled Clinical Trial[ptyp] OR Letter[ptyp] OR Multicenter Study[ptyp]) AND Humans[Mesh] AND English[lang]) AND ((Clinical Trial[ptyp] OR Case Reports[ptyp] OR Controlled Clinical Trial[ptyp] OR Letter[ptyp] OR Multicenter Study[ptyp]) AND Humans[Mesh] AND English[lang])) until January 1st 2022, and assessment reports provided by the EMA and FDA registration authorities [11,12]. The search was limited to DDIs between anticancer drugs and non-anticancer drugs. This only concerns authorized drugs since these are included in the electronic medication surveillance system. In addition, some Over-The-Counter

Table 1
Categories for quality of scientific evidence for DDIs¹⁵

Level	Quality of Scientific Evidence
-	No evidence
0	Pharmacodynamic animal studies; in vitro studies with a limited predictive value for the human in vivo situation; data on file
1	Incomplete, published case reports
2	Well-documented, published case reports; retrospective analyses of case series; case control studies
3	Controlled, published interaction studies in patients or healthy volunteers, surrogate end points.
4	Controlled, published interaction studies in patients or healthy volunteers, clinically relevant end points

Table 2
Categories for clinical effects of DDI with example¹⁵

Severity level and examples of clinical effects per category
A: Clinically irrelevant effect Increase or decrease in drug level without direct clinical consequences: tyrosine kinase inhibitors, endoxifen (active metabolite of tamoxifen)
B: Temporarily adverse effect Decrease simvastatin/zolpidem level (by induction enzalutamide)
C: Longer-lasting adverse effect Increase of everolimus level (by CYP3A4-inhibitors) Increase of phenytoin plasma concentration (by capecitabine/5FU)
D: Long-lasting or permanent adverse effect Increased toxicity capecitabine/5FU (by folic acid/metronidazole) Decrease of plasma concentration of carbamazepine/phenytoin/valproic acid by certain anticancer agents
E: Severe adverse effect: Increased toxicity mercaptopurine (by allopurinol/febuxostat) Neuromuscular toxicity vinblastine/vincristine (by CYP3A4-inhibitors) Hepatic veno-occlusive disease by busulfan (by itraconazole/ketoconazole)
F: Potentially fatal effect Multi-organ failure (by combination of busulfan and metronidazole) Death (by combination of methotrexate and trimethoprim or co-trimoxazole)

(OTC)- and herbal drugs often used by patients on their own initiative that may jeopardize effective and safe anticancer therapy were considered for drug-drug interacting potential. For instance, St. John's Wort (hypericum) can significantly lower irinotecan exposure [13]. Therefore, some of the OTC- and herbal drugs have been included in the structural assessment of DDIs, for instance hypericum as an inducer of drug metabolizing enzymes, and supplements with calcium or magnesium for potential effects on drug absorption. Special attention was given to the extrapolation of a certain DDI to other drugs (also called the “group effect”). An example concerns the DDIs with CYP3A4 inhibitors and inducers. When an anticancer drug shows a DDI with a certain strong CYP3A4 inhibitor (eg, ketoconazole), this DDI generally also applies to other strong CYP3A4 inhibitors (eg, other azoles) [14]. As a consequence, the DDI-information and advice for management is extrapolated to these other CYP3A4 inhibitors. However, this extrapolation is more complicated when moderate CYP3A4 inhibitors (eg, fluconazole) are involved since data on DDIs with moderate CYP3A4 inhibitors are often not available. In such cases a thorough assessment by the Expert Group is needed. DDIs between anticancer drugs were not considered since anticancer drugs are frequently prescribed deliberately in combination in accordance with current guidelines. All data on potential DDIs were evaluated and categorized, with regard to quality, level of evidence and clinical significance as presented in Table 1 and Table 2 [15]. For each potential DDI a standardized data sheet for risk analysis and concise advice was predefined by a pharmacist and presented to the Multidisciplinary Expert Group. Subsequently, the Expert Group assessed, weighed and (if needed) adjusted the provided risk analyses to assess a final advice for management of the potential DDIs. After assessment, the results were grouped into 3 categories (Table 3). Thereafter, the risk analysis and the results of the as-

Table 3
Categories for potential DDIs and advice for managing DDIs

Category	Advice
Drug-drug interaction (DDI) has been established, and the effect is clinically relevant.	Intervention is required, alert is generated
DDI has been established, but the effect is not clinically relevant	No intervention is required, no alert is generated
DDI has not been established	No intervention is required, no alert is generated

assessments are made available for healthcare professionals by integration into the national drug database for clinical decision support and is incorporated in all national electronic prescribing systems alerting medical doctors and pharmacists on clinically relevant DDIs. To optimize anticancer therapy, a clear and practical advice on the management of DDIs was given and, by this, solid medication surveillance for DDIs during prescribing of anticancer agents was guaranteed. The workflow as described above is summarized in Fig. 1, in which DDIs are distinguished between (1) 2 single drugs, (2) between 2 groups of drugs or (3) between a single drug and a group of drugs.

Results

A total of 290 potential DDIs were identified in the literature and presented to the Expert Group. By assessment, 94 DDIs (32%) were considered as clinically relevant and a practical advice for management was given (eg, dose modification, discontinuation of treatment or additional monitoring). Table 4 Among these, 12 DDIs with OTC- and herbal drugs, that is, antacids, folic acid and hypericum as a CYP3A4-inducer, were identified and assessed as clinically relevant. In contrast, 110 potential DDIs (38%) were identified as not clinically relevant, and consequently not requiring an alert nor advice regarding an intervention. These DDIs mostly concerned minor alterations in pharmacokinetics and/or additional toxicities. For the remaining 86 potential DDIs (39%) that were mentioned in literature, the Expert Group considered the evidence for a DDI

insufficient and not relevant, so neither alert nor advice was generated.


The clinically relevant interactions requiring interventions mainly concern pharmaceutical interactions regarding decreased drug absorption in from the gastrointestinal tract, pharmacokinetic interactions with CYP450 metabolizing enzymes influencing efficacy or toxicity, and pharmacodynamic interactions leading to additional toxicities. Furthermore, several DDIs with specific underlying mechanisms were established and assessed as in need interventions. All results of the structured assessment of DDIs with anticancer drugs are presented in Supplementary Tables 5, 6, 7.

Discussion

Since the start of the Multidisciplinary Expert Group, 290 potential DDIs have been identified with almost half assessed as clinically relevant. The DDI assessments and management guidelines are integrated in all national electronic prescribing systems, thereby providing a practical tool for clinicians and (hospital)pharmacists for the management of DDIs involving anticancer therapies. More importantly, solid medication surveillance on DDIs is facilitated and guaranteed and patient safety improved.

With the increasing complexity of anticancer therapy and aging of the population, an adequate medication surveillance of DDIs in this population is increasingly important. Multiple studies show that DDIs are often not recognized as such by prescribers and that the prevalence of (clinically relevant) DDIs in cancer patients is high [5,16]. Although most DDIs are mild, some can be severe or even life threatening and an intervention is needed [10,16,17]. This can also apply to OTC- an herbal drugs, for which more research is warranted into their effects on the efficacy and toxicity of anticancer drugs.

The assessment of DDIs in anticancer therapy is an ongoing process since new insights in (the management of) DDIs are continuously presented in literature. Therefore, new information is continuously discussed and reassessed by the Expert Group and may lead to updates in the recommendations for management of a DDI. For instance, regarding the management of the DDI between tyrosine kinase inhibitors (TKIs) and proton pump inhibitors (PPIs), the use of the acidic beverage cola was added to the advice [18].



*DDI = Drug Drug Interaction

Fig. 1. Workflow Multidisciplinary Expert Group

Table 4

Clinically relevant drug-drug interactions, intervention required [See bottom of table for abbreviations and lists of drugs]

Anticancer agent	Interacting agent	Potential Effect	Management
- Abemaciclib - Palbociclib - Ribociclib	CYP3A4-inducers	Therapeutic failure of abemaciclib, palbociclib or ribociclib	- Avoid combination, or - Monitor effect of abemaciclib, palbociclib or ribociclib
- Abemaciclib - Palbociclib - Ribociclib	CYP3A4-inhibitors	↑ toxicity of abemaciclib, palbociclib or ribociclib	- Avoid combination, or ↓ abemaciclib, palbociclib or ribociclib dose
Abiraterone	CYP3A4-inducers	↓ serum level of abiraterone	- Use alternative for the CYP3A4-inducer, or ↑ abiraterone dose
Afatinib	- Ritonavir - Lopinavir - Cobicistat	↑ serum level of afatinib	Administer afatinib at least 1 hour before administration of HIV-medication
- Anagrelide - Arsenic trioxide - Oxaliplatin - Vandetanib	QTDrug List	↑ QTc interval	Avoid combination, or monitor ECG
Anthracyclines	Cyclosporine	↑ anthracycline serum levels	Avoid combination, or monitor serum level of anthracycline
Anti-androgens	Androgens (androstanolone, nandrolone, prasterone, testosterone)	Counteracting effect	Avoid combination
- Anti-estrogens - Aromatase inhibitors	Estrogens	Counteracting effect	Avoid combination
- Azathioprine - Mercaptopurine - Thioguanine	- Allopurinol - Febuxostat	Myelosuppression	- Use alternative for allopurinol or febuxostat or - Reduce mercaptopurine/azathioprine dose - Monitor hematological parameters and liver function
- Azathioprine - Mercaptopurine - Thioguanine	Ribavirin	Reversible myelosuppression	Monitor adverse effects (myelosuppression and pancytopenia) and hematological parameters
Bendamustine	Allopurinol	↑ toxicity (Stevens-Johnson syndrome and toxic epidermal necrolysis)	Avoid allopurinol, or consider rasburicase in tumor lysis syndrome/rapid tumor lysis
Bleomycin	Oxygen	↑ bleomycin lung toxicity	Inform the anesthesiologist about current or past bleomycin administration
Busulfan	- Itraconazole - Ketoconazole	↑ busulfan serum level	- Avoid combination - Interrupt itraconazole or ketoconazole
Busulfan	Metronidazole	↑ busulfan serum level	- Avoid combination - Interrupt metronidazole
- Capecitabine - Fluorouracil	Folic acid (5 mg)	Capecitabine or fluorouracil toxicity	Avoid concomitant folic acid
- Capecitabine - Fluorouracil - Tegafur	Phenytoin	↑ serum level of phenytoin	- Avoid combination - Monitor phenytoin serum level
- Capecitabine - Fluorouracil - Tegafur	Metronidazole	Capecitabine or fluorouracil or tegafur toxicity	- Avoid combination - Interrupt capecitabine, fluorouracil or tegafur - Use alternative for metronidazole
Certain cytostatic agents (bleomycin, carboplatin, carmustine, cisplatin, cyclophosphamide, dacarbazine, doxorubicin, mercaptopurine, vinblastine, vincristine)	Phenytoin	Therapeutic failure of phenytoin	Monitor serum level of phenytoin, adjust dose accordingly

(continued on next page)

Table 4 (continued)

Anticancer agent	Interacting agent	Potential Effect	Management
Certain cytostatic agents (bleomycin, cisplatin, cyclophosphamide, cytarabine, doxorubicin, etoposide, ifosfamide, methotrexate, paclitaxel)	Valproic acid	Therapeutic failure of valproic acid	Monitor serum level of valproic acid, adjust dose accordingly
Certain cytostatic agents (cisplatin, cyclophosphamide, cytarabine, daunorubicin, doxorubicin, hydroxycarbamide, thioguanine, vincristine)	Carbamazepine	Therapeutic failure of carbamazepine	Monitor serum level of carbamazepine, adjust dose accordingly
Cisplatin	Aminoglycosides/ amphotericin-B	Nephrotoxicity	Monitor adverse effects (nephrotoxicity)
Cisplatin	Loop diuretics	Nephrotoxicity	- Avoid administration of loop diuretics up to a month after discontinuation of cisplatin. - Short term loop diuretic use during cisplatin infusion is allowed.
Cladribine	- Lamivudine - Emtricitabine	Therapeutic failure of cladribine	Avoid lamivudine or emtricitabine use
Cyclophosphamide	Cyclosporine	↑ cyclosporine serum level	Monitor serum level of cyclosporine
Cytostatic agents	VKA's	Stronger fluctuation of coagulation time	- Monitor INR - Alert anticoagulation clinic
Dabrafenib	Gemfibrozil	↑ dabrafenib serum level	Avoid gemfibrozil
- Dabrafenib - Lorlatinib - Apalutamide	Midazolam	↓ midazolam serum level	Use alternative for midazolam (temazepam, flurazepam or oxazepam)
Dacomitinib	Dextromethorphan	↑ dextromethorphan serum level	Avoid combination
Darolutamide	CYP3A4-inducers	↓ darolutamide serum level	Avoid combination
- Dexamethasone (≥ 5 mg/day) - Methylprednisolone			
(≥ 100 mg/day)	Voriconazole	↓ voriconazole serum level	- Avoid combination, or - Monitor for symptoms of therapeutic failure of voriconazole and monitor voriconazole Cmin
Doxorubicin	HIV protease inhibitors	Doxorubicin toxicity	Monitor doxorubicin toxicity
- Duvelisib - Idelalisib - Olaparib	CYP3A4-inducers	↓ serum level of duvelisib, idelalisib or olaparib	- Avoid combination, or - Monitor effect of duvelisib, idelalisib or olaparib
Duvelisib	CYP3A4-inhibitors	↑ duvelisib serum level	- Consider dose reduction of duvelisib - Monitor for symptoms of toxicity (e.g. infections, diarrhea, colitis)
- Duvelisib - Fedratinib - Idelalisib - Ribociclib	Midazolam	↑ midazolam serum level	- Use alternative for midazolam, or - Monitor adverse effects (↑ sedation)
Enzalutamide			
Mitotane	Non-anticancer CYP3A4 substrates: Antipsychotics, antiarrhythmic drugs, DOACs, contraceptives, HCV drugs, HIV drugs, immunosuppressants, opioids, other drugs	↓ serum level of non anticancer CYP3A4 substrate	- Avoid combination or - Monitor effect of the non anticancer CYP3A4 substrate and adjust dose accordingly

(continued on next page)

Table 4 (continued)

Anticancer agent	Interacting agent	Potential Effect	Management
Enzalutamide	Gemfibrozil	↑ enzalutamide serum level	- Avoid gemfibrozil, or - Consider ↓ enzalutamide dose
Enzalutamide			
Apalutamide	VKA's	↓ effect of VKA	Alert anticoagulation clinic
- Enzalutamide - Apalutamide	- Omeprazole - Esomeprazole	↓ (es)omeprazole effect	↑ (es)omeprazole dose
Etoposide	Cyclosporine	↑ etoposide serum level	↓ etoposide dose
Everolimus	Cyclosporine	↑ everolimus serum level	- Avoid combination, or - Consider ↓ everolimus dose
Everolimus	- CYP3A4-inducers - Dabrafenib	↓ serum level of everolimus	- Avoid combination, or - Consider ↑ everolimus dose
Everolimus	- CYP3A4-inhibitors - Fluconazole - Imatinib - Verapamil	↑ everolimus serum level	- Avoid combination, or - Consider ↓ everolimus dose - No action needed if using 150 mg single dose or 150 mg once a week fluconazole
Everolimus	Flucloxacilline	↓ everolimus serum level	Avoid combination, or monitor everolimus
Ibrutinib	CYP3A4-inhibitors	↑ ibrutinib serum level	- Avoid combination (ketoconazole) or - Consider decreasing dose ibrutinib (other CYP3A4-inhibitors)
- Ibrutinib - Acalabrutinib	- DOACs - TAls - Heparins - NSAID's - SSRIs	↑ bleeding tendency	Consultation with hematologist
- Ibrutinib - Acalabrutinib	VKA's	↑ bleeding tendency, without affecting INR	Alert anticoagulation clinic
- Imatinib - Ribociclib	Simvastatin	↑ statin serum level	- Imatinib: Use alternative for simvastatin - Ribociclib: Stop simvastatin (patient is in palliative phase)
Imatinib	Cyclosporine	↑ cyclosporine serum level	Monitor serum level of cyclosporine and renal function
Immunomodulatory monoclonal antibodies	Live attenuated vaccines	Generalized infection or ↓ effect of vaccine	Avoid combination
- Immuno-suppressant oncolytics - TKIs (immunosuppressive) - Monoclonal antibodies (immunosuppressant)	Inactivated vaccines	↓ effect of vaccine	Consider repeated administration
- Immuno-suppressant oncolytics - TKIs (immunosuppressive) - Monoclonal antibodies (immunosuppressant)	Live attenuated vaccines	Generalized infection or ↓ effect of vaccine	Avoid combination
Irinotecan	CYP3A4-inhibitors	↑ serum level of active irinotecan metabolite, SN-38	- Use alternative for CYP3A4-inhibitor, or - Monitor AUC of SN-38
Irinotecan	- Enzalutamide - Mitotane	↓ serum level of irinotecan and active irinotecan metabolite, SN-38	- Use alternative for CYP3A4-inhibitor, or - Consider dose increase for irinotecan
- Ixazomib - Sonidegib	Rifampicin	↓ serum level of ixazomib or sonidegib	- Avoid combination or - Monitor effect of ixazomib or sonidegib
- Lapatinib - Vemurafenib	Digoxin	↑ digoxin serum level of	Monitor serum level of digoxin and adverse effects

(continued on next page)

Table 4 (continued)

Anticancer agent	Interacting agent	Potential Effect	Management
• Methotrexate • Etoposide • Teniposide	• Carbamazepine • Phenytoin • Phenobarbital	• ↓ serum levels of carbamazepine, phenytoin, or phenobarbital	• Monitor serum level of antiepileptic drug
• Methotrexate	• Cotrimoxazole • Trimethoprim	• Methotrexate toxicity	• Avoid combination, except in acute lymphoblastic leukaemia (ALL) • Prophylaxis with low dose cotrimoxazole
• Methotrexate	• Cyclosporine	• ↑ serum level of both methotrexate and cyclosporine	• Monitor adverse effects of methotrexate • Monitor serum levels of methotrexate, and cyclosporine, hematological parameters, and renal function.
• Methotrexate	• Immunocyanin	• ↓ immunocyanin effect	• Avoid combination
• Methotrexate	• NSAID's	• ↑ methotrexate serum level	• Avoid combination, or • Monitor methotrexate serum level and adverse effects and renal function
• Methotrexate	• Probenecid	• ↑ methotrexate serum level	• Avoid combination, or • Monitor adverse effects of methotrexate and monitor renal function
• Methotrexate	• Tenofovir disoproxil	• ↑ risk of tenofovir disoproxil (nephro)toxicity	• Monitor renal function • Consider replacing tenofovir disoproxil by tenofovir alafenamide as part of combination antiretroviral therapy
• Methotrexate, high dose	• Levetiracetam	• ↑ methotrexate serum level	• Monitor adverse effects of methotrexate and renal function
• Methotrexate, high dose	• PPI's	• ↑ methotrexate serum level	• Interrupt PPI use
• Methotrexate, high dose	• Voriconazole	• ↑ risk of phototoxicity	• Monitor for symptoms of phototoxicity • Instruct patient to avoid sun exposure
• Mitotane	• Spironolactone	• ↓ mitotane effect	• Avoid combination, or • Monitor serum level of mitotane
• Olaparib	• CYP3A4-inhibitors	• Olaparib toxicity	• Avoid combination, or • ↓ olaparib dose
• Panobinostat	• CYP3A4-inducers	• ↓ panobinostat serum level	• Use alternative for panobinostat, or • Monitor effect of panobinostat
• Panobinostat	• CYP3A4-inhibitors	• ↑ panobinostat serum level	• Use alternative for panobinostat, or • Consider decreasing panobinostat dose
• Regorafenib • Darolutamide	• Rosuvastatin	• Rosuvastatin toxicity (myopathy)	• Instruct patient on possible symptoms of rosuvastatin toxicity
• Regorafenib • Vandetanib	• Rifampicin	• Alteration of regorafenib or vandetanib effect	• Use alternative for regorafenib or vandetanib, or • Monitor effect regorafenib or vandetanib
• Rolapitant	• Rifampicin	• ↓ rolapitant serum level	• Monitor effect of rolapitant
• Ruxolitinib	• Fluconazole	• ↑ ruxolitinib serum level	• Avoid combination or • Consider reduction of ruxolitinib dose • No action needed with fluconazole dosages of 150 mg single dose or 150 mg once a week

(continued on next page)

Table 4 (continued)

Anticancer agent	Interacting agent	Potential Effect	Management
• Talazoparib	• Itraconazole	• ↑ talazoparib toxicity	• Avoid combination or • Consider dose reduction of talazoparib
• Tamoxifen	• VKA's	• ↑ effect of VKA	• Monitor INR • Alert anticoagulation clinic
• Tamoxifen	• CYP2D6-inhibitors	• ↓ formation of active endoxifen metabolite	• Avoid combination
• Tamoxifen	• CYP3A4-inducers	• ↓ serum level of tamoxifen and active metabolite endoxifen	• Avoid combination, or • Monitor serum level of tamoxifen and endoxifen and consider dose-adjustment
• Tamoxifen	• Hydroxychloroquine • Chloroquine	• ↑ tamoxifen toxicity (irreversible retinopathy)	• Avoid combination or • Monitor toxicity (retinopathy)
• Temsirolimus	• CYP3A4-inducers	• ↓ (tem)sirolimus serum level	• Use alternative for CYP3A4-inducer, or • Consider increasing temsirolimus dose
• Temsirolimus	• CYP3A4-inhibitors	• ↑ (tem)sirolimus serum level	• Use alternative for CYP3A4-inhibitor, or • Consider reducing temsirolimus dose
• Temsirolimus	• Disulfiram • Metronidazole	• Torisel© solution for injection contains alcohol, resulting in ↑ alcohol toxicity when combined with metronidazole or disulfiram	• Avoid combination
• Tivozanib • Vemurafenib	• Rifampicin	• ↓ serum level of tivozanib or vemurafenib	• Avoid combination, or • Monitor effects of tivozanib or vemurafenib
• Trabectedin	• CYP3A4-inhibitors	• ↑ trabectedin serum level	• Avoid combination, or • Monitor adverse effects
• Trabectedin	• Rifampicin	• ↓ trabectedin serum level	• Avoid combination
• TKIs (all except acalabrutinib, ibrutinib (separate DDI))	• VKA's	• Stronger fluctuation of coagulation time	• Monitor INR • Alert the anticoagulation clinic
• TKIs (acalabrutinib, afatinib, avapritinib, axitinib, bosutinib, brigatinib, cabozantinib, ceritinib, cobimetinib, crizotinib, dabrafenib, dasatinib, entrectinib, erlotinib, gefitinib, gilteritinib, ibrutinib, imatinib, lapatinib, larotrectinib, lorlatinib, midostaurine, neratinib, nilotinib, nintedanib, osimertinib, pazopanib, ponatinib, ruxolitinib, sorafenib, sunitinib)	• CYP3A4-inducers	• ↓ TKI serum level	• Avoid combination, or • Monitor effect of TKI, or • ↑ dose of TKI • Therapeutic drug monitoring of certain TKIs is an option
• TKIs (acalabrutinib, avapritinib, axitinib, bosutinib, brigatinib, ceritinib, cobimetinib, crizotinib, dabrafenib, dasatinib, encorafenib, entrectinib, erlotinib, fedratinib, gefitinib, gilteritinib, lapatinib, larotrectinib, lorlatinib, midostaurine, neratinib, nilotinib, pazopanib, regorafenib, ruxolitinib, sunitinib)	• CYP3A4-inhibitors	• ↑ TKI serum level	• Avoid combination, or • Reduce dose of tyrosine kinase inhibitor • Therapeutic drug monitoring of certain TKIs is an option.
• TKIs, various (acalabrutinib, bosutinib, ceritinib, dacomitinib, dasatinib, erlotinib, gefitinib, lapatinib, neratinib, pazopanib)	• Antacids	• ↓ absorption of TKI	• Separate the dose: TKI at least 2 hours before or 4 hours after the antacid
	• PPIs • H2-receptor antagonists	• ↓ availability of TKI	• Consider temporarily stop of the PPI or H2-recept antagonist, or separate the dose: TKI, 2 hours before PPI or H2-receptor antagonist; if this is not possible, TKI directly followed by PPI or H2-recept antagonist, or TKI concomitantly with Coca-Cola

(continued on next page)

Table 4 (continued)

Anticancer agent	Interacting agent	Potential Effect	Management
• Vandetanib	• Metformin	• ↑ metformin serum level	• Adjust metformin dose • Instruct patient on possible symptoms
• Venetoclax	• CYP3A4-inducers	• Therapeutic failure of venetoclax	• Avoid combination
• Venetoclax	• CYP3A4-inhibitors • Diltiazem • Fluconazole • Verapamil	• ↑ venetoclax serum level	• Avoid combination or decrease venetoclax dose • No action needed with fluconazole dosages of 150 mg single dose or 150 mg once a week
• Vinblastine	• CYP3A4-inhibitors	• Vinblastine (neuro)toxicity	• Avoid combination, or • Monitor toxicity of vinblastine
• Vincristine	• CYP3A4-inhibitors	• Vincristine (neuro)toxicity	• Avoid combination, or • Monitor toxicity of vincristine • No action needed with fluconazole dosages of 150 mg single dose or 150 mg once a week

Abbreviations: DOAC, direct oral anticoagulant; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitor; SSRI, selective serotonin reuptake inhibitor; TAI, thrombocyte aggregation inhibitor; TKI, tyrosine kinase inhibitor; VKA, vitamin K antagonist

Anthracyclines: Daunorubicin, doxorubicin, epirubicin, idarubicin, mitoxantrone, pixantrone

Anti-androgens: Abiraterone, apalutamide, bicalutamide, darolutamide, enzalutamide, nilutamide

Aromatase inhibitors: Anastrozole, exemestane, letrozole

CDK4/6 inhibitors: Abemaciclib, palbociclib, ribociclib

CYP3A4-inducers: Carbamazepine, efavirenz, enzalutamide, phenobarbital, phenytoin, hypericum, mitotane, nevirapine, primidone, rifabutin, rifampicin

CYP3A4-inhibitors: Clarithromycin, cobicistat, erythromycin, itraconazole, ketoconazole, posaconazole, voriconazole, voriconazole, ritonavir

CYP2D6-inhibitors: Bupropion, cinalcacet, fluoxetine, quinidine, paroxetine, terbinafine

DOACs - Direct oral anticoagulants: Apixaban, dabigatran, edoxaban, rivaroxaban

HIV protease inhibitors: Atazanavir, darunavir, fosamprenavir, indinavir, lopinavir/ritonavir, nelfinavir, ritonavir, saquinavir, tipranavir

Histamine H2-receptor antagonists: Cimetidine, famotidine, nizatidine

Immunomodulatory monoclonal antibodies: Atezolizumab, avelumab, blinatumomab, cemiplimab, dinutuximab, dinutuximab beta, durvalumab, ipilimumab, nivolumab, pembrolizumab

Immunosuppressant oncolytics: Cytostatics, alpelisib, bortezomib, carfilzomib, CDK4/CDK6-inhibitors, duvelisib, ixazomib, idelalisib, panobinostat, PARP-inhibitors poma-lidomide, CAR-T-celtherapy, talimogene laherparepvec (T-VEC)

NSAIDs - Non-steroidal anti-inflammatory drugs: Celecoxib, diclofenac, etoricoxib, ibuprofen, indomethacin, mefenamic acid, naproxen

PPIs - Proton pump inhibitors: Dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole, omeprazole/sodium bicarbonate (Zegerid®)

QTDrug List: Amiodarone, azithromycin, chlorpromazine, chlorpromazine, chloroquine, (es)citalopram, claritromycin, disopyramide, donepezil, droperidol, erythromycin, flecainid, fluconazole, haloperidol, hydroxychloroquine, ibutilide, ketanserin, kinidine, levofloxacin, levomepromazine, methadone, moxifloxacin, ondansetron, papaverine, pentamidine, pimozide, procainamide, roxithromycin, sertindol, sotalol, sulphuride

Source: Arizona Center for Education and Research on Therapeutics (AZCERT): QTDrug List – Drugs with potential to prolong the QT interval and known risk of torsades de pointes (TdP)

SSRIs - Selective serotonin reuptake inhibitors:

• Citalopram, dapoxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline

• Duloxetine, milnacipram, venlafaxine (noradrenaline and serotonin reuptake inhibitors [SNRI])

• Trazodone (serotonin antagonist and reuptake inhibitor [SARI])

TAIs - Thrombocyte aggregation inhibitors: Acetylsalicylic acid, clopidogrel, dipyridamole, eptifibatide, prasugrel, ticagrelor, tirofiban

VKAs - Vit K antagonists: Acenocoumarol, phenprocoumon, warfarin

In addition, the extrapolation of the DDI between TKIs and PPIs must be considered carefully. Although many TKIs show a clinically relevant DDI with a PPI based on altered pH dependant solubility, it is unknown whether this DDI can be extrapolated to all TKIs with known pH dependant solubility. Therefore, a solid in-depth evaluation of known literature and a clear understanding of pharmacology is needed for the risk assessment of this DDI [19].

The past decade there has seen a shifting paradigm towards therapeutic drug monitoring (TDM) in managing DDIs. Especially for most TKIs, there is a clear relationship between exposure, toxicity and treatment efficacy and recommendations have been developed for the integration of TDM for this class of anticancer drugs [20–23]. For certain anticancer agents TDM offers a practical way to manage DDIs, where dose adjustments can be made if drug plasma levels are outside the therapeutic window. To explore further application of TDM, more research is needed to confirm the relevance of TDM as an instrument to be leveraged in the management of DDIs in oncology.

Current studies clearly show that DDIs frequently occur in cancer patients and are often clinically relevant [24–27]. The assess-

ment and advice given by the Multidisciplinary Expert Group contributes to an efficient and evidence-based electronic DDI medication surveillance, where these guidelines provide an important tool in the prescribing of anticancer drugs, thereby optimizing treatment efficacy and improving patient safety.

Conclusion

As the complexity of anticancer therapy increases, more specific screening tools for the detection of DDIs are necessary to increase the efficacy and limit the toxicity of anticancer therapy. This article gives a clear overview of clinically relevant DDIs with anticancer therapy. Furthermore, a straightforward approach for assessment and integration of DDI guidelines into the national electronic prescribing system is provided and may offer a structured, evidence-based, and consensus-based tool for DDI medication surveillance during anticancer therapy. The overview and specific universal tools given in this study may help (hemo-)oncologists and pharmacists to be more aware of DDIs during anticancer therapy and may lead to a closer collaboration in the assessment, management,

and integration of these DDIs into national electronic prescribing systems.

Study highlights

Drug-drug interactions (DDIs) with anticancer drugs are common and can significantly affect efficacy and toxicity of treatment. Since 2006, a Dutch Multidisciplinary Expert Group has assessed the clinical significance of DDIs in oncology and provides recommendations for the management of these DDIs.

A transparent methodology and outcomes of an evidence- and consensus-based assessment of DDIs concerning anticancer drugs is presented. Integration of DDI guidelines into the national electronic prescribing system is essential to achieve optimal efficacy and minimal toxicity in patients receiving anticancer therapy. A clear overview of clinically relevant DDIs with anticancer therapy provides clinicians with a structured, evidence-based, and consensus-built tool for anticancer therapy surveillance and management.

Author contributions

RvL: Formal analysis; Investigation; Methodology; Validation; Visualization; Writing – original draft. **MIC:** Data curation; Formal analysis; Methodology; Project administration; Software; Validation; Writing – review & editing. **AR:** Formal analysis; Methodology; Validation; Writing – review & editing. **AT:** Formal analysis; Methodology; Validation; Writing – review & editing. **BV:** Formal analysis; Methodology; Validation; Writing – review & editing. **WK:** Formal analysis; Methodology; Validation; Writing – review & editing. **BW:** Data curation; Formal analysis; Methodology; Project administration; Software; Validation; Visualization; Writing – review & editing. **NS:** Formal analysis; Methodology; Validation; Writing – review & editing. **OV:** Formal analysis; Methodology; Validation; Writing – review & editing. **TvG:** Formal analysis; Methodology; Validation; Writing – review & editing. **FI:** Conceptualization; Formal analysis; Funding acquisition; Investigation; Methodology; Supervision; Validation; Writing – original draft; Writing – review & editing.

Conflict of interest

RvL: Research (unrestricted) grants Roche, Bayer, Pfizer, Boehringer Ingelheim, Astellas, BMS. Consultancy: MSD, BMS, Sanofi, Astellas, Pfizer, Roche Travel grants: Roche, Novartis, Pfizer, Astellas NS: provided consultation or attended advisory boards for AIMM Therapeutics, Boehringer Ingelheim, Ellipses Pharma. N Steeghs received research grants for the institute from AB Science, Abbvie, Actuate Therapeutics, Amgen, Array, AstraZeneca/MedImmune, Bayer, Blueprint Medicines, Boehringer Ingelheim, Bristol-Myers Squibb, Cantargia, Cytovation, Deciphera, Genentech/Roche, GlaxoSmithKline, Incyte, InteRNA, Lilly, Merck Sharp & Dohme, Merus, Novartis, Pfizer, Pierre Fabre, Roche, Sanofi, Taiho, Takeda (outside the submitted work) TvG: In the last 3 years TvG has received lecture fees and study grants from Chiesi and Astellas, in addition to consulting fees from Roche Diagnostics, Vitaeris, CSL Behring, Astellas, Aurinia Pharma and Novartis. In all cases money has been transferred to hospital accounts, and none has been paid to his personal bank accounts. TvG does not have employment or stock ownership at any of these companies, and neither does he have patents or patent applications MIC, AR, AvdT, BvV, WK, BW, OV, and FJ report no conflict of interest or funding information

Acknowledgments

Dr Meta Diekstra, PharmD, and Florentine Hogenhuis, PharmD, are acknowledged for their administrative contribution to the manuscript.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1053/j.seminoncol.2022.03.002](https://doi.org/10.1053/j.seminoncol.2022.03.002).

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68(6):394–424.
- Signaleringscommissie Kanker KWF Kankerbestrijding. Kanker in Nederland tot 2020. Trends en prognoses (2011).
- Alkan A, Yaşar A, Karci E, Köksöy EB, Ürün M, Şenler FÇ, et al. Severe drug interactions and potentially inappropriate medication usage in elderly cancer patients. *Support Care Cancer* 2017;25:229–36.
- Sharma M, Loh KP, Nightingale G, Mohile SG, Holmes HM. Polypharmacy and potentially inappropriate medication use in geriatric oncology. *J Geriatr Oncol* 2016;7(5):346–53.
- Van Leeuwen RWF, Brundel DH, Neef C, van Gelder T, Mathijssen RH, Burger DM, et al. Prevalence of potential drug–drug interactions in cancer patients treated with oral anticancer drugs. *Br. J. Cancer* 2013;108(5):1071–1078.
- Riechelmann RP, Krzyzanowska MK. Drug interactions and oncological outcomes: a hidden adversary. *Ecancermedicalscience* 2019 Mar 28;13:ed88.
- Weingart, S.N. et al. JNCCN NCCN Task Force Report: Oral Chemotherapy.
- Leveque D, Delpeuch A, Gourieux B. New anticancer agents: Role of clinical pharmacy services. *Anticancer Res* 2014;34:1573–8.
- Van Leeuwen RWF, van Gelder T, Mathijssen J, R H, Jansman A, F G. Review Drug–drug interactions with tyrosine-kinase inhibitors: a clinical perspective. *Lancet Oncol* 2014;15:e315–26.
- Jansman FGA, Reyners AK, van Roon EN, Smorenburg CH, Helgason HH, le Comte M, et al. Consensus-Based Evaluation of Clinical Significance and Management of Anticancer Drug Interactions. *Clin Ther* 2011;33:305–14.
- European Medicines Agency, 2022 available on: <https://www.ema.europa.eu/en>.
- U.S. Food and Drug Administration, 2022 available on: <https://www.fda.gov>.
- Mathijssen RH, Verwij J, de Bruijn P, Loos WJ, Sparreboom A. Effects of St. John's wort on irinotecan metabolism. *J Natl Cancer Inst* 2002;94(12):1247–9.
- Flockhart DA, Oesterheld JR. Cytochrome P450-mediated drug interactions. *Child Adolesc Psychiatr Clin N Am* 2000;9(1):43–76.
- Van Roon EN, et al. Clinical relevance of drug–drug interactions: a structured assessment procedure. *Drug Saf* 2005;28:1131–9.
- Van Leeuwen RWF, et al. Drug–drug interactions in patients treated for cancer: a prospective study on clinical interventions. *Ann. Oncol.* 2015;26(5):992–997.
- The Royal Dutch Pharmacists Association (KNMP). Working group for pharmacotherapy and drug information. Decision map for interaction assessment (2015), available on: <https://www.knmp.nl/producten/gebruiksrecht-g-standaard/medicatiebewaking-g-standaard/interacties>.
- Van Leeuwen RWF, et al. Influence of the acidic beverage cola on the absorption of erlotinib in patients with non-small-cell lung cancer. *J Clin Oncol* 2016;34(12):1309–14.
- Van Leeuwen RWF, Jansman FGA, Hunfeld NG, Peric R, Reyners AKL, Imholz ALT, Brouwers JRB, Aerts JG, van Gelder T, Mathijssen RHJ. Tyrosine Kinase Inhibitors and Proton Pump Inhibitors: An Evaluation of Treatment Options. *Clin Pharmacokinet* 2017;56:683–8.
- IJzerman NS, Groenland SL, Koenen AM, Kerst M, van der Graaf WTA, Rosing H, Beijnen JH, Huitema ADR, Steeghs N. Therapeutic drug monitoring of imatinib in patients with gastrointestinal stromal tumours – Results from daily clinical practice. *Eur J Cancer* 2020;136:140–8 Sep.
- Westerdijk K, Desai IM, Steeghs N, van der Graaf WTA, van Erp NP. Dutch Pharmacology and Oncology Group (DPOG). Imatinib, sunitinib and pazopanib: from flat-fixed dosing towards a pharmacokinetically guided personalized dose. *Br J Clin Pharmacol* 2020;86(2):258–73 Feb.
- Groenland SL, van Eerden RAG, Verheijen RB, et al. Therapeutic drug monitoring of oral anticancer drugs: the Dutch pharmacology oncology group-therapeutic drug monitoring protocol for a prospective study. *Ther Drug Monit* 2019;41(5):561–7 Oct.
- Verheijen RB, Yu H, Schellens JHM, Beijnen JH, Steeghs N, Huitema ADR. Practical recommendations for therapeutic drug monitoring of kinase inhibitors in oncology. *Clin Pharmacol Ther* 2017;102(5):765–76 Nov.
- Bulsink A, Imholz ALT, Brouwers JRB, Jansman FGA. Characteristics of potential drug-related problems among oncology patients. *Int. J. Clin. Pharm.* 2013. doi:[10.1007/s11096-012-9747-7](https://doi.org/10.1007/s11096-012-9747-7).

- [25] Ko Y, et al. Prevalence of the coprescription of clinically important interacting drug combinations involving oral anticancer agents in singapore: a retrospective database study. *Clin Ther* 2012;34:1696–704.
- [26] Ramos-Esquivel A, Víquez-Jaikel Á, Fernández C. Potential Drug-Drug and Herb-Drug Interactions in Patients With Cancer: A Prospective Study of Medication Surveillance. *J. Oncol. Pract.* 2017;13:e613–22.
- [27] Van Loveren F, Van Berlo-Van de Laar IRF, Imholz ALT, Van 't Riet E, Taxis K, Jansman FGA. Prevalence and follow-up of Potentially Inappropriate Medication and Potentially Omitted Medication in older patients with cancer – the PIM-POM study. *J Geriatr Oncol* 2021;12:80–4.